



Clinical Microbiology

Performance of rapid tests for carbapenemase detection among Brazilian *Enterobacteriaceae* isolates



Lisiane Rech Pancotto^a, Carolina Silva Nodari^a, Franciéli Pedrotti Rozales^{a,b}, Tatiane Soldi^a, Carolina Gomes Siqueira^c, Ana Lúcia Freitas^c, Afonso Luís Barth^{a,b,c,*}

^a Hospital de Clínicas de Porto Alegre, Laboratório de Pesquisa em Resistência Bacteriana (LABRESIS), Porto Alegre, RS, Brazil

^b Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Ciências Farmacêuticas, Porto Alegre, RS, Brazil

^c Universidade Federal do Rio Grande do Sul, Faculdade de Farmácia, Porto Alegre, RS, Brazil

ARTICLE INFO

Article history:

Received 9 April 2017

Accepted 7 July 2018

Available online 16 August 2018

Associate Editor: Ana Lucia Darini

Keywords:

Carbapenemases

Rapid tests

Carba NP

Blue-Carba

CIM

ABSTRACT

The global emergence of carbapenemases led to the need of developing new methods for their rapid detection. The aim of this study was to evaluate the performance of the rapid tests for carbapenemase-producing and non-producing *Enterobacteriaceae*. Carbapenem non-susceptible *Enterobacteriaceae* from a surveillance study submitted to a multiplex real time PCR for carbapenemase detection were included in this study. The isolates were subjected to the rapid phenotypic tests Carba NP, Blue-Carba and Carbapenem Inactivation Method (CIM). A total of 83 carbapenemase-producing (43) and non-producing (40) isolates were included in the study. The sensitivity/specificity were 62.7%/97.5%, 95.3%/100%, and 74.4%/97.5% for Carba NP, Blue-Carba and CIM, respectively. Both Carba NP and Blue-Carba presented their final results after 75 min of incubation; the final results for CIM were obtained only after 8 h. Failure to detect OXA-370 carbapenemase was the main problem for Carba NP and CIM assays. As the Blue-Carba presented the highest sensitivity, it can be considered the best screening test. Conversely, CIM might be the easiest to perform, as it does not require special reagents. The early detection of carbapenemases aids to establish infection control measures and prevent carbapenemases to spread reducing the risk of healthcare associated infections and therapeutic failure.

© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Nowadays carbapenem resistance among *Enterobacteriaceae* is a major concern. These bacteria become resistant to this

antibiotic class mainly due to three mechanisms, either isolated or combined: efflux pumps; permeability loss; and carbapenemase production.¹ The latter represents a growing concern, since these enzymes hydrolyze not only carbapenems but also other β -lactams and their dissemination is usually plasmid-mediated.²

Carbapenemases can be classified as class A, B or D β -lactamases,³ based on their molecular structure. In Brazil, different carbapenemase classes have emerged over the years

* Corresponding author.

E-mail: albarth@hcpa.edu.br (A.L. Barth).

<https://doi.org/10.1016/j.bjm.2018.07.002>

1517-8382/© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

but the *Klebsiella pneumoniae* carbapenemase (KPC) is already considered endemic.² We have also observed the spread of the New Delhi Metallo-β-lactamase (NDM) in recent years^{4,5} and the emergence of OXA-370 in some regions.⁵ Moreover, scattered reports of GES-5 and IMP-1 have been described over the country.^{6,7}

The global emergence of carbapenemases brought to light the need of developing new methods for its rapid detection. The enzymatic activity of carbapenemases is explored as an alternative for their detection in some recent studies. This activity might be detected by biochemical assays such as Carba NP and Blue-Carba methods,^{8,9} in which the hydrolysis of the β-lactam ring is visualized by the colour change of a pH indicator, or using mass spectroscopy in which the absence of the carbapenem molecular ion is observed in the presence of a carbapenemase-producing bacteria.^{10,11} Van der Zwaluw et al. also proposed a methodology (the Carbapenem Inactivation Method – CIM) in which a carbapenem-susceptible *E. coli* strain is able to grow near a meropenem disk after its incubation with a carbapenemase-producing strain.¹² Recently, the Clinical & Laboratory Standards Institute (CLSI) proposed the use of Carba NP and the CIM assays as screening methods for carbapenemase detection.¹³

The aim of this study was to evaluate the performance of the rapid tests Carba NP, Blue-Carba and CIM against carbapenemase-producing and -non-producing Enterobacteriaceae.

Material and methods

Bacterial isolates

Carbapenem non-susceptible Enterobacteriaceae, from a previous surveillance study, were identified by biochemical characterization and, when necessary, confirmed using VITEK2 system (BioMérieux, France). Isolates resistant to, at least, imipenem and/or meropenem were selected for this study.

Genotypic detection of carbapenemases

Carbapenemase genes were detected by a multiplex real-time PCR with specific primers for the *bla*_{KPC}, *bla*_{GES}, *bla*_{IMP}, *bla*_{NDM}, *bla*_{VIM} and *bla*_{OXA-48-like}.¹⁴ PCR products of isolates positive for *bla*_{GES} and *bla*_{OXA-48-like} were purified using the ExoStar kit (GE Healthcare) and sequenced using a BigDye Terminator kit (version 3.1) and an ABI 3500 Genetic Analyzer (Applied Biosystems).

Phenotypic detection of carbapenemases

Carba NP

The carbapenemase detection by Carba NP was performed and interpreted as previously described by Nordmann et al.⁸ Briefly, overnight cultured colonies grown in Mueller Hinton agar were submitted to extraction with B-PERII (Bacterial Protein Extraction Reagent, Thermo Scientific Pierce, Rockford, USA) and incubated for 2 h in a solution containing phenol-red and imipenem. The bacterial extract was also incubated in a

phenol-red solution without antibiotic. The result was considered positive when the solution containing imipenem became orange or yellow. The result was not validated if the solution without antibiotic presented a colour change more expressive than the solution containing imipenem. A known KPC-2 producer was used as positive control, and *E. coli* ATCC 25922 was tested as negative control.

Blue-Carba

Carbapenemase detection by Blue-Carba was performed and interpreted as previously described by Pires et al.⁹ Briefly, overnight cultured colonies grown in Mueller Hinton agar were incubated for 2 h in a solution containing bromothymol blue and imipenem. The bacterial colonies were also incubated in a bromothymol blue solution without antibiotic. The result was considered positive when the solution containing imipenem became green or yellow, and its color was different from the one observed in the negative control. The result was not validated if the solution without antibiotic presented the same or stronger colour change as the solution containing imipenem. A known KPC-2 producer was utilized as positive control, and a test tube containing only bacteria inoculum and Blue-Carba solution was used as negative control for each isolate tested.

Carbapenemase inhibition method (CIM)

The CIM was performed as previously described by van der Zwaluw et al.¹² Briefly, overnight cultured colonies were incubated for 2 h with a meropenem disk in 400 μL of water, with zinc sulfate supplementation for metallo-beta-lactamase-producing isolates. After the incubation, the disk was placed on a Mueller-Hinton agar plate previously inoculated with a susceptible *E. coli* (ATCC 25922) and incubated at 35 °C up to the growth of the strain (around 6 h). The result was considered positive when the *E. coli* presented an inhibition zone ≤16 mm around the meropenem disk, according to the CLSI interpretation criteria.¹³ A known KPC-2 producer was utilized as positive control, and *E. coli* ATCC 25922 was tested as negative control.

Results

A total of 83 carbapenem-non-susceptible Enterobacteriaceae isolates were selected for this study: 12 KPC, 13 NDM, 12 OXA-370, four GES-5, and two IMP producers, as well as 40 carbapenemase non-producers. All isolates were submitted to the rapid tests Carba NP, Blue-Carba and CIM. The results for carbapenemase-producing isolates, as well as the two carbapenemase-non-producing isolates with at least one false positive result, are presented in Table 1. The 38 carbapenemase-non-producing isolates that were not included in the table presented negative results for all tests performed. The Carba NP test presented a sensitivity of 62.7% and a specificity of 97.5%. The vast majority of isolates with positive results presented a clear color change within 15 min of incubation; only one isolate required the maximum time of 75 min to become positive (Table 2). The sensitivity of Blue-Carba was 95.3% and its specificity was 100%. As observed in Carba NP, most isolates with positive results in the Blue-Carba presented the final result within 15 min of incubation; only

Table 1 – Results of rapid tests Carba NP, Blue-Carba and CIM for carbapenemase-producing and non-producing Enterobacteriaceae.

Isolate ID	Carbapenemase	Carba NP	Time to positivity (min)	Blue-Carba	Time to positivity (min)	CIM
1101F	KPC	–	NA	+	30	+
1345F	KPC	+	15	+	15	+
1373F	KPC	+	15	+	15	+
1388F	KPC	+	15	+	15	+
1389F	KPC	+	45	+	15	+
1390F	KPC	+	15	+	15	+
3401F	KPC	+	15	+	15	+
3409F	KPC	+	15	+	15	+
3436F	KPC	+	15	+	15	+
3443F	KPC	+	15	+	15	+
3446F	KPC	+	15	+	15	+
3818F	KPC	+	15	+	15	+
821F	NDM	+	30	+	15	+
871F	NDM	+	15	+	45	+
1233F	NDM	+	15	+	30	+
2007F	NDM	+	15	+	15	+
2130F	NDM	+	15	+	15	+
2610F	NDM	–	NA	+	15	+
2612F	NDM	+	45	+	15	+
2748F	NDM	–	NA	+	15	+
3035F	NDM	–	NA	+	15	+
3304F	NDM	–	NA	+	15	+
3320F	NDM	+	15	+	15	+
3763F	NDM	+	15	+	15	+
3768F	NDM	+	15	+	15	+
1888F	IMP	+	15	+	15	+
3349F	IMP	+	15	+	15	+
1047F	GES-5	+	60	–	NA	+
1597F	GES-5	+	15	+	15	–
2818F	GES-5	+	15	+	15	+
3691F	GES-5	+	75	+	75	+
1534F	OXA-370	–	NA	+	15	–
1636F	OXA-370	–	NA	+	15	+
2169F	OXA-370	–	NA	+	60	–
2246F	OXA-370	–	NA	–	NA	–
2494F	OXA-370	–	NA	+	60	–
2592F	OXA-370	–	NA	+	15	+
2729F	OXA-370	–	NA	+	15	–
2807F	OXA-370	–	NA	+	30	–
3023F	OXA-370	–	NA	+	30	–
3149F	OXA-370	+	30	+	15	–
3284F	OXA-370	–	NA	+	30	–
3704F	OXA-370	–	NA	+	30	–
3413F	Negative	+	30	NV	NA	–
3452F	Negative	–	NA	–	NA	+

NA, not applicable; NV, not validated.

one isolate required the maximum time of 75 min to become positive (Table 2). The Carbapenemase Inactivation Method presented a sensitivity of 74.4% and specificity of 97.5%. The CIM required a standard time of at least 480 min between the beginning and the final results of the test (Table 2).

Discussion

Lately a variety of novel methods for carbapenemase detection have been proposed.^{9,12,15–17} All methods explore the carbapenem-hydrolysing activity of the β-lactamases and are

Table 2 – Summary of the characteristics of the rapid tests for carbapenemase detection Carba NP, Blue-Carba and CIM.

	Sensitivity (%)	Specificity (%)	Maximum time until final result (min)
Carba NP	62.7	97.5	75
Blue-Carba	95.3	100.0	75
CIM	74.4	97.5	480

supposed to present high sensitivity and specificity within a short period of time, with results interpretable in a single day of work. In this study, we evaluated the rapid tests Carba NP, Blue-Carba and CIM.

Carba NP presented high specificity, but its sensitivity was the lowest among the other methodologies. The reduced sensitivity of Carba NP was mostly due to OXA-370 isolates. This finding was also reported in other studies that evaluated the performance of Carba NP with OXA-48-producing isolates.^{18,19} The Carba NP method proved to be a very fast methodology as most positive results were observed in 15 min. In fact, although the bacterial extract and the carbapenem were incubated for 120 min, no significant colour change was observed after 75 min, which was considered the final time results for this test.

The Blue-Carba methodology presented the highest sensitivity and specificity among the three methodologies. Only one OXA-370 positive isolate and one GES positive isolate were not detected by Blue-Carba. Pasteran et al. observed very similar results of sensitivity and specificity for this assay when they tested isolates from very diverse locations in Latin America.²¹ As found in the Carba NP test, most isolates presented positive results in 15 min and the final result for Blue-Carba was observed after 75 min of incubation.

Regarding CIM, we have modified the original protocol of van der Zwaluw et al.¹² and added zinc sulphate in the suspension of bacteria with the meropenem disk to achieve a higher sensitivity of the method regarding the detection of metallo-beta-lactamase. As observed with Carba NP, a lower sensitivity was obtained among the OXA-370 producers. The major setback of this methodology, compared to the other two evaluated in this study, is the time needed for the final result, considering that at least 6 h are necessary for a proper growth of the carbapenem-susceptible *E. coli*, which is mandatory for the test interpretation. On the other hand, the CIM seems to be the simplest test, considering that no special reagents are necessary for its execution.

The difficulty for detection of isolates harboring OXA-48-like is probably related to the least expressive hydrolytic capacity of class D carbapenemases.²⁰ Also, it is still not clear if OXA-370 (a variant of OXA-48) is a true carbapenemase, considering that low carbapenem MIC values are usually observed in clinical samples presenting this enzyme. In this matter, the detection of imipenem hydrolysis, in which the tests are based, is even harder. If OXA-370 producing isolates were not included in this study, the sensitivity for CarbaNP test would be of 83.8% and of 96.7% for CIM. However, the OXA-370 seems to be the variant of OXA-48 that is prevalent in Brazil and, therefore, it is important that the rapid tests be evaluated against these isolates.

Noteworthy, minimal color changes, yielded by lower carbapenem hydrolysis levels, were easier to observe using the Blue-Carba test, considering that blue-green changes are easier to observe, compared to red-orange ones. In some cases, especially regarding OXA-370- and GES-5-producing isolates, the interpretation of tests results were challenging, and a second opinion was requested. Thus, the correct interpretation relies on capacitated technicians.

The global spread of carbapenemases highlights the importance of their rapid detection. Considering that carbapenems

are the last resort for serious *Enterobacteriaceae* infections, controlling the dissemination of the mechanisms of resistance to these antibiotics is mandatory.²² Although PCR is considered the gold standard for carbapenemase detection,¹⁶ most laboratories are not able to perform this methodology due to costs and the special equipment required. The methodologies evaluated in this study provide reliable results in a single work-day. The carbapenemase detection might also be done directly from clinical samples, as described by Dortet et al.¹⁸

In summary, the Blue-Carba presented the highest sensitivity and, therefore, can be considered the best test to be used as a screening phenotypic methodology, considering that the misdetection of carbapenemase producers has a greater impact for clinicians than a false positive result, and differences observed between sensitivity values are much greater than the differences observed in specificity. On the other hand, CIM might be the easiest test to perform, as it does not require any special reagent and it is easier to be interpreted. The early detection of carbapenemases helps to establish infection control measures and may prevent carbapenemases to spread reducing the risk of healthcare associated infections and therapeutic failure.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

1. Paterson DL. Resistance in gram-negative bacteria: *Enterobacteriaceae*. *Am J Med*. 2006;119:S20–S28 [discussion S62–S70].
2. Nordmann P, Naas T, Poirel L. Global spread of Carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis*. 2011;17:1791–1798.
3. Ambler RP. The structure of beta-lactamases. *Philos Trans R Soc Lond B Biol Sci*. 1980;289:321–331.
4. Rozales FP, Ribeiro VB, Magagnin CM, Pagano M, Lutz L, Falci DR, et al. Emergence of NDM-1-producing *Enterobacteriaceae* in Porto Alegre, Brazil. *Int J Infect Dis*. 2014;25:79–81.
5. Magagnin CM, Rozales FP, Antochevis L, Nunes LS, Martins AS, Barth AL, et al. Dissemination of bla (OXA-370) gene among several *Enterobacteriaceae* species in Brazil. *Eur J Clin Microbiol Infect Dis*. 2017 [Epub ahead of print].
6. Picao RC, Santos AF, Nicoletti AG, Furtado GH, Gales AC. Detection of GES-5-producing *Klebsiella pneumoniae* in Brazil. *J Antimicrob Chemother*. 2010;65:796–797.
7. Lincopan N, McCulloch JA, Reinert C, Cassetari VC, Gales AC, Mamizuka EM. First isolation of metallo-beta-lactamase-producing multiresistant *Klebsiella pneumoniae* from a patient in Brazil. *J Clin Microbiol*. 2005;43:516–519.
8. Nordmann P, Poirel L, Dortet L. Rapid detection of carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis*. 2012;18:1503–1507.
9. Pires J, Novais A, Peixe L. Blue-carba, an easy biochemical test for detection of diverse carbapenemase producers directly from bacterial cultures. *J Clin Microbiol*. 2013;51:4281–4283.
10. Carvalhaes CG, Cayo R, Visconde MF, Barone T, Frigatto EA, Okamoto D, et al. Detection of carbapenemase activity directly from blood culture vials using MALDI-TOF MS: a

- quick answer for the right decision. *J Antimicrob Chemother.* 2014;69:2132–2136.
11. Lasserre C, De Saint Martin L, Cuzon G, Bogaerts P, Lamar E, Glupczynski Y, et al. Efficient detection of carbapenemase activity in *Enterobacteriaceae* by matrix-assisted laser desorption ionization-time of flight mass spectrometry in less than 30 minutes. *J Clin Microbiol.* 2015;53:2163–2171.
 12. van der Zwaluw K, de Haan A, Pluister GN, Bootsma HJ, de Neeling AJ, Schouls LM. The carbapenem inactivation method (CIM), a simple and low-cost alternative for the Carba NP test to assess phenotypic carbapenemase activity in gram-negative rods. *PLoS ONE.* 2015;10:e0123690.
 13. Clinical & Laboratory Standards Institute – CLSI. *Performance Standards of Antimicrobial Susceptibility Testing; 27th Informational Supplement; M100-S27.* Wayne, PA: Clinical and Laboratory Standards Institute; 2017.
 14. Monteiro J, Widen RH, Pignatari AC, Kubasek C, Silbert S. Rapid detection of carbapenemase genes by multiplex real-time PCR. *J Antimicrob Chemother.* 2012;67:906–909.
 15. Dorette L, Poirel L, Nordmann P. Rapid identification of carbapenemase types in *Enterobacteriaceae* and *Pseudomonas* spp. by using a biochemical test. *Antimicrob Agents Chemother.* 2012;56:6437–6440.
 16. Martino MD, Koga PC, Pasternak J, Doi AM, Ciola CS, da Silva CB, et al. Evaluation of a new rapid test for carbapenemase detection in carbapenem resistant *Enterobacteriaceae*. *J Microbiol Methods.* 2015;115:20–21.
 17. Saito R, Koyano S, Dorin M, Higurashi Y, Misawa Y, Nagano N, et al. Evaluation of a simple phenotypic method for the detection of carbapenemase-producing *Enterobacteriaceae*. *J Microbiol Methods.* 2015;108:45–48.
 18. Dorette L, Brechard L, Poirel L, Nordmann P. Rapid detection of carbapenemase-producing *Enterobacteriaceae* from blood cultures. *Clin Microbiol Infect.* 2014;20:340–344.
 19. Huang TD, Berhin C, Bogaerts P, Glupczynski Y. Comparative evaluation of two chromogenic tests for rapid detection of carbapenemase in *Enterobacteriaceae* and in *Pseudomonas aeruginosa* isolates. *J Clin Microbiol.* 2014;52:3060–3063.
 20. Evans BA, Amyes SG. OXA beta-lactamases. *Clin Microbiol Rev.* 2014;27:241–263.
 21. Pasteran F, Veliz O, Ceriana P, Lucero C, Rapoport M, Albornoz E, et al. Evaluation of the blue-carba test for rapid detection of carbapenemases in gram-negative bacilli. *J Clin Microbiol.* 2015;53:1996–1998.
 22. Walsh TR. Emerging carbapenemases: a global perspective. *Int J Antimicrob Agents.* 2010;36(Suppl 3):S8–S14.