



Hyperventilation increases the rate of the rise in arterial blood desflurane concentration during induction - a Gas Man® simulation

Junyong In

Dongguk University, Ilsan Hospital, Department of Anesthesiology and Pain Medicine, Goyang, Gyeonggi-do/Republic of Korea.

Email: dragona1@dumc.or.kr

Tel.: 82 31 9617875

Lu et al. described that hyperventilation accelerated the desflurane uptake from the alveolar to the arterial blood (1). It is a very intriguing study of a direct determination of the desflurane concentration in the arterial blood (Ades). In comparison to the end-tidal volatile anesthetic concentration, the arterial concentration of a volatile anesthetic is more likely to reflect its concentration in the brain (1-3).

A pharmacokinetic model for volatile anesthetics has been developed and used for over 25 years (4). Therefore, the effect of ventilation on Ades can be simulated with this model. Gas Man® (Version 4.1, Med Man Simulations, Inc., Chestnut Hill, MA, USA) is a computer simulation program which contains a five-compartment system built statistically which simulates the pharmacokinetics for volatile anesthetic uptake and distribution (4) with embedded parameters (5). This program helps users and readers understand anesthetic uptake and distribution with graphs. If the embedded parameters are adjusted prior to a simulation, simulated results can be seen in graphs individually.

For the simulations, the necessary parameters and patient data were taken from the report of Lu et al. (1): an anesthesia breathing circuit volume of 6 L; a fresh gas flow of 6 L/min; a functional residual capacity of 2.4 L; 61.8, 62.0, and 58.0 kg patient (hyperventilation, normal ventilation, and hypoventilation, mean values from each group); an alveolar minute ventilation of 3.33, 2.38, and 1.80 L/min, mean values from each group; cardiac output which was derived from the cardiac index of the report of Lu et al. with Du Bois method (5.38, 5.56, and 5.36 L/min, mean values from each group). Other parameters were not manipulated. Therefore, the values for compartment volumes (vessel rich group, muscle, fat, and venous group) and the values for regional blood flow percentages (vessel rich group, muscle, and fat) showed some differences depending on the entered weight. At 0, 1, 3, 5, 10, 20, 30, and 40 minutes, the simulated Ades were recorded.

In these simulations, despite having some limitations on the data used (e.g. using mean values of weight and alveolar ventilation) in the report of Lu et al, Gas Man® shows




Figure 1 - Simulated arterial concentration of desflurane (%) time curves under three different ventilatory settings, including hyperventilation, normal ventilation, and hypoventilation in Gas Man® (Version 4.1, Med Man Simulations, Inc., Chestnut Hill, MA, USA).

similar trends with the simulated Ades just as those of Lu et al (1). Hyperventilation seems to produce a higher Ades-over-time curve than normal ventilation and hypoventilation. Simulated Ades in the hyperventilation group also seems to increase more rapidly than those in the normal ventilation and the hypoventilation groups during the first 5 minutes (Figure 1).

REFERENCES

1. Lu CC, Lin TC, Hsu CH, Yu MH, Chen TL, Chen RM, et al. Hyperventilation accelerates the rise of arterial blood concentrations of desflurane in gynecologic patients. Clinics. 2012;67(9):1029-34, <http://dx.doi.org/10.6061/clinics/2012/0908>.
2. Lu CC, Tsai CS, Ho ST, Chueng CM, Wang JJ, Wong CS, et al. Pharmacokinetics of desflurane uptake into the brain and body. Anaesthesia. 2004;59(3):216-21, <http://dx.doi.org/10.1111/j.1365-2044.2003.03654.x>.
3. Lu CC, Lin TC, Yu MH, Chen TL, Lin CY, Chiang JS, et al. Effects of changes in alveolar ventilation on isoflurane arterial blood concentration and its uptake into the human body. Pharmacology. 2009;83(3):150-6, <http://dx.doi.org/10.1159/000187719>.
4. Philip JH. Gas Man--an example of goal oriented computer-assisted teaching which results in learning. Int J Clin Monit Comput. 1986;3(3):165-73, <http://dx.doi.org/10.1007/BF01716358>.
5. Eger EI, 2nd. Anesthetic uptake and action, Baltimore: Williams & Wilkins. 1974.

Copyright © 2013 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

DOI: [10.6061/clinics/2013\(03\)LE02](http://dx.doi.org/10.6061/clinics/2013(03)LE02)