

Molecular and antigenic characterisation of ribosomal phosphoprotein P0 from *Babesia bovis*

Carlos AN Ramos^{1,2}, Flávio R Araújo^{2/+}, Ingrid IF Souza², Renato HM Oliveira², Carina Elisei², Cleber O Soares², Ana MS Sacco³, Grácia MS Rosinha², Leucio C Alves¹

¹Departamento de Medicina Veterinária, Universidade Federal Rural de Pernambuco, Recife, PE, Brasil

²Embrapa Gado de Corte, BR262 km4 CP154, 79002970 Campo Grande, MS, Brasil ³Embrapa Pecuária Sul, Bagé, RS, Brasil

Babesia bovis is a tick-borne pathogen that remains an important constraint for the development of cattle industries in tropical and subtropical regions of the world. Effective control can be achieved by vaccination with live attenuated phenotypes of the parasite. However, these phenotypes have a number of drawbacks, which justifies the search for new, more efficient immunogens based mainly on recombinant protein technology. In the present paper, ribosomal phosphoprotein P0 from a Brazilian isolate of *B. bovis* was produced and evaluated with regard to conservation and antigenicity. The protein sequence displayed high conservation between different Brazilian isolates of *B. bovis* and several Apicomplexa parasites such as *Theileria*, *Neospora* and *Toxoplasma*. IgG from cattle experimentally and naturally infected with *B. bovis* as well as IgG₁ and IgG₂ from naturally infected cattle reacted with the recombinant protein. IgG from cattle experimentally infected with *Babesia bigemina* cross-reacted with *B. bovis* recombinant P0. These characteristics suggest that P0 is a potential antigen for recombinant vaccine preparations against bovine babesiosis.

Key words: *Babesia bovis* - babesiosis - bovine - recombinant ribosomal phosphoprotein

Bovine babesiosis caused by *Babesia bovis* and *Babesia bigemina* continues to be an economically important disease with wide distribution in tropical and subtropical regions, where the common arthropod vector *Rhipicephalus microplus* (Murrell & Barker 2003) occurs (Bock et al. 2004). *B. bovis* causes the most pathogenic disease among parasites of the genus, resulting from sequestration of infected erythrocytes in the microcapillary endothelia of vital organs and a hypotensive shock syndrome denominated cerebral babesiosis, which is often fatal to susceptible cattle (Wright & Goodger 1988, Clark & Jacobsen 1998). Cattle that survive initial infection, either naturally or following chemotherapy, remain persistently infected and resistant to clinical disease, thereby allowing the use of attenuated phenotypes as live vaccines. However, these vaccines have limitations, including the transmission of other blood-borne pathogens and failure due to vaccine breakdowns (de Vos & Bock 2000), which justifies the search for new, more efficient immunogens.

Advances in molecular techniques have led to the discovery and evaluation of many subunit antigens of *B. bovis* in recent years, which have provided antigens of the parasite with immunogenic and antigenic potential (Gaffar et al. 2004, Brown et al. 2006, Brayton et al. 2007, Bono et al. 2008). P0 is a neutral protein found in

all eukaryotic organisms (Rich & Steitz 1987) and is a component of a family of ribosomal phosphoproteins, together with P1 and P2 (Rajeshwari et al. 2004). P0 (but not P1 and P2) is of vital importance to cells, as demonstrated in *Saccharomyces cerevisiae* (Santos & Ballesta 1994). Although a ribosomal component, this protein has been located on the surface of many eukaryotic cells, including many protozoan parasites (Singh et al. 2002), which may at least partially explain the antigenicity and immunogenicity of P0 against *Plasmodium yoellii*, *Leishmania major* and *Babesia microti* (Chatterjee et al. 2000, Iborra et al. 2005, Terkawi et al. 2007b). Moreover, in vitro studies have demonstrated that anti-P0 antiserum can neutralise *Toxoplasma gondii*, *Neospora caninum* and *B. bovis* parasites by either inhibiting their growth or blocking cell invasion (Sehgal et al. 2003, Terkawi et al. 2007a, Zhang et al. 2007). Thus, we hypothesise that P0 is a potential candidate as an immunogen against bovine babesiosis.

The aims of the present paper were to analyse the genetic conservation of the P0 gene among Brazilian isolates of *B. bovis* and produce recombinant P0 (rP0) from a Brazilian isolate and evaluate its antigenicity with sera from cattle naturally or experimentally infected with *B. bovis* and sera from cattle experimentally infected with *B. bigemina* in an enzyme-linked immunosorbent assay (ELISA).

MATERIAL AND METHODS

Isolates - Five Brazilian isolates of *B. bovis* were used. The isolates were from the states of Bahia, São Paulo, Rio Grande do Sul (RS), Mato Grosso do Sul and Rondônia, representing all physiographic Regions of Brazil (Northeast, Southeast, South, Midwest and North, respectively).

Financial support: EC (INCO 003691-MEDLABAB), CNPq, FUNDECT, FACEPE

+ Corresponding author: flabio@cnpqg.embrapa.br

Received 26 May 2009

Accepted 23 September 2009

Gene sequencing and analysis - Whole ORFs of the P0 gene from the five Brazilian isolates of *B. bovis* were amplified by polymerase chain reaction (PCR) and cloned into *pGEM-T Easy* (Promega, USA) for sequence analysis, following the manufacturer's instructions. Amplifications of the whole gene were performed in two steps using the primer sets P0-1F 5'-ATGGCTCGCATGAGCAAG-3' and P0-1R 5'-AC-CCTGTCGTTAACCTTGATGA-3' and P0-2F 5'-AT-GTCCACCTCATCAAGGTTAAC-3' and P0-2R 5'-TTAGTCAAAGAGGGAGAATCCC-3' to produce two amplicons. Amplification reactions were performed in a volume of 25 μ L containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 0.2 mM of each deoxynucleoside triphosphate, 12 pmol of each primer, 100 ng of genomic DNA and 1.25 U of *Taq* DNA polymerase (Cenbiot, Brazil). Amplifications were performed in a Mastercycler thermocycler (Eppendorf, Germany) as follows: 94°C for 1 min (denaturation); 30 cycles of 94°C for 1 min (denaturation), 53°C for 30 sec (annealing) and 72°C for 40 sec (extension) and a final cycle at 72°C for 4 min (extension). PCR products were analysed by electrophoresis on 1% agarose gels stained with SybrGold (Invitrogen, USA).

The DNA sequences of P0 from five Brazilian isolates of *B. bovis* were obtained using the BigDye Terminator kit (Applied Biosystems, USA). Sequences were assembled using the Sequencher v.4.1.4 software program (Gene Codes, USA) and submitted to BLASTn search (<http://www.ncbi.nlm.nih.gov>) to verify the sequence identity. Multiple sequence alignment was performed with the ClustalW algorithm (www.ebi.ac.uk/Tools/clustalw2/index.html).

Phylogenetic analyses were conducted with deduced amino acid sequences of P0 from Brazilian isolates of *B. bovis*, other species of *Babesia*, other Apicomplexa and *Bos taurus* using the MEGA software program, version 4.0 (Tamura et al. 2007). A phylogenetic tree was generated using the neighbour-joining method (Saitou & Nei 1987) and a bootstrap resampling technique of 1,000 replications was performed to statistically support the reliabilities of the nodes on the trees (Felsenstein 1985).

Cloning and expression - DNA from the southern Brazilian isolate of *B. bovis* was used to amplify the complete ORF of P0 by PCR using the primer set P0-1F and P0-2R. The amplification reaction was performed in a volume of 25 μ L, as described above, with an initial denaturing step of 95°C for 3 min followed by 35 cycles of 95°C for 1 min, 58°C for 30 sec and 72°C for 1 min and a final extension step at 72°C for 4 min.

The amplicon was initially cloned into *pGEM-T Easy* (Promega, USA) and subsequently sub-cloned into the *pRSET-c* (Invitrogen, USA) *EcoRI* site. Chemically competent *Escherichia coli* host strain BL-21 cells were transformed with recombinant plasmid (*pRSET-c/P0*) and the transformation was spread on selective plates containing Luria Bertani (LB) medium supplemented with 100 μ g/mL ampicillin and 34 μ g/mL chloramphenicol. The expression of the P0 gene was carried out in LB broth with the addition of 1 mM isopropyl- β -D-galactopyranoside.

The purification of rP0 was carried out with agarose-Ni⁺ resin (ProBond, Invitrogen, USA), following the manufacturer's instructions. The purified rP0 was analysed by sodium dodecyl-polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting with monoclonal anti-histidine antibody (GE, UK).

Protein concentration was determined by SDS-PAGE stained with Coomassie blue by comparison with known concentrations of bovine serum albumin using the LabImage v.3.3.2 image analysis software (Loccus, Brazil).

Serum samples - Cattle serum samples negative for antibodies to *B. bovis* and *B. bigemina* were collected from tick-free areas of RS (n = 33) and Paraíba (n = 29), Brazil and from cattle kept in a tick-free isolation area of Embrapa Beef Cattle, Campo Grande, MS, Brazil (n = 67). Serum samples from cattle naturally infected with *B. bovis* were obtained from state of Pará (PA), Brazil (n = 133). These serum samples were analysed by a crude *B. bovis* antigen ELISA (Madruga et al. 2000). Serum samples from cattle experimentally infected with *B. bovis* (n = 77) or *B. bigemina* (n = 36) were obtained from the South Embrapa Cattle and Sheep Research Centre, Bagé, RS, Brazil. These animals were raised in a tick-free area and were periodically tested for antibodies to *B. bovis*, *B. bigemina* and *Anaplasma marginale* by the immunofluorescent antibody test and examination of stained blood smears. Experimental infections with *B. bovis* or *B. bigemina* were carried out by the subcutaneous route with pure isolates kept in liquid nitrogen and sera were collected 30 days post-inoculation.

Standardisation of ELISA with rP0 - Optimal dilutions of antigen, sera and conjugate were determined using six sera negative and six sera positive for antibodies against *B. bovis*. Then, 96-well Costar 3590 plates (Corning, USA) were adsorbed with 1 ng/mL of rP0 (100 μ L/well) diluted in phosphate buffer containing 0.1% Tween 20 (PBST) for 12 h at 4°C. Plates were blocked with PBST with 5% skim milk for 1 h at 37°C. After five washes with PBST, control and test sera (in duplicate) - diluted 1:800 in PBST - were incubated for 60 min at 37°C. Plates were then washed five times with PBST and 100 μ L/well of rabbit anti-bovine IgG horseradish peroxidase conjugate (Sigma A5295, USA) diluted 1:10,000 in PBST was added. Plates were incubated for 30 min at 37°C and, after five washes with PBST, reactions were revealed with Fast-OPD (50 μ L/well; Sigma, USA). The reaction was stopped by the addition of 100 μ L/well of H₂SO₄ (2.5 N) and results were read on an EL-800 ELISA reader (Bio-Tek, USA) with a 490-nm filter.

IgG₁ and IgG₂ titration - For titration of anti-rP0 IgG₁ and IgG₂, sera from 30 cattle naturally infected with *B. bovis* from PA were tested as described above, except that sera were diluted from 1:100-1:3,800, anti-bovine IgG₁ and IgG₂ monoclonal antibodies (Serotec, USA) were diluted 1:5,000 and rabbit anti-mouse IgG horseradish conjugate (Sigma A-9044, USA) was diluted 1:10,000. Titres were defined as the reciprocal of the highest dilution giving a positive result.

Cutoff selection - The cutoffs of the ELISA tests were determined based on those described by Frey et al. (1998).

Statistical analysis - The means of the optical densities of sera tested for IgG were analysed by ANOVA followed by Tukey's test, with a 95% confidence level. Titres of IgG₁ and IgG₂ were analysed by Wilcoxon signed-rank test, with a 95% confidence level.

RESULTS AND DISCUSSION

The P0 gene sequences (939 nucleotides) obtained in the present study were deposited in the Genbank database under the following accession: FJ588004 Midwestern isolate, FJ588005 Northeastern isolate, FJ588006 Northern isolate, FJ588007 Southeastern isolate and FJ588008 Southern isolate of *B. bovis*.

The multiple alignment of P0 DNA sequences revealed a high degree of identity among the Brazilian isolates of *B. bovis* ($\geq 99\%$). The best hit by BLASTn was the Genbank entry AF498365 (99% nucleotide identity). The amino acid sequence homology among Brazilian isolates was 100%. This demonstrates that P0 is highly conserved among isolates of *B. bovis*, which is a key prerequisite for any subunit vaccine candidate.

The P0-deduced amino acid sequences of the Brazilian isolates of *B. bovis* were aligned with P0-deduced amino acid sequences of *B. bovis* USA AAM18123, *Babesia caballi* BAF91359, *B. bigemina* BAF91357, *Babesia gibsoni* BAF45423, *Babesia equi* BAF91358, *Theileria annulata* XP_954430, *N. caninum* BAF62528, *T. gondii* AAK69358 and *B. taurus* AAI51696. A considerable level of homology was observed between the P0 sequences of the Brazilian isolates of *B. bovis* and *B. caballi* (85%), *B. bigemina* (84%), *B. gibsoni* (80%), *B. equi* (67%), *T. annulata* (62%), *N. caninum* (58%) and *T. gondii* (58%). This high degree of homology may explain the serological cross-reactivity or cross-protective immunity between the P0 proteins of *B. gibsoni* and *B. microti* (Terkawi et al. 2007b), *B. gibsoni* and *B. rodhaini* (Terkawi et al. 2008), *B. gibsoni* and *B. bovis* (Terkawi et al. 2007a) and others. The high degree of homology also suggests that P0 may be a universal immunogen against bovine babesiosis caused by *B. bovis* and *B. bigemina*. Nevertheless, there was considerable homology between the P0 proteins from *B. bovis* and *B. taurus* (46%). This indicates the possibility that immunisation with rP0 from *B. bovis* may generate autoimmune antibodies in cattle. The phylogenetic analysis resulted in a tree with two branches (Fig. 1). The first one included Apicomplexa parasites, with a subgroup containing all the isolates of *B. bovis* (Brazilian and North American), another subgroup with *B. bigemina*, *B. caballi* and *B. gibsoni*, a third subgroup with *T. annulata* and *B. equi* and the last one with the Sarcocystidae *N. caninum* and *T. gondii*. The second branch was of *B. taurus*.

The high degree of P0 conservation among protozoa and even in mammals, suggests that this protein plays an important role in cell metabolism. P0 has been associated with DNA repair, cell development, apoptosis and carcinogenesis (Brockstedt et al. 1998, Frolov & Birchler 1998, Liliensiek et al. 1998). Moreover, P0 is a structural

component, together with P1 and P2, of the 60S ribosomal subunit of eukaryotes (Uchiumi et al. 1987) and, in knockout studies on *S. cerevisiae*, P0 was shown to be vital to cell survival (Santos & Ballesta 1994).

rP0 from *B. bovis* was produced as a 6x-histidine-tag fusion, resulting in a protein of approximately 37 kDa viewed by SDS-PAGE and western blotting (Fig. 2). This molecular mass is consistent with the fusion protein,




Fig. 1: phylogenetic analysis of deduced amino acid sequences of P0 from Brazilian isolates of *Babesia bovis*, other species of *Babesia*, other Apicomplexa and *Bos taurus*. a: NCBI Genbank accession; MW: Midwest Region; N: North Region; NE: Northeast Region; S: South Region; SE: Southeast Region.




Fig. 2A: Coomassie blue-stained sodium dodecyl-polyacrylamide gel electrophoresis (SDS-PAGE) profile of *Escherichia coli* BL-21 extracts expressing P0 gene from a Brazilian isolate (Southern Region) of *Babesia bovis*. Lanes 1: molecular mass marker (Promega ref V8491); 2: extract of *E. coli* BL-21; 3: BL-21 cells transformed with pR7/P0; 4: semi-purified recombinant P0; B: Western blot analysis with recombinant P0 (rP0) from a Brazilian isolate (Southern Region) of *Babesia bovis* and anti-6x histidine monoclonal antibody (GE, UK). Lanes 1: extract of *E. coli* BL-21; 2: BL-21 cells transformed with pRSETc/P0; 3: purified rP0.

including the his-tag (~3 kDa) and P0 (34 kDa). Using anti-sera against *B. gibsoni* P0, Terkawi et al. (2007a) detected protein bands of 31-34 kDa in lysates of *B. gibsoni*, *B. bovis*, *B. bigemina*, *B. equi* and *B. caballi* through western blot analysis.

The recognition of rP0 from *B. bovis* by cattle IgG was evaluated by ELISA (Fig. 3). Among the 77 sera from cattle experimentally infected with *B. bovis*, 75 (97.4%) were positive in the rP0 ELISA. Out of 133 sera from cattle naturally infected with *B. bovis*, 127 (95.5%) were positive. Out of 129 sera from cattle negative for *B. bovis* by ELISA with crude *B. bovis* antigen, 128 (99.2%) were also negative in the rP0 ELISA.

To investigate whether rP0 from *B. bovis* is recognised by *B. bigemina* antibodies, sera from cattle exclusively infected with *B. bigemina* were tested with rP0 by ELISA (Fig. 3). Out of 36 sera analysed, 34 (94.4%) were positive in the rP0 ELISA. Serological cross-reactions against P0 have been described among *Babesia* species (Terkawi et al. 2007a) and between *Plasmodium falciparum* and *T. gondii* (Sehgal et al. 2003). The cross-reaction found in the present paper suggests that P0 proteins from *B. bovis* and *B. bigemina* share immunodominant epitopes and raises the possibility that such epitopes may be involved in a cross-protective immune response.

rP0 from *B. bovis* was also recognised by IgG₁ and IgG₂ antibodies from 30 cattle naturally infected with *B. bovis* from PA, with a significantly higher titre of IgG₂ ($p = 0.0121$) (Fig. 4). Th1 responses, including the production of interferon- γ , activation of macrophages and production of the IgG₂ isotype by B cells, have been associated with immunity against intra-erythrocytic pathogens (Brown et al. 1998, Aguilar-Delfin et al. 2003).

There is the possibility that the IgG, IgG₁ and IgG₂ responses found in the present study may be elicited by cross-reactions with antibodies against P0 proteins from other Apicomplexa parasites, such as *N. caninum*. Nevertheless, only 0.8% of the sera collected from cattle raised in




Fig. 3: distribution of optical densities of cattle sera in ELISA test with *Babesia bovis* recombinant P0. 1: cattle negative for *Babesia bovis*; 2: cattle naturally infected with *B. bovis*; 3: cattle experimentally infected with *B. bovis*; 4: cattle experimentally infected with *Babesia bigemina*. Different letters indicate significant statistical differences by ANOVA and Tukey's test (95% confidence level).




Fig. 4: IgG₁ and IgG₂ titers anti-recombinant P0 of *Babesia bovis* in 30 cattle naturally infected from the state of Pará, Brazil. Titers of IgG₁ and IgG₂ were analyzed by Wilcoxon signed-rank test, with a 95% confidence level.

tick-free areas did not react with rP0, suggesting that cross-reactions are relevant only at the *Babesia* genus level.

In summary, the results of the present paper indicate the high conservation of *B. bovis* P0 not only between different *B. bovis* Brazilian isolates, but also between many protozoan parasites, as well as the antigenicity of rP0 with *B. bovis* and *B. bigemina*-infected sera. These two characteristics are desirable for the subunit vaccine.

REFERENCES

- Aguilar-Delfin I, Wettstein PJ, Persing DH 2003. Resistance to acute babesiosis is associated with Interleukin-12 and gamma interferon mediated responses and requires macrophages and natural killer cells. *Infect Immun* 71: 2002-2008.
- Bock R, Jacksson L, de Vos A, Jorgensen W 2004. Babesiosis of cattle. *Parasitology* 129: 247-269.
- Bono MF, Mangold AJ, Baravalle ME, Valentini BS, Thompson CS, Wilkowsky SE, Echaide IE, Farber MD, Torioni De Echaide SM 2008. Efficiency of a recombinant MAS-2c-based ELISA to establish the persistence of antibodies in cattle vaccinated with *Babesia bovis*. *Vet Parasitol* 157: 203-210.
- Brayton KA, Lau AO, Herndon DR, Hannick L, Kappmeyer LS, Berens SJ, Bidwell SL, Brown WC, Crabtree J, Fadrosch D, Feldblum T, Forberger HA, Haas BJ, Howell JM, Khouri H, Koo H, Mann DJ, Norimine J, Paulsen IT, Radune D, Ren Q, Smith RKJr, Suarez CE, White O, Wortman JR, Knowles DP Jr, McElwain TF, Nene VM 2007. Genome sequence of *Babesia bovis* and comparative analysis of apicomplexan hemoproteins. *PLoS Pathog* 3: 1401-1413.
- Brockstedt E, Rickers A, Kostka S, Laubersheimer A, Dorken B, Wittmann-Liebold B, Bommert K, Otto A 1998. Identification of apoptosis-associated proteins in a human Burkitt lymphoma cell line Cleavage of heterogeneous nuclear ribonucleoprotein A1 by caspase 3. *J Biol Chem* 273: 28057-28064.
- Brown WC, Norimine J, Goff WL, Suarez CE, McElwain TF 2006. Prospects for recombinant vaccines against *Babesia bovis* and related parasites. *Parasite Immunol* 28: 315-327.
- Brown WC, Shkap V, Zhu D, McGuire TC, Tuo W, McElwain TF, Palmer GH 1998. CD4⁺ T-lymphocyte and immunoglobulin G2 responses in calves immunized with *Anaplasma marginale* outer

- membranes and protected against homologous challenge. *Infect Immun* 66: 5406-5413.
- Chatterjee S, Singh S, Sohoni R, Kattige V, Deshpande C, Chiplunkar S, Kumar N, Sharma S 2000. Characterization of domains of the phosphoriboprotein P0 of *Plasmodium falciparum*. *Mol Biochem Parasitol* 107: 143-154.
- Clark IA, Jacobsen L 1998. Do babesiosis and malaria share a common disease process? *Ann Trop Med Parasitol* 92: 483-488.
- de Vos A, Bock RE 2000. Vaccination against bovine babesiosis. *Ann NY Acad Sci* 916: 540-545.
- Felsenstein J 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783-791.
- Frey A, Di Canzio J, Zurawski D 1998. A statistically defined endpoint titer determination method for immunoassays. *J Immunol Methods* 221: 35-41.
- Frolov MV, Birchler JA 1998. Mutation in P0, a dual function ribosomal protein/apurinic/aprimidinic endonuclease, modifies gene expression and position effect variegation in *Drosophila*. *Genetics* 150: 1487-1495.
- Gaffar FR, Yatsuda AP, Franssen FF, Vries EA 2004. *Babesia bovis* merozoite protein with a domain architecture highly similar to the thrombospondin-related anonymous protein (TRAP) present in *Plasmodium* sporozoites. *Mol Biochem Parasitol* 136: 25-34.
- Iborra S, Carrión J, Anderson C, Alonso C, Sacks D, Soto M 2005. Vaccination with the *Leishmania infantum* acidic ribosomal P0 protein plus CpG oligodeoxynucleotides induces protection against cutaneous leishmaniasis in C57BL/6 mice but does not prevent progressive disease in BALB/C mice. *Infect Immun* 73: 5842-5852.
- Liliensiek B, Rocha M, Umansky V, Benner A, Lin J, Ziegler R, Nawroth PP, Schirmacher V 1998. Identification of four genes in endothelial cells whose expression is affected by tumor cells and host immune status a study in *ex vivo*-isolated endothelial cells. *Blood* 92: 3394-3404.
- Madrugá CR, Araújo FR, Marques APC, Carvalho CME, Cusinato FQ, Crocci AJ, Kessler RH, Miguita M 2000. Desenvolvimento de uma prova de imunoadsorção enzimática para detecção de anticorpos contra *Babesia bovis*. *Pesq Vet Bras* 20: 167-170.
- Murrel A, Barker SC 2003. Synonymy of *Boophilus* Curtice, 1891 with *Rhipicephalus* Koch, 1844 (Acari: Ixodidae). *Syst Parasitol* 56: 169-172.
- Rajeshwari K, Patel K, Nambeesan S, Mehta M, Sehgal A, Chakraborty T, Sharma S 2004. The P domain of the P0 protein of *Plasmodium falciparum* protects against challenge with malaria parasites. *Infect Immun* 72: 5515-5521.
- Rich BE, Steitz JA 1987. Human acidic ribosomal phosphoproteins P0, P1 and P2: analysis of cDNA clones, *in vitro* synthesis and assembly. *Mol Cell Biol* 7: 4065-4074.
- Saitou N, Nei M 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406-425.
- Santos C, Ballesta JPG 1994. Ribosomal protein P0 contrary to phosphoproteins P1 and P2 is required activity and *Saccharomyces cerevisiae* viability. *J Biol Chem* 269: 15689-15696.
- Sehgal A, Kumar N, Carruthers V, Sharma S 2003. Translocation of ribosomal protein P0 on *Toxoplasma gondii* tachyzoite surface. *Int J Parasitol* 33: 1589-1594.
- Singh S, Sehgal A, Waghmare S, Chakraborty T, Goswami A, Sharma S 2002. Surface expression of the conserved ribosomal protein P0 on parasite and other cells. *Mol Biochem Parasitol* 119: 121-124.
- Tamura K, Dudley J, Nei M, Kumar S 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 24: 1596-1599.
- Terkawi MA, Jia H, Gabriel A, Goo YK, Nishicawa Y, Yokoyama N, Igarashi I, Fujisaki K, Xuan X 2007a. A shared antigen among *Babesia* species: ribosomal phosphoprotein P0 as a universal babesia vaccine candidate. *Parasitol Res* 102: 35-40.
- Terkawi MA, Jia H, Zhou J, Lee E, Igarashi I, Fujisaki K, Nishikawa Y, Xuan X 2007b. *Babesia gibsoni* ribosomal phosphoprotein P0 induces cross-protective immunity against *B. microti* infection in mice. *Vaccine* 25: 2027-2035.
- Terkawi MA, Zhang G, Jia H, Aboge G, Goo YK, Nishikawa Y, Yokoyama N, Igarashi I, Kawazu SI, Fujisaki K, Xuan X 2008. C3 contributes to the cross-protective immunity induced by *Babesia gibsoni* phosphoriboprotein P0 against a lethal *B. rodhaini* infection. *Parasite Immunol* 30: 365-370.
- Uchiumi T, Wahha AJ, Traut RR 1987. Topography and stoichiometry of acidic proteins in large ribosomal subunits from *Artemia salina* as determined by cross linking. *Proc Natl Acad Sci USA* 84: 5580-5584.
- Wright IG, Goodger BV 1988. Pathogenesis of babesiosis. In M Ristic, *Babesiosis of domestic animals and man*, CRC Press, Boca Raton, 99-118.
- Zhang H, Lee E, Liao M, Compaore MKA, Zhang G, Kawase O, Fujisaki K, Sugimoto C, Nishikawa Y, Xuan X 2007. Identification of ribosomal phosphoprotein P0 of *Neospora caninum* as a potential common vaccine candidate for the control of both neosporosis and toxoplasmosis. *Mol Biochem Parasitol* 153: 141-148.