

Molecular markers and genetic diversity of *Plasmodium vivax*

Cristiana Ferreira Alves de Brito^{1/+}, Marcelo Urbano Ferreira²

¹Laboratório de Malária, Instituto de Pesquisas René Rachou-Fiocruz, Av. Augusto de Lima 1715, 30190-002 Belo Horizonte, MG, Brasil

²Departamento de Parasitologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, SP, Brasil

Enhanced understanding of the transmission dynamics and population genetics for Plasmodium vivax is crucial in predicting the emergence and spread of novel parasite phenotypes with major public health implications, such as new relapsing patterns, drug resistance and increased virulence. Suitable molecular markers are required for these population genetic studies. Here, we focus on two groups of molecular markers that are commonly used to analyse natural populations of P. vivax. We use markers under selective pressure, for instance, antigen-coding polymorphic genes, and markers that are not under strong natural selection, such as most minisatellite and microsatellite loci. First, we review data obtained using genes encoding for P. vivax antigens: circumsporozoite protein, merozoite surface proteins 1 and 3a, apical membrane antigen 1 and Duffy binding antigen. We next address neutral or nearly neutral molecular markers, especially microsatellite loci, providing a complete list of markers that have already been used in P. vivax populations studies. We also analyse the microsatellite loci identified in the P. vivax genome project. Finally, we discuss some practical uses for P. vivax genotyping, for example, detecting multiple-clone infections and tracking the geographic origin of isolates.

Key words: malaria - *Plasmodium vivax* - genetic markers - genetic diversity - multiple-clone infections - microsatellites

Plasmodium vivax, a relatively neglected human malaria parasite, is a major public health challenge for Central and South America, the Middle East, Central, South and Southeast Asia, Oceania and East Africa, where 2.85 billion people are currently at risk of infection and 70-80 million clinical cases are reported each year (Mueller et al. 2009, Guerra et al. 2010). The emergence of drug-resistant strains and severe (sometimes fatal) disease challenges the traditional view of *vivax* malaria as a benign infection (Price et al. 2009). Recent epidemiological trends in Brazil illustrate the importance of *P. vivax* as a re-emerging pathogen. The annual incidence of *Plasmodium falciparum* (the predominant malaria parasite species between 1985-1990) decreased steadily during the 1990s, while that of *P. vivax* maintained an upward trend. Both *P. falciparum* and *P. vivax* are still transmitted across the Amazon Basin, with rare *Plasmodium malariae* infections. However, *P. vivax* now causes 85% of the 315,000 malaria cases reported in this country each year (Oliveira-Ferreira et al. 2010), suggesting that this species may be less susceptible to the malaria control strategies currently used in Brazil.

Understanding the genetic structure of *P. vivax* is essential to accurately describing the transmission dynamics of *vivax* malaria. Population genetics data are crucial, for example, to predict how fast phenotypes of interest, such as novel antigenic variants, particular relapsing patterns or drug resistance, arise and spread

in natural populations (Zilversmit & Hartl 2005). The current data on *P. vivax* population genetics lags behind that of *P. falciparum*, primarily due to the scarcity of appropriate genetic markers. As we cannot propagate the parasite in continuous culture in vitro (Udomsangpetch et al. 2008), the phenotypic diversity of natural human infections from *P. vivax* remained unexplored until the early 1990s. Advances in molecular methods, especially implementation of polymerase chain reaction (PCR)-based protocols to amplify parasite DNA, allowed for genotyping of malaria parasite samples obtained directly from patient blood, without a preceding in vitro culture step (Kimura et al. 1990). The relative paucity of genetic markers hampers further investigation into the genotypes that underlie phenotypes such as relapse pattern, virulence and drug resistance.

In this review, we broadly classify the available molecular markers into two groups: (i) markers that are clearly under natural selection, that is, markers that map to genome regions associated with adaptive traits and (ii) markers that are neutral or nearly neutral, that is, those that are not obviously influenced by natural selection. The extensive polymorphism found in loci associated with *P. vivax* adaptive traits, such as those coding for surface antigens and drug-resistance (Cui et al. 2003a), reflects the combined effects of the parasite's population history and selective constraints imposed by the host's immunity and drug usage (Escalante et al. 2004). Thus, these loci do not provide much information on the *P. vivax* population structure. Neutral and nearly neutral markers, in contrast, may allow for unbiased estimates of genetic variation, population structure and gene flow in natural parasite populations.

When comparing different markers and parasite typing systems, a clear quantitative definition of genetic diversity is required. One of the most popular methods of summarising genetic diversity levels is to report vir-

Financial support: Rede Malária/PRONEX, MS/DECIT, FAPEMIG, FAPEMAT, NIAID, NIH (R01 AI075416-04) (to MUF)

+ Corresponding author: cristiana@cpqrr.fiocruz.br

CFADB and MUF were supported by CNPq fellowships.

Received 26 April 2011

Accepted 8 June 2011

tual heterozygosity (H_E) values. Virtual H_E is the average probability that a pair of alleles randomly obtained from the population is different. Whenever appropriate, in this review we provide H_E estimates obtained using different markers for different populations.

Molecular markers under selection - Until recently, most genetic markers available for characterising natural *P. vivax* populations were orthologues of previously identified *P. falciparum* antigen-coding genes. The well-characterised polymorphic regions in these genes have been extensively used to analyse genetic diversity patterns in *P. falciparum* (Roy et al. 2008). A similar approach has been explored in *vivax*-oriented studies (Cui et al. 2003a). The following single-copy genes in *P. vivax* have often been used in these studies: *gam1*, coding for the gametocyte antigen 1, *csp*, coding for the circumsporozoite protein (CSP), *mssl* and *mssl3alpha*, coding for the merozoite surface proteins (MSP)-1 and 3 α , respectively, *amal*, coding for the apical membrane antigen 1, and *dbp*, coding for the Duffy binding protein (DBP) (Table I).

A notable feature of several *P. vivax* surface antigens (including major vaccine-candidate molecules) is tandem arrays of relatively short amino acid motifs. The CSP in *P. vivax* (PvCSP), an abundant antigen on the surface of sporozoites that has been extensively used as a vaccine development target, has immunodominant B-cell epitopes that map to central repeats between nonrepetitive sequences (Nardin & Zavala 1998). PvCSP displays two major types of nonapeptide repeats [most commonly GDRA(D/A)GQPA and ANGA(G/D)(N/D)QPG], which define the variants known as VK210 and VK247, respectively (Arnot et al. 1985, Rosenberg et al. 1989) (Fig. 1). Both VK210 and VK247 variants are globally distributed, but geographic biases have been described (Cochrane et al. 1990, Qari et al. 1993a, b). Similar repetitive sequences are found in Brazil, Southeast Asia and Papua New Guinea (Qari et al. 1991, 1992). However, markedly divergent sequence variants were found in isolates from China and North Korea (Mann et al. 1994). Although VK210 and VK247-type sequences often occur in sympatric parasite populations, there are no examples of a hybrid repeat array with both repeat types (Lim et al. 2005).

A third type of repeat unit [APGANQ(E/G)GAA], identical to that described for *Plasmodium simiovale*, characterises the so-called *P. vivax*-like parasites (Qari et al. 1993a). Although *P. vivax*-like CSP repeats have been found in parasite isolates across the world, its global distribution remains unconfirmed (Gopinath et al. 1994). In Brazil, *P. vivax*-like parasites have been identified only in mixed-clone infections with VK210-type or VK247-type parasites (Machado & Póvoa 2000). More recently, complex mixtures of parasites harbouring several PvCSP types and co-infecting the same hosts have been demonstrated by Henry-Halldin et al. (2011) in Papua New Guinea.

The nonapeptide sequences in PvCSP are repeated ~20 times in the array. Insertions and deletions in the central repeat domain, from either sexual recombination during meiosis or intrahelical strand-slippage events during mitotic DNA replication (McConkey et al. 1990), generate novel CSP variants that may be positively select-




Fig. 1: schematic representation of studied antigens: A: the circumsporozoite protein of *Plasmodium vivax* (PvCSP) consists of a conserved region (light grey) flanking a repeated central region (RCR) (blank box), which contains immunodominant B-cell epitopes and is flanked by pre and post-repeat specific nonrepetitive sequences (NR) (black boxes). PvCSP displays three types of repeats [GDRA(D/A)GQPA (VK210), ANGA(G/D)(N/D)QPG (VK247)] which are the most prevalent and APGANQ[E/G]GGAA (*P. vivax*-like); B: the *P. vivax* merozoite surface protein 1 (PvMSP-1) sequence comprises seven conserved blocks (amino acid similarity among PvMSP-1 alleles, 71-85%) (light grey boxes) and six variable blocks (amino acid similarity, 21-34%) (black boxes), described by Putaporntip et al. (2002). For comparison, we also show the original division of PvMSP-1 into interspecies conserved blocks (ICBs) (amino acid similarity > 48% in pairwise comparisons of MSP-1 orthologues in *Plasmodium falciparum*, *P. vivax* and *Plasmodium yoelii*) (light grey boxes), conserved blocks (CBs) (amino acid similarity > 50% between *P. falciparum* and *P. vivax*, but lower in other pairwise comparisons) (dark grey boxes) and polymorphic blocks (amino acid similarity < 45%) (black boxes) (del Portillo et al. 1991). These two methods of portioning PvMSP-1 differ in that Putaporntip et al. (2002) compared intraspecific sequences, while del Portillo et al. (1991) compared between-specific sequences. The F1, F2 and F3 were defined by Imwong et al. (2005) (horizontal bold lines); C: the *P. vivax* merozoite surface protein 3 α (PvMSP-3 α) contains an alanine-rich central polymorphic region (blank box), which could be partially deleted, generating three types of proteins with different sizes. This region is flanked by semi-conserved regions (SC) (grey boxes) and by conserved regions (CR) (light grey boxes); D: the *P. vivax* apical membrane antigen 1 (PvAMA-1) displayed three domains in its ectodomain, referred to as domain I, II and III; E: the *P. vivax* Duffy binding protein (PvDBP) is schematically divided in seven domains: II [Duffy binding-like domain (DBL)] and VI are cysteine-rich domains. The DBL domain is the most polymorphic and responsible for the interaction with the Duffy antigen at the erythrocyte surface; CT: cytoplasmic tail; PS: pro sequence; SS: signal peptide; TM: transmembrane domain.

ed if the mutant parasites evade a host's immunity. The central repetitive domain of *P. falciparum* CSP (PfCSP) contains extensive variation. This domain consists of a variable number of four-mer NANP and NVDP motif copies without breaking down the tight linkage between polymorphic sites in flanking sequences, as in interhelical exchanges during meiosis (Rich et al. 1997). This pattern suggests that these repeats undergo frequent in-

TABLE I
Genes encoding antigens used as molecular markers to study field populations of *Plasmodium vivax*

Antigen	Protein fragment	Population studied (n)	Total ^a (n)	References
PvAMA-1	Domain I	BRA	105	Grynberg et al. (2008)
			20	Rodrigues et al. (2005)
		KOR	30	Chung et al. (2003)
			22	Han et al. (2002)
		MYA	76	Moon et al. (2010)
		AFR (5), CHI (5), IND (18), INDO (5), PHI (148), PNG (23), SI (7), THAI (6)	219	Figtree et al. (2000)
	Domain I, II	IND	61	Thakur et al. (2008)
	Whole	IND	11	Rajesh et al. (2007)
		SRL	23	Gunasekera et al. (2007)
		THAI	231	Putaporntip et al. (2009)
VEN		73	Ord et al. (2008)	
PvCSP	CR	AFG	202	Zakeri et al. (2010)
		AZE	36	Leclerc et al. (2004)
		BRA	48	Machado and Póvoa (2000)
			45	Patil et al. (2010)
			32	Santos-Ciminera et al. (2007)
			155	Storti-Melo et al. (2009)
			126	Gopinath et al. (1994)
			24	Hernández-Martínez et al. (2011)
			61	Bonilla et al. (2006)
			151	Kim et al. (2006)
	Central AME, South AME, AFR, Southeast ASI, IND	COL	24	Hernández-Martínez et al. (2011)
		GUY	61	Bonilla et al. (2006)
		IND	151	Kim et al. (2006)
		IRA	144	Zakeri et al. (2006)
		KOR	632	Choi et al. (2010)
		MYA	116	Moon et al. (2010)
		PAK	187	Zakeri et al. (2006)
		PAK (187), IRA (150)	337	Zakeri et al. (2010)
		PNG	40	Kolakovich et al. (1996)
			507	Henry-Halldin et al. (2011)
		THAI	90	Cui et al. (2003b)
			100	Imwong et al. (2005)
	171	Kain et al. (1993)		
	17	Rongnoparut et al. (1995)		
CR, NR	BRA (30), PNG (14)	30	Qari et al. (1992)	
N, C-end	PHI, CHI, SI, PNG	18	Huang et al. (1994)	
Whole	CHI	16	Mann et al. (1994)	
PvDBP	Whole	COL	23	Martinez et al. (2004)
	Domain II-IV	South KOR	30	Kho et al. (2001)
	Domain II	BRA	40	Sousa et al. (2006)
			122	Sousa et al. (2010)
		COL	20	Ampudia et al. (1996)
		COL (18), South KOR (14), PNG (68)	100	Cole-Tobian et al. (2002)
		IRA	75	Babaeekho et al. (2009)
		PNG	40	Kolakovich et al. (1996)
			358	Cole-Tobian et al. (2002)
			50	Tsuboi et al. (1994)
			24	Xainli et al. (2000)
		THAI	30	Gosi et al. (2008)
	Domain II, III	KOR	8	Suh et al. (2001)

Antigen	Protein fragment	Population studied (n)	Total ^a (n)	References	
PvMSP-1	Pv200L	COL	26	Valderrama-Aguirre et al. (2011)	
	925 pb	PHI, CHI, SI, PNG	18	Cheng et al. (1993)	
	BL 2	BRA (28), VIET (23)	51	Hoffmann et al. (2003)	
	ICBs 2-4	COL	20	Gutierrez et al. (2000)	
	ICBs 4, 5	AFR (5), CHI (10), IND (21), INDO (11), PHI (97), PNG (21), SI(6), THAI (4)	KOR	30	Kim et al. (2009)
			AZE	36	Leclerc et al. (2004)
	ICBs 5, 6 or BL 5	IND (9), COL (11)	BRA	32	Santos-Ciminera et al. (2007)
			French GUY	120	Veron et al. (2009)
			IND	25	Farooq et al. (2009)
			KOR	632	Choi et al. (2010)
				25	Lim et al. (2000)
			MYA	116	Moon et al. (2009)
			PNG	40	Kolakovich et al. (1996)
			PAK (187), IRA (150)	337	Zakeri et al. (2010)
			SRL	22	Premawansa et al. (1993)
			THAI	15	Putaporntip et al. (1997)
	F1-F3	IND	THAI	100	Imwong et al. (2005)
			PNG	93-108	Koepfli et al. (2009)
	MSP-1 ₄₂	SRL	IND	33	Thakur et al. (2008)
			BRA	28	Soares et al. (1999)
MSP-1 ₁₉	GUY		61	Bonilla et al. (2006)	
		PHI (4) THAI (3) SI (8), PNG (1), South East ASI (4)	20	Pasay et al. (1995)	
C-terminal Whole	THAI (20), BRA (8), BAN (5), South KOR (4), VAN (2), IND (1)	KOR	45	Putaporntip et al. (2002)	
		TUR	30	Han et al. (2011)	
				Zeyrek et al. (2010)	
PvMSP-3 α	NA	IND	151	Kim et al. (2006)	
		COL	55	Cristiano et al. (2008)	
		French GUY	120	Veron et al. (2009)	
		IND	27	Prajapati et al. (2010)	
		IRA	144	Zakeri et al. (2006)	
		MYA	116	Moon et al. (2009)	
		PAK	50	Khatoon et al. (2010)	
		PAK (187), IRA (150)	337	Zakeri et al. (2010)	
		PER	186	Sutton et al. (2009)	
		PNG	28	Bruce et al. (2000)	
			11	Mueller et al. (2002)	
		PNG (39), IND (4), Sudan (1), SRL (1)	45	Bruce et al. (1999)	
		South KOR	24	Han et al. (2004)	
		THAI	90	Cui et al. (2003b)	
VEM	58	Ord et al. (2005)			
Whole	THAI	17	Mascorro et al. (2005)		

a: number of *P. vivax* infected patients studied in genotyping; AFG: Afghanistan; AFR: Africa; AMA: apical membrane antigen; AME: America; ASI: Asia; AZE: Azerbaijan; BAN: Bangladesh; BL: polymorphic block (Putaporntip et al. 1997); BRA: Brazil; CHI: China; COL: Colombia; CR: central repetitive; CSP: circumsporozoite protein; DBP: Duffy binding protein; ICB: interspecies conserved blocks (del Portillo et al. 1991); IND: India; INDO: Indonesia; IRA: Iran; KOR: Korea; MSP: merozoite surface protein; MYA: Myanmar; NA: not applied restriction fragment length polymorphism-polymerase chain reaction; NR: non-repetitive; PAK: Pakistan; PER: Peru; PHI: Philippines; PNG: Papua New Guinea; Pv: *Plasmodium vivax*; SI: Solomon Islands; SRL: Sri Lanka; THAI: Thailand; TUR: Turkey; VAN: Vanuatu; VEN: Venezuela; VIET: Vietnam.

trahelical recombination that either add or delete repeat units during mitotic DNA replication (Fig. 2), as other short repeats, such as microsatellite and minisatellite-type sequences (Levinson & Gutman 1987). Frequent mitotic recombination coupled with positive selection of new variants might accelerate sequence evolution even where meiotic recombination and outcrossing are relatively uncommon in malaria parasite populations (Rich et al. 2000). In length variation arrays, whether length variation affects B-cell epitope recognition is uncertain, but the conformational nature of repetitive epitopes in PfCSP (Monette et al. 2001) supports this hypothesis.

We have recently examined whether similar recombination mechanisms create significant variation in the PvCSP repeat arrays, which consist of longer (9-mer) repetitive motifs (Patil et al. 2010). We investigated patterns of sequence diversity in PvCSP encoding gene alleles in sympatric *P. vivax* isolates from an area of low malaria endemicity in Brazil. In these isolates, we used single-nucleotide polymorphism typing to examine the haplotype structure of chromosome 8, where the PvCSP encoding gene is located. We sought to determine whether the repetitive domain in the *csp* locus was significantly diverse under conditions of low malaria endemicity, which reduces effective meiotic recombination rates in local parasites. We confirmed previous findings (Arnot et al. 1985, 1990, Qari et al. 1992, 1994, Mann et al. 1994) of substantial nucleotide sequence diversity in repeat arrays from 45 isolates. They were analysed using nine different *csp* alleles, each with a unique arrangement of six different nonapeptide repeat sequences, and all were of the VK210 type. Notably, different nucleotide motifs, termed repeat allotypes, coded for the same VK210-type repeat units, indicating that some conservation is maintained at the amino acid level (putatively due to functional constraints), but not necessarily at the nucleotide level (Rich et al. 1997, 2000). Most repeat units in the same *csp* allele were either identical or nearly identical to each other, consistent with their recent expansion in the repeat array. We found strong linkage disequilibrium at sites across the chromosome 8 segment flanking the *csp* locus, consistent with rare meiotic recombination in this region. This suggests that repeat array diversity may not be severely constrained by low meiotic recombination rates that are typical in areas with low malaria endemicity. New repeat variants may be readily created by nonhomologous recombination events with potential implications for PvCSP-based vaccine development.

The MSP-1 and 3 α are additional examples of blood-stage vaccine-candidate antigens with repetitive arrays. The 200-kDa MSP 1 in *P. vivax* (PvMSP-1), originally described at the molecular level by del Portillo et al. (1991), is a major target of naturally acquired (Nogueira et al. 2006) and vaccine-induced immunity (Yang et al. 1999, Herrera et al. 2007). The protein contains six highly polymorphic domains (4 of them repetitive) flanked by seven fairly conserved sequences (Putapornpip et al. 2002) (Fig. 1). Sequence analysis has confirmed that blocks 1, 3 and 5 in PvMSP-1 are conserved at the protein level, while blocks 2, 4 and 6 are highly variable




Fig. 2: a strand-slippage event in a repetitive sequence during mitotic DNA replication. The nascent (upper) strand becomes displaced from the parental (lower) strand during DNA replication and re-anneals out of phase, inserting a repeat motif (grey square) without changing non-repetitive flanking sequences (grey rectangles). A similar slippage process in the parental strand would delete a one repeat motif from the nascent strand. These processes are suggested to generate variation in repeat arrays of surface malarial antigens, such as circumsporozoite protein of *Plasmodium vivax* (PvCSP) and in minisatellite and microsatellite-type sequences.

(Kolakovich et al. 1996, Figtree et al. 2000, Hoffmann et al. 2003, Bastos et al. 2007, Farooq et al. 2009, Kim et al. 2009, Veron et al. 2009, Zakeri et al. 2010).

The extensive sequence divergence in variable domains of PvMSP-1 (amino acid similarity, 21-34%) has been maintained by balanced selection, most likely as a result of variant-specific immune pressure (Putapornpip et al. 2006). However, the extent to which PvMSP-1 sequence diversity affects immune recognition for this major malaria-vaccine candidate antigen remains uncertain. To address this topic, we have recently compared naturally acquired antibody responses to three polymorphic domains and the highly conserved PvMSP-1 C-terminal domain in a rural, malaria-exposed Amazonian cohort. We expressed 15 recombinant proteins corresponding to PvMSP-1 variants commonly found in local parasites and showed that less than one-third had detectable IgG antibodies to at least one expressed variant of blocks 2, 6 and 10. However, 54.3% recognised the invariant C-terminal domain PvMSP-1₁₉. Although the proportion of PvMSP-1 variant responders increased substantially during subsequent acute *P. vivax* infections, the specificity of IgG antibodies did not necessarily match the PvMSP-1 variant(s) found in infecting parasites (Bastos et al. 2007). The mechanisms underlying the limited immune recognition of PvMSP-1 variants that are known to circulate in a given endemic area remain unclear.

P. vivax evidences an allelic form diversity across the *mspl* gene. In contrast, *P. falciparum* exhibits a clear allelic dimorphism over a long region in the *mspl* locus (that is, sequences can be grouped into 1 of 2 major allelic types, MAD20 and K1) (Roy et al. 2008). Fre-

quent meiotic recombination events that shuffle different sequence types to generate new alleles are responsible for the mosaic structure of PvMSP-1 (Putaporntip et al. 1997, 2002). Recently, Han et al. (2011) compared PvMSP-1 alleles from re-emergent *P. vivax* parasites collected in the Republic of Korea over the past decade. They showed an increase in allelic diversity for variable domains of this protein, suggesting that recombination independently changed protein domains. Moreover, the greater similarity between allelic sequences within *P. vivax* than between *P. vivax* and closely related species was taken as evidence that polymorphisms in *P. vivax* arose recently. This is in stark contrast to the *P. falciparum* pattern, wherein different alleles are estimated to have diverged more than 25 million years ago (Polley et al. 2005). However, the cause for the large differences in this genetic history for these highly divergent parasites remains obscure. It may reflect either the significantly different sequence structure of the orthologues, different molecular functions for the gene in the two species, a different population history, or another event.

Proteolytic processing of PvMSP-1 generates a C-terminal fragment of 42 kDa (PvMSP-1₄₂), which is subsequently cleaved to generate two smaller fragments with apparent molecular masses 33 kDa (PvMSP-1₃₃) and 19 kDa (PvMSP-1₁₉). Only the small, PvMSP-1₁₉ fragment remains at the merozoite surface during and after erythrocyte invasion (Holder 2009). Limited polymorphism has been observed in the C-terminal 19-kD PvMSP-1 fragment, with only two polymorphic sites (Soares et al. 1999, Dias et al. 2011). PvMSP-1₄₂, like PfMSP-1₄₂, exhibits extensive genetic polymorphism in natural infections (Escalante et al. 1998, Conway et al. 2000, Putaporntip et al. 2002, Pacheco et al. 2007, Thakur et al. 2008, Dias et al. 2011).

The *P. vivax msp3a* gene encodes a MSP with an apparent molecular mass ranging from 148-150 kDa. The protein contains an alanine-rich central domain that is predicted to form a coiled-coil tertiary structure (Galinski et al. 1999) (Fig. 1). The gene coding for the MSP-3 α (PvMSP-3 α) is highly polymorphic and its variability has been assessed by restriction fragment length polymorphism-PCR analysis (Bruce et al. 1999). The suitability of PvMSP-3 α as a molecular marker has been confirmed in several geographic parasite populations: Papua New Guinea (Bruce et al. 1999), Thailand (Cui et al. 2003b, Mascorro et al. 2005), Pakistan and Iran (Zakeri et al. 2006, 2010, Khatoon et al. 2010), Venezuela (Ord et al. 2005), Peru (Sutton et al. 2009), Colombia (Cristiano et al. 2008), India (Prajapati et al. 2010) and French Guiana (Veron et al. 2009). However, classifying restriction fragments according to size after electrophoresis is relatively subjective, which hampers large-scale analyses and reduces comparisons between studies. In addition, complex restriction patterns generated by mixed-clone infections may be difficult to interpret.

The *P. vivax* apical membrane antigen-1 (PvAMA-1) is an immunogenic, type I integral membrane protein, which is expressed at the apical surface of merozoites and sporozoites. It seems to play a role in erythrocyte and hepatocyte invasion (Hodder et al. 2001, Silvie et al. 2004). A number of studies have addressed polymorphism in that PvAMA-1 encoding gene in various parts

of the world, such as Asia, Oceania and South America (Cheng & Saul 1994, Figtree et al. 2000, Han et al. 2002, Chung et al. 2003, Rodrigues et al. 2005, Gunasekera et al. 2007, Rajesh et al. 2007, Grynberg et al. 2008, Thakur et al. 2008, Moon et al. 2010). The PvAMA-1 protein includes three extracellular domains, known as domains I, II and III (Pizarro et al. 2005) (Fig. 1). Most studies show a high degree of genetic variability in domain I (Ord et al. 2008, Thakur et al. 2008, Putaporntip et al. 2009), while the Sri Lanka domain II is the most polymorphic (Gunasekera et al. 2007). A comparison of PvAMA-1 domain I sequences from 320 worldwide isolates revealed a striking divergence between Old World populations and those from Brazil (Grynberg et al. 2008, Putaporntip et al. 2009). Sequence analysis of Sri Lankan isolates suggested that balancing selection maintains polymorphism at the *P. vivax ama1* locus (Gunasekera et al. 2007), while no strong evidence for balancing selection was found in Thailand (Putaporntip et al. 2009). A number of studies have reported a substantially lower variability at this locus in *P. vivax* than in *P. falciparum*, implying that natural selection acts differently at this locus in the two species (Gunasekera et al. 2007, Ord et al. 2008, Putaporntip et al. 2009).

P. vivax DBP (PvDBP) plays an important role in the formation of an irreversible junction between *P. vivax* merozoites and its receptor, Duffy antigen/receptor for chemokines (DARC), at the surface of immature red blood cells (reticulocytes). This is a key step in host cell invasion (Singh et al. 2005, Galinski & Barnwell 2009). In vitro, antibodies against PvDBP inhibit DBP binding to DARC and block invasion of human erythrocytes (Michon et al. 2000, Grimberg et al. 2007, Ceravolo et al. 2008). The most polymorphic segment of this protein is the erythrocyte-binding motif, a 170 amino-acid stretch located in the cysteine-rich domain II (Ranjan & Chitnis 1999, VanBuskirk et al. 2004, Hans et al. 2005) (Fig. 1). This segment of the protein has been used as molecular marker in genetic population studies of field isolates from Papua New Guinea (Tsuboi et al. 1994, Xainli et al. 2000), Colombia (Ampudia et al. 1996), South Korea (Kho et al. 2001), Brazil (Sousa et al. 2006, 2010) and Thailand (Gosi et al. 2008). Recently, the importance of selective diversifying pressure has been shown for specific regions of the protein structure (Sousa et al. 2010).

Although molecular epidemiological studies using *P. vivax gam1* gene have revealed variability in *P. vivax* isolates from Sri Lanka (Snewin et al. 1995), Korea (Kho et al. 2001) and India (Prajapati et al. 2006), amplification of this gene has been associated with artefacts (Imwong et al. 2001, 2005). For this reason, *P. vivax gam1* is no longer used as a reliable genetic marker.

Neutral or nearly neutral molecular markers - Feng et al. (2003) described 33 tandem repeat (TR) arrays consisting of repeat units with four or more nucleotides. However, not much diversity was revealed by analysing these arrays in four monkey-adapted *P. vivax* strains (India VII, Salvador-I, Belem, Thai-NYU). Leclerc et al. (2004) used the eight most polymorphic TR loci described by Feng et al. (2003) to show much more variability in TRs than microsatellites in isolates from dif-

ferent endemic areas. However, little TR diversity was found within populations (Tables II, III). Rezende et al. (2009) selected the five most polymorphic loci from Leclerc's study for their analysis of field isolates from four regions in Brazil. This set of TR markers again revealed relatively little diversity in Brazilian isolates (Table III). Most TR markers had a clearly predominant allele. Moreover, these markers were inefficient in detecting multiple-clone infections and were unable to group the isolates according to their geographical origin (Rezende et al. 2009). As these TR markers map to a 100-kb region in chromosome 8 with many putative protein-coding genes (including *csp*), natural selection is expected to affect them, even if indirectly, as a result of hitchhiking (Feng et al. 2003). Thus, these loci may not be effectively neutral given their proximity to sequences under strong diversifying or purifying selection (Leclerc et al. 2004). An alternative explanation for the low variability in TR loci may include their structure, as they are relatively short arrays consisting of long (> 6 bp) repeat units. These sequences are less prone to strand slippage events (Fig. 2) than typical microsatellite-type sequences consisting of long arrays of short repeat motifs.

The current markers of choice for large-scale population genetic studies in eukaryotes are highly polymorphic and short (1-6 bp-long) TRs, known as microsatellites (Schlötterer 1998). In contrast to the ~1,000 polymorphic microsatellite loci currently available for *P. falciparum* typing, it was demonstrated that a few dozen microsatellite loci are polymorphic in *P. vivax* field isolates (Table II). Despite these limitations, microsatellite-based stud-

ies of the parasite have provided valuable information on *P. vivax* population structure and diversity in the Americas, Asia and Africa (Ferreira et al. 2007, Karunaweera et al. 2007, 2008, Koepfli et al. 2009, Orjuela-Sánchez et al. 2009a, Gunawardena et al. 2010, Rezende et al. 2010, Van den Eede et al. 2010a, b, 2011).

The first microsatellite described for *P. vivax* was a dinucleotide (AT) repeat, used to identify high variability in 89 isolates from Papua New Guinea (Gómez et al. 2003) (Tables II, III). Leclerc et al. (2004) reported low variability in dinucleotide microsatellite loci. In this study, only one of 13 microsatellites showed extensive variability and nine were entirely monomorphic among the eight *P. vivax* populations analysed (Table III). Using a draft of the unpublished *P. vivax* genome, Imwong et al. (2006) designed primer pairs for 11 dinucleotide microsatellites and showed extensive variability in *P. vivax* populations from areas with intermediate to high levels of endemicity. They compared their findings with those of Leclerc et al. (2004) and explained the discrepancy in microsatellite diversity levels as the result of repeat array length at the loci studied. They suggested that microsatellite diversity correlates positively repeat array length, as the rate of strand-slippage events that create diversity increases exponentially with repeat array lengths. Recently, Rezende et al. (2010) corroborated this hypothesis using a set of microsatellites with di, tri and tetra nucleotide repeat units. They highlighted the importance of microsatellite loci structure in genetic diversity studies using malaria parasites. Karunaweera et al. (2007) described a new set of highly polymorphic

TABLE II
Description of molecular markers: minisatellites and microsatellites used in genetic population analysis of *Plasmodium vivax* field isolates

Group of molecular markers	Genomic regions			Number of microsatellites				References
	Coding (hypothetical) ^a	Non-coding	Total	Types of repeat unit (imperfect and/or composed) ^b				
				Di	Tri	Tetra	> 4	
I	4 (2)	4	8 ^c	0	1 (1)	3	4	Feng et al. (2003)
II	1	0	1	1	0	0	0	Gomez et al. (2003)
III	1	12	13	13 (9)	0	0	0	Feng et al. (2003)
IV	2 (1)	6	11 ^d	11	0	0	0	Imwong et al. (2006)
V	2 (1)	6	8	0	1	3	4	Imwong et al. (2007)
VI	13 (9)	1	14	0	13 (7)	1 (1)	0	Karunaweera et al. (2007)
VII	7 (2)	17	24	7 (2)	10 (3)	6	1	Joy et al. (2008)
VIII	4 (3)	7	11	4	3	4	0	Rezende et al. (2010)
IX	3 (2)	0	3	0	3 (1)	0	0	Van den Eede et al. (2010a)

a: number of microsatellites lying in hypothetical proteins among the microsatellites which lie in coding regions; *b*: number of microsatellites imperfect and/or composed among the total number of microsatellites, with different repeat unit sizes [dinucleotides (Di), trinucleotides (Tri) and tetranucleotides (Tetra) or more than 4 nucleotides]; *c*: in the original manuscript, these molecular markers were designated as tandem repeats, including minisatellites and microsatellites; *d*: three loci could not be assigned to chromosomes.

tricleotide and tetranucleotide microsatellites, which are expected to yield more accurate allele scores than dinucleotide markers (Anderson et al. 1999).

P. vivax genome sequencing allowed for further characterisation of a panel including ~160 microsatellites (Carlton et al. 2008). While searching for a repeat unit with 2-6 nucleotides, the authors found only one microsatellite locus with repeats consisting of more than four nucleotides. Most (76%) loci had perfect repeat units (Fig. 3A), in contrast with the markers described by Karunaweera et al. (2007). All except one microsatellite were assigned to chromosomes, but their chromosome distribution was clearly heterogeneous. Chromosomes 1 and 4 had the lowest marker density, whereas chromosomes 14 and 9 had the highest density (Fig. 3B). Despite the relatively low AT content in the *P. vivax* genome, when compared with that of *P. falciparum*, *P. vivax* microsatellite-type repeats are particularly AT-rich (Fig. 3C). To characterise polymorphic microsatellite markers, these loci were analysed in eight monkey-adapted *P. vivax* strains (Brazil I, Miami II, Pakchong, Panama I, Nicaragua, Thai II, Vietnam IV and Indonesia XIX). Nineteen markers were monomorphic among these samples, none were polymorphic in all strains and the large majority of markers were polymorphic only in 2-4 strains (Fig. 3D). Further, only one-fourth of the markers shown to be polymorphic in natural *P. vivax* populations were also identified as polymorphic using these eight *P. vivax* strains.

High variability in *P. vivax* isolates has been demonstrated using different sets of microsatellites (Table III). However, this comparison is complicated, as it is difficult to determine how much of the variation depends on the set of markers and how much is due to genetic diversity in the studied populations (Imwong et al. 2007a, Koepfli et al. 2009). Currently, the most widely used markers are a set of 14 polymorphic microsatellites described by Karunaweera et al. (2007), which were used to analyse the population structures of *P. vivax* in Brazil, Vietnam, Sri Lanka, Myanmar, Ethiopia and Peru (Ferreira et al. 2007, Karunaweera et al. 2008, Gunawardena et al. 2010, Van den Eede et al. 2010a, b, 2011). Using the same set of markers, similar variability was detected in different geographical areas, with H_E ranging from 0.71 in Brazil to 0.86 in Myanmar and Vietnam; only Peru showed a slightly lower variability ($H_E = 0.58$). More recently, Joy et al. (2008) selected 24 microsatellites of ~160, described by Carlton et al. (2008), and used them to show local adaptation between mosquito vectors and different genetic populations of *P. vivax* from Mexico. There was no correlation between the number of alleles identified by Carlton et al. (2008) and the variability identified using a large number of *P. vivax* field isolates using the same set of markers by Joy et al. (2008). These data reinforce the uncertainty in using laboratory strains to identify polymorphic markers (Fig. 4). Koepfli et al. (2009) showed high variability in Papua New Guinea isolates using this

TABLE III

Genetic variability analysis of *Plasmodium vivax* field populations using minisatellites and microsatellites as molecular markers

MM group (n)	n	Geographical origin of isolates (n)	Alleles ^a				References
			n	H_E	MI		
I	96	AZE (37), THAI (15), TUR (16), VEN(16), ETH(8)	1-8	0.26-0.67	NA	Leclerc et al. (2004)	
I (5)	44	BRA	2-9	0.57	0-66	Rezende et al. (2009)	
II	89	PNG	15	0.85	4.5	Gomez et al. (2003)	
III	108	AZE (40), THAI (20), TUR (19), VEN (18), ETH (7), MOZ (1), PNG (1), SRL (1)	1-9	0.35-0.80	NA	Leclerc et al. (2004)	
IV	82	COL (27), IND (27), THAI (28)	7-18	0.64-0.77	NA	Imwong et al. (2006)	
V	345	IND (90), LAO (81), THAI (92), COL (82)	9-35	0.72-0.79	40-80	Imwong et al. (2007)	
VI	25	SRL	6-13	0.78	NA	Karunaweera et al. (2007)	
VI	74	BRA	3-14	0.71	48	Ferreira et al. (2007)	
VI	132	SRL (50), BRA (74), VIET (8)	9-43	0.65-0.908	60 ^b	Karunaweera et al. (2008)	
VI (12)	425	SRI (140), MYA (167), ETH (118)	3-44	0.75-0.86	65	Gunawardena et al. (2010)	
VII	234	MEX	4-8 ^d	0.61	16	Joy et al. (2008)	
V (2), VI (1)	93-108	PNG	13-67	0.93	NA	Koepfli et al. (2009)	
VIII	53	BRA	4-31	0.71	60	Rezende et al. (2010)	
VI, IX	69	VIET	6-41	0.86	100 ^c	Van den Eede et al. (2010a)	
VI, IX	94	PER	2-8	0.58	11-70 ^c	Van den Eede et al. (2010b)	

a: number of alleles per locus only in *P. vivax* isolates; b: data for samples collected in 2005; c: including all observed alleles above 100 arbitrary fluorescence units (except for MS16 -> 25%); d: average per locus; AZE: Azerbaijan; BRA: Brazil; COL: Colombia; ETH: Ethiopia; H_E : heterozygosity expected per population using the set of polymorphic markers; IND: India; LAO: Lao; MEX: Mexico; MI: multiple clone infections; MM: molecular marker group defined in Table I (number of markers used was provided only if was not used all markers of the group); MOZ: Mozambique; MYA: Myanmar; n: number of isolates; NA: not available; PER: Peru; PNG: Papua New Guinea; SRL: Sri Lanka; THAI: Thailand; TUR: Turkey; VEN: Venezuela; VIET: Vietnam.

set of microsatellites and three selected from other studies (Table II). Rezende et al. (2010) described a set of 11 microsatellites used to show a positive correlation between transmission levels and genetic variability in *P. vivax* isolates from geographical regions in Brazil. Taken together, these data suggest that, regardless of endemicity in the isolates' geographical origin, most studies show that *P. vivax* populations are highly variable across the globe.

Practical uses for *P. vivax* genotyping - The vast field of infectious disease molecular epidemiology has produced a number of examples involving practical applications for molecular markers in addressing issues of clear biological and public health importance. Here, we briefly review four areas of potential interest: (i) detection of naturally occurring multiple-clone infections, (ii) after-treatment follow-up in clinical trials, (iii) between-species comparisons of genetic diversity levels and (iv) tracking of the geographic origin of infections.

Over the last two decades, molecular markers have revealed that many natural *P. vivax* infections comprise a complex mixture of clonal populations; that is, multiple-clone *vivax* infections are fairly common in human hosts (Havryliuk & Ferreira 2009). The earliest attempts to in-

vestigate the clonal diversity of *P. vivax* isolates in human infections focused on characterisation of phenotypes using monoclonal antibodies or isoenzymes (Udagama et al. 1987, 1990, Joshi et al. 1989), an approach that required blood samples with relatively high parasitaemias. As a rule, multilocus analysis, such as microsatellite typing, detects more multiple-clone infections than single-locus analysis, and augmenting the number of markers increases the chance of detecting multiple clones. For example, in 90 *P. vivax* isolates from Thailand, Cui et al. (2003b) found a multiple-clone infection rate of 25.6% using the PvCSP encoding gene alone, 19.3% using PvMSP-3 α alone and 35.6% when the markers were combined. Particularly surprising is the extensive clonal diversity of *P. vivax* infections in areas with relatively low malaria transmission, such as Brazil, Colombia and Sri Lanka (Ferreira et al. 2007, Imwong et al. 2007b, Karunaweera et al. 2008, Orjuela-Sánchez et al. 2009a, Gunawardena et al. 2010). The evolutionary and epidemiological consequences of multiple-clone *P. vivax* infections have not been well-investigated. However, experimental rodent malaria models suggest that, for two or more genetically distinct clones in the same host, intra-host competition for limited resources may select for *P. vivax* traits repre-




Fig. 3: characterisation of microsatellites identified in the *Plasmodium vivax* genome project (Carlton et al. 2008). A: number of microsatellites with repeat units composed of either 2, 3, 4 or > 4 nucleotides. Microsatellites were also divided into either perfect (with only 1 type of tandem repeat unit), composed (comp) (with more than 1 type of repeat unit) or imperfect (imp) (repeat units interspersed with nucleotides); B: distribution of the number of microsatellites among *P. vivax* chromosomes, including those identified by the genome project, previously described, and present in both; C: positive correlation between percentage of microsatellites and AT content in the repeat units; D: percentage of microsatellites with different numbers of alleles, analysing eight *P. vivax* laboratory lines (Carlton et al. 2008).




Fig. 4: correlation between number of alleles in eight *Plasmodium vivax* laboratory lines (Carlton et al. 2008) and the heterozygosity expected (H_E) in 234 isolates from Mexico (Joy et al. 2008), using the same set of 24 microsatellites.

senting major public health challenges, such as increased virulence, transmissibility and antimalarial drug resistance (reviewed by Havryliuk & Ferreira 2009).

Molecular markers have also been used to type *P. vivax* infections after drug treatment. Reappearance of parasitaemia after drug treatment can result from either recrudescence of surviving asexual blood-stage parasites; relapse from dormant liver stages, known as hypnozoites; or new infections with unrelated parasites. Paired parasite molecular genotyping distinguishes between recrudescences (with the same genotype as the initial infection) and new infections (with a different genotype). Until recently, relapses were thought to be caused by hypnozoites that are genetically identical to the blood-stage parasites found in primary infections (Craig & Kain 1996, Kirchgatter & del Portillo 1998). This suggested that molecular methods could easily discriminate relapses, which would have the same genotype as the primary infection, from new infections, which would have a different genotype. This view has been challenged by the recent discovery of different parasite genotypes in primary infections and relapses for 72% of *P. vivax*-infected patients from Thailand, India and Myanmar (Imwong et al. 2007b). Accurate detection of multiple-clone *P. vivax* infections is even more important in light of this report. If the primary infection comprises multiple clones, then some of them may have been missed or partially characterised or, even worse, had their alleles combined to create artificial haplotypes during genotyping. Likewise, the relapsing clone could have been either missed or incorrectly typed during the primary infection, leading to a false conclusion that different genotypes were present during the primary and relapse infections. The current consensus is that relapses may originate from reactivation of either the same parasite clone found in the primary bloodstream infection (homologous hypnozoites) or another, genetically different clone (heterologous hypnozoites). Our recent microsatellite analysis of 28 paired acute-infection and recurrence parasites from rural Amazonia revealed only two pairs of identical haplotypes (consistent with recrudescences or reactivation of homologous hypnozoites) and four pairs of related haplotypes (sharing alleles at 11-13 of 14 microsatellites analysed) (Orjuela-Sánchez et al. 2009a).

As the characteristics of endemic settings may influence the genetic diversity of the parasite, noteworthy findings could be obtained from sympatric *P. vivax* and *P. falciparum* isolates that are analysed using similar protocols. Two comparisons of microsatellite diversity in human malaria parasite species co-circulating in the same area in Brazil have been published. Both suggest that *P. vivax* infections are more diverse (that is, they comprise a higher number of alleles and virtual H_E) and comprise multiple clones more often than *P. falciparum* infections in rural Amazonia (Ferreira et al. 2007, Orjuela-Sánchez et al. 2009b). As *P. vivax* currently predominates in Brazil, differences in species-specific transmission levels might translate into differences in genetic diversity. However, we recently found more microsatellite diversity in *P. vivax* compared with *P. falciparum* populations in western Cambodia, where both species are similarly prevalent (MU Ferreira and RM Fairhurst, unpublished observations). Thus, we suggest that the higher microsatellite diversity found in *P. vivax* isolates may reflect greater plasticity in microsatellite-type short repeats for this species.

Finally, standardised molecular methods applied to worldwide populations may also help characterise population or region-specific molecular barcodes. This is essential for tracking the geographic origin of infections. A microsatellite-based analysis of *P. vivax* samples from Sri Lanka, Ethiopia and Myanmar recently showed that parasites may be classified as originating from Asia or Africa with 70-80% accuracy. This suggests that microsatellite data might be useful in predicting the origin of *P. vivax* parasites (Gunawardena et al. 2010). Further analyses are required to confirm whether microsatellite-based molecular barcodes are useful for determining whether particular infections were either transmitted locally or imported into areas approaching the elimination phase.

REFERENCES

- Ampudia E, Patarroyo M, Patarroyo M, Murillo L 1996. Genetic polymorphism of the Duffy receptor binding domain of *Plasmodium vivax* in Colombian wild isolates. *Mol Biochem Parasitol* 78: 269-272.
- Anderson TJC, Su XZ, Bockarie M, Lagog M, Day KP 1999. Twelve microsatellite markers for characterization of *Plasmodium falciparum* from finger-prick blood samples. *Parasitology* 119: 113-125.
- Arnot DE, Barnwell JW, Tam JP, Nussenzweig V, Nussenzweig RS, Enea V 1985. Circumsporozoite protein of *Plasmodium vivax*: gene cloning and characterization of the immunodominant epitope. *Science* 230: 815-818.
- Arnot DE, Stewart MJ, Barnwell JW 1990. Antigenic diversity in Thai *Plasmodium vivax* circumsporozoite proteins. *Mol Biochem Parasitol* 43: 147-149.
- Babaeekho L, Zakeri S, Djajid ND 2009. Genetic mapping of the duffy binding protein (DBP) ligand domain of *Plasmodium vivax* from unstable malaria region in the Middle East. *Am J Trop Med Hyg* 80: 112-118.
- Bastos MS, da Silva-Nunes M, Malafronte RS, Hoffmann EH, Wunderlich G, Moraes SL, Ferreira MU 2007. Antigenic polymorphism and naturally acquired antibodies to *Plasmodium vivax* merozoite surface protein 1 in rural Amazonians. *Clin Vaccine Immunol* 14: 1249-1259.

- Bonilla JA, Validum L, Cumming R, Palmer CJ 2006. Genetic diversity of *Plasmodium vivax* Pvcp3 and Pvmsp1 in Guyana, South America. *Am J Trop Med Hyg* 75: 830-835.
- Bruce MC, Galinski MR, Barnwell JW, Donnelly CA, Walmsley M, Alpers MP, Walliker D, Day KP 2000. Genetic diversity and dynamics of *Plasmodium falciparum* and *P. vivax* populations in multiply infected children with asymptomatic malaria infections in Papua New Guinea. *Parasitology* 121: 257-272.
- Bruce MC, Galinski MR, Barnwell JW, Snounou G, Day KP 1999. Polymorphism at the merozoite surface protein-3 locus of *Plasmodium vivax*: global and local diversity. *Am J Trop Med Hyg* 61: 518-525.
- Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, Caler E, Crabtree J, Anguoli SV, Merino EF, Amedeo P, Cheng Q, Coulson RM, Crabb BS, del Portillo HA, Essien K, Feldblyum TV, Fernandez-Becerra C, Gilson PR, Gueye AH, Guo X, Kang'a S, Kooij TW, Korsinczyk M, Meyer EV, Nene V, Paulsen I, White O, Ralph SA, Ren Q, Sargeant TJ, Salzberg SL, Stoeckert CJ, Sullivan SA, Yamamoto MM, Hoffman SL, Wortman JR, Gardner MJ, Galinski MR, Barnwell JW, Fraser-Liggett CM 2008. Comparative genomics of the neglected human malaria parasite *Plasmodium vivax*. *Nature* 455: 757-763.
- Cheng Q, Stowers A, Huang TY, Bustos D, Huang YM, Rzepczyk C, Saul A 1993. Polymorphism in *Plasmodium vivax* *MSA1* gene the result of intragenic recombinations? *Parasitology* 106: 335-345.
- Ceravolo I, Souza-Silva F, Fontes CJ, Braga EM, Madureira A, Kretzli AU, Souza JM, Brito CF, Adams JH, Carvalho LH 2008. Inhibitory properties of the antibody response to *Plasmodium vivax* Duffy binding protein in an area with unstable malaria transmission. *Scand J Immunol* 67: 270-278.
- Cochrane AH, Nardin EH, Arruda M, Maracic M, Clavijo P, Collins WE, Nussenzeig RS 1990. Widespread reactivity of human sera with variant repeat of the circumsporozoite protein of *Plasmodium vivax*. *Am J Trop Med Hyg* 43: 446-451.
- Cole-Tobian JL, Cortés A, Baisor M, Kastens W, Xainli J, Bockarie M, Adams JH, King CL 2002. Age-acquired immunity to a *Plasmodium vivax* invasion ligand, the Duffy binding protein. *J Infect Dis* 186: 531-539.
- Conway DJ, Cavanagh DR, Tanabe K, Roper C, Mikes ZS, Sakihama N, Bojang KA, Oduola AM, Kremsner PG, Arnot DE, Greenwood BM, McBride JS 2000. A principal target of human immunity to malaria identified by molecular population genetic and immunological analyses. *Nat Med* 6: 689-692.
- Craig AA, Kain KC 1996. Molecular analysis of strains of *Plasmodium vivax* from paired primary and relapse infections. *J Infect Dis* 174: 373-379.
- Cristiano FA, Pérez MA, Nicholls RS, Guerra AP 2008. Polymorphism in the *Plasmodium vivax* *msp 3a* gene in field samples from Tierralta, Colombia. *Mem Inst Oswaldo Cruz* 103: 493-496.
- Cui L, Escalante AA, Imwong M, Snounou G 2003a. The genetic diversity of *Plasmodium vivax* populations. *Trends Parasitol* 19: 220-226.
- Cui L, Mascorro CN, Fan Q, Rzomp KA, Khuntirat B, Zhou G, Chen H, Yan G, Sattabongkot J 2003b. Genetic diversity and multiple infections of *Plasmodium vivax* malaria in western Thailand. *Am J Trop Med Hyg* 68: 613-619.
- Cheng Q, Saul A 1994. Sequence analysis of the apical membrane antigen I (AMA-1) of *Plasmodium vivax*. *Mol Biochem Parasitol* 65: 183-187.
- Choi YK, Choi KM, Park MH, Lee EG, Kim YJ, Lee BC, Cho SH, Rhie HG, Lee HS, Yu JR, Lee JS, Kim TS, Kim JY 2010. Rapid dissemination of newly introduced *Plasmodium vivax* genotypes in south Korea. *Am J Trop Med Hyg* 82: 426-432.
- Chung JY, Chun EH, Chun JH, Kho WG 2003. Analysis of the *Plasmodium vivax* apical membrane antigen-1 gene from re-emerging Korean isolates. *Parasitol Res* 90: 325-329.
- del Portillo HA, Longacre S, Khouri E, David PH 1991. Primary structure of the merozoite surface antigen 1 of *Plasmodium vivax* reveals sequences conserved between different *Plasmodium* species. *Proc Natl Acad Sci USA* 88: 4030-4034.
- Dias S, Longacre S, Escalante AA, Udagama-Randeniya PV 2011. Genetic diversity and recombination at the C-terminal fragment of the merozoite surface protein-1 of *Plasmodium vivax* (PvMSP-1) in Sri Lanka. *Infect Genet Evol* 11: 145-156.
- Escalante AA, Cornejo OE, Rojas A, Udhayakumar V, Lal AA 2004. Assessing the effect of natural selection in malaria parasites. *Trends Parasitol* 20: 388-395.
- Escalante AA, Lal AA, Ayala FJ 1998. Genetic polymorphism and natural selection in the malaria parasite *Plasmodium falciparum*. *Genetics* 149: 189-202.
- Farooq U, Malla N, Dubey ML 2009. Polymorphism in merozoite surface protein-1 gene in north and northwest Indian field isolates of *Plasmodium vivax*. *Indian J Med Res* 130: 736-741.
- Feng X, Carlton JM, Joy DA, Mu J, Furuya T, Suh BB, Wang Y, Barnwell JW, Su XZ 2003. Single-nucleotide polymorphisms and genome diversity in *Plasmodium vivax*. *Proc Natl Acad Sci USA* 100: 8502-8507.
- Ferreira MU, Karunaweera ND, da Silva-Nunes M, da Silva NS, Wirth DF, Hartl DL 2007. Population structure and transmission dynamics of *Plasmodium vivax* in rural Amazonia. *J Infect Dis* 195: 1218-1226.
- Figtree M, Pasay CJ, Slade R, Cheng Q, Cloonan N, Walker J, Saul A 2000. *Plasmodium vivax* synonymous substitution frequencies, evolution and population structure deduced from diversity in AMA 1 and MSP 1 genes. *Mol Biochem Parasitol* 108: 53-66.
- Galinski M, Barnwell J 2009. Monkey malaria kills four humans. *Trends Parasitol* 25: 200-204.
- Galinski MR, Corredor-Medina C, Povoia M, Crosby J, Ingravalle P, Barnwell JW 1999. *Plasmodium vivax* merozoite surface protein-3 contains coiled-coil motifs in an alanine-rich central domain. *Mol Biochem Parasitol* 101: 131-147.
- Gómez JC, McNamara T, Bockarie MJ, Baird JK, Carlton JM, Zimmerman PA 2003. Identification of a polymorphic *Plasmodium vivax* microsatellite marker. *Am J Trop Med Hyg* 69: 377-389.
- Gopinath R, Wongsrichanalai C, Córdón-Rosales C, Mirabelli L, Kyle D, Kain KC 1994. Failure to detect a *Plasmodium vivax*-like malaria parasite in globally collected blood samples. *J Infect Dis* 170: 1630-1633.
- Gosi P, Khushmith S, Khalambaheti T, Lanar D, Schaecher K, Fukuda M, Miller S 2008. Polymorphism patterns in Duffy-binding protein among Thai *Plasmodium vivax* isolates. *Malar J* 7: 112.
- Grimberg BT, Udomsangpetch R, Xainli J, McHenry A, Panichakul T, Sattabongkot J, Cui L, Bockarie M, Chitnis C, Adams J, Zimmerman PA, King CL 2007. *Plasmodium vivax* invasion of human erythrocytes inhibited by antibodies directed against the Duffy binding protein. *PLoS Med* 4: e337.
- Grynberg P, Fontes CJF, Hughes AL, Braga EM 2008. Polymorphism at the apical membrane antigen 1 locus reflects the world population history of *Plasmodium vivax*. *BMC Evolutionary Biology* 8: 123.
- Guerra CA, Howes RE, Patil AP, Gething PW, Van Boeckel TP, Temperley WH, Kabaria CW, Tatem AJ, Manh BH, Elyazar IR, Baird JK, Snow RW, Hay SI 2010. The international limits and population at risk of *Plasmodium vivax* transmission in 2009. *PLoS Negl Trop Dis* 4: e774.

- Gunasekera AM, Wickramarachchi T, Neafsey DE, Ganguli I, Perera L, Premaratne PH, Hartl D, Handunnetti SM, Udagama-Randeniya PV, Wirth DF 2007. Genetic diversity and selection at the *Plasmodium vivax* apical membrane antigen-1 (PvAMA-1) locus in a Sri Lankan population. *Mol Biol Evol* 24: 939-947.
- Gunawardena S, Karunaweera ND, Ferreira MU, Phone-Kyaw M, Pollack RJ, Alifrangis M, Rajakaruna RS, Konradsen F, Amerasinghe PH, Schousboe ML, Galappaththy GN, Abeyasinghe RR, Hartl DL, Wirth DF 2010. Geographic structure of *Plasmodium vivax*: microsatellite analysis of parasite populations from Sri Lanka, Myanmar, and Ethiopia. *Am J Trop Med Hyg* 82: 235-242.
- Gutierrez A, Vicini J, Patarroyo ME, Murillo LA, Patarroyo MA 2000. *Plasmodium vivax*: polymorphism in the merozoite surface protein 1 gene from wild Colombian isolates. *Exp Parasitol* 95: 215-219.
- Han ET, Park JH, Shin EH, Choi MH, Oh MD, Chai JY 2002. Apical membrane antigen-1 (AMA-1) gene sequences of re-emerging *Plasmodium vivax* in South Korea. *Korean J Parasitol* 40: 157-162.
- Han ET, Song TE, Park JH, Shin EH, Guk SM, Kim TY, Chai JY 2004. Allelic dimorphism in the merozoite surface protein-3alpha in Korean isolates of *Plasmodium vivax*. *Am J Trop Med Hyg* 71: 745-749.
- Han ET, Wang Y, Lim CS, Cho JH, Chai JY 2011. Genetic diversity of the malaria vaccine candidate merozoite surface protein 1 gene of *Plasmodium vivax* field isolates in Republic of Korea. *Parasitol Res* [Epub ahead of print]. Available from springerlink.com/content/x085hm23431201g1.
- Hans D, Pattnaik P, Bhattacharyya A, Shakri A, Yazdani S, Sharma M, Choe H, Farzan M, Chitnis C 2005. Mapping binding residues in the *Plasmodium vivax* domain that binds Duffy antigen during red cell invasion. *Mol Microbiol* 55: 1423-1434.
- Havryliuk T, Ferreira MU 2009. A closer look at multiple-clone *Plasmodium vivax* infections: detection methods, prevalence and consequences. *Mem Inst Oswaldo Cruz* 104: 67-73.
- Henry-Halldin CN, Sepe D, Susapu M, McNamara DT, Bockarie M, King CL, Zimmerman PA 2011. High-throughput molecular diagnosis of circumsporozoite variants VK210 and VK247 detects complex *Plasmodium vivax* infections in malaria endemic populations in Papua New Guinea. *Infect Genet Evol* 11: 391-398.
- Hernández-Martínez MA, Escalante AA, Arévalo-Herrera M, Herrera S 2011. Antigenic diversity of the *Plasmodium vivax* circumsporozoite protein in parasite isolates of Western Colombia. *Am J Trop Med Hyg* 84: 51-57.
- Herrera S, Corradin G, Arévalo-Herrera M 2007. An update on the search for a *Plasmodium vivax* vaccine. *Trends Parasitol* 23: 122-128.
- Hodder AN, Crewther PE, Anders RF 2001. Specificity of the protective antibody response to apical membrane antigen 1. *Infect Immun* 69: 3286-3294.
- Hoffmann EHE, Ribolla PEM, Ferreira MU 2003. Genetic relatedness of *Plasmodium falciparum* isolates and the origin of allelic diversity at the merozoite surface protein-1 (MSP-1) locus in Brazil and Vietnam. *Malar J* 2: 24.
- Holder AA 2009. The carboxy-terminus of merozoite surface protein 1: structure, specific antibodies and immunity to malaria. *Parasitology* 136: 1445-1456.
- Huang T, Cheng Q, Allan S, Huang Y 1994. DNA sequencing of circumsporozoite protein genes of *Plasmodium vivax* from four different countries in west Pacific region: comparative study on the flank sequences. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* 12: 85-92.
- Imwong M, Nair S, Pukrittayakamee S, Sudimack D, Williams JT, Mayxay M, Newton PN, Kim JR, Nandy A, Osorio L, Carlton JM, White NJ, Day NPJ, Anderson TJ 2007a. Contrasting genetic structure in *Plasmodium vivax* populations from Asia and South America. *Int J Parasitol* 37: 1013-1022.
- Imwong M, Pukrittayakamee S, Grüner AC, Rénia L, Letourneur F, Looareesuwan S, White NJ, Snounou G 2005. Practical PCR genotyping protocols for *Plasmodium vivax* using *Pvcs* and *Pvmsp1*. *Malar J* 4: 20.
- Imwong M, Pukrittayakamee S, Looareesuwan S, Poirriez J, Pasvol G, White NJ, Snounou G 2001. *Plasmodium vivax*: polymerase chain reaction amplification artifacts limit the suitability of *pvgam1* as a genetic marker. *Exp Parasitol* 99: 175-179.
- Imwong M, Snounou G, Pukrittayakamee S, Tanomsing N, Kim JR, Nandy A, Guthmann JP, Nosten F, Carlton J, Looareesuwan S, Nair S, Sudimack D, Day NP, Anderson TJ, White NJ 2007b. Relapses of *Plasmodium vivax* infection usually result from activation of heterologous hypnozoites. *J Infect Dis* 195: 927-933.
- Imwong M, Sudimack D, Pukrittayakamee S, Osório L, Carlton JM, Day NPJ, White NJ, Anderson TJ 2006. Microsatellite variation, repeat array length and population history of *Plasmodium vivax*. *Mol Biol Evol* 23: 1016-1018.
- Joshi H, Subbarao SK, Raghavendra K, Sharma VP 1989. *Plasmodium vivax*: enzyme polymorphism in isolates of Indian origin. *Trans R Soc Trop Med Hyg* 83: 179-181.
- Joy DA, Gonzalez-Cerón L, Carlton JM, Gueye A, Fay M, McCutchan TF, Su XZ 2008. Local adaptation and vector-mediated population structure in *Plasmodium vivax* malaria. *Mol Biol Evol* 25: 1245-1252.
- Kain KC, Brown AE, Mirabelli L, Webster HK 1993. Detection of *Plasmodium vivax* by polymerase chain reaction in a field study. *J Infect Dis* 168: 1323-1326.
- Karunaweera ND, Ferreira MU, Hartl DL, Wirth DF 2007. Fourteen polymorphic microsatellite DNA markers for the human malaria parasite *Plasmodium vivax*. *Mol Ecol Notes* 7: 172-175.
- Karunaweera ND, Ferreira MU, Munasinghe A, Barnwell JW, Collins WE, King CL, Kawamoto F, Hartl DL, Wirth DF 2008. Extensive microsatellite diversity in the human malaria parasite *Plasmodium vivax*. *Gene* 410: 105-112.
- Khatoun L, Baliraine FN, Bonizzoni M, Malik SA, Yan GY 2010. Genetic structure of *Plasmodium vivax* and *Plasmodium falciparum* in the Bannu district of Pakistan. *Malar J* 9: 112.
- Kho W, Chung J, Sim E, Kim D, Chung W 2001. Analysis of polymorphic regions of *Plasmodium vivax* Duffy binding protein of Korean isolates. *Korean J Parasitol* 39: 143-150.
- Kim JR, Imwong M, Nandy A, Chotivanich K, Nontprasert A, Tonomsing N, Maji A, Addy M, Day NP, White NJ, Pukrittayakamee S 2006. Genetic diversity of *Plasmodium vivax* in Kolkata, India. *Malar J* 5: 71.
- Kim SH, Hwang SY, Shin JH, Moon CS, Kim DW, Kho WG 2009. Molecular genetic characterization of the merozoite surface protein 1 gene of *Plasmodium vivax* from reemerging Korean isolates. *Clin Vaccine Immunol* 16: 733-738.
- Kimura E, Mattei D, di Santi SM, Scherf A 1990. Genetic diversity in the major merozoite surface antigen of *Plasmodium falciparum*: high prevalence of a third polymorphic form detected in strains derived from malaria patients. *Gene* 91: 57-62.
- Kirchgatter K, del Portillo HA 1998. Molecular analysis of *Plasmodium vivax* relapses using the MSP1 molecule as a genetic marker. *J Infect Dis* 177: 511-515.
- Koepfli C, Mueller I, Marfurt J, Goroti M, Sie A, Oa O, Genton B, Beck HP, Felger I 2009. Evaluation of *Plasmodium vivax* genotyping markers for molecular monitoring in clinical trials. *J Infect Dis* 199: 1074-1080.

- Kolakovich KA, Ssengoba A, Wojcik K, Tsuboi T, Al-Yaman F, Alpers M, Adams JH 1996. *Plasmodium vivax*: favored gene frequencies of the merozoite surface protein-1 and the multiplicity of infection in a malaria endemic region. *Exp Parasitol* 83: 11-18.
- Leclerc MC, Durant P, Gauthier C, Patot S, Billote N, Menegon M, Severini C, Ayala FJ, Renaud F 2004. Meager genetic variability of the human malaria agent *Plasmodium vivax*. *Proc Natl Acad Sci USA* 101: 14455-14460.
- Levinson G, Gutman GA 1987. Slipped-strand mispairing: a major mechanism for DNA sequence evolution. *Mol Biol Evol* 4: 203-221.
- Lim CS, Kim SH, Kwon SI, Song JW, Song KJ, Lee KN 2000. Analysis of *Plasmodium vivax* merozoite surface protein-1 gene sequences from resurgent Korean isolates. *Am J Trop Med Hyg* 62: 261-265.
- Lim CS, Tazi L, Ayala FJ 2005. *Plasmodium vivax*: recent world expansion and genetic identity to *Plasmodium simium*. *Proc Natl Acad Sci USA* 102: 15523-15528.
- Machado RLD, Póvoa MM 2000. Distribution of *Plasmodium vivax* variants (VK210, VK247 and *P. vivax*-like) in three endemic areas of the Amazon Region of Brazil and their correlation with chloroquine treatment. *Trans R Soc Trop Med Hyg* 94: 377-381.
- Maestre A, Sunil S, Ahmad G, Mohammed A, Echeverri M, Corredor M, Blair S, Chauhan VS, Malhotra P 2004. Inter-allelic recombination in the *Plasmodium vivax* merozoite surface protein 1 gene among Indian and Colombian isolates. *Malar J* 3: 4.
- Mann VH, Huang T, Cheng Q, Saul A 1994. Sequence variation in the circumsporozoite protein gene of *Plasmodium vivax* appears to be regionally biased. *Mol Biochem Parasitol* 68: 45-52.
- Martinez P, Suarez CF, Cardenas PP, Patarroyo MA 2004. *Plasmodium vivax* Duffy binding protein: a modular evolutionary proposal. *Parasitology* 128: 353-366.
- Mascorro CN, Zhao K, Khuntirat B, Sattabongkot J, Yan G, Escalante AA, Cui L 2005. Molecular evolution and intragenic recombination of the merozoite surface protein MSP-3 alpha from the malaria parasite *Plasmodium vivax* in Thailand. *Parasitology* 131: 25-35.
- McConkey GA, Waters AP, McCutchan TF 1990. The generation of genetic diversity in malaria parasites. *Annu Rev Microbiol* 44: 479-498.
- Michon P, Fraser T, Adams J 2000. Naturally acquired and vaccine-elicited antibodies block erythrocyte cytoadherence of the *Plasmodium vivax* Duffy binding protein. *Infect Immun* 68: 3164-3171.
- Monette M, Opella SJ, Greenwood J, Willis AE, Perham RN 2001. Structure of a malaria parasite antigenic determinant displayed on filamentous bacteriophage determined by NMR spectroscopy: implications for the structure of continuous peptide epitopes of proteins. *Protein Sci* 10: 1150-1159.
- Moon SU, Lee HW, Kim JY, Na BK, Cho SH, Lin K, Sohn WM, Kim TS 2009. High frequency of genetic diversity of *Plasmodium vivax* field isolates in Myanmar. *Acta Trop* 109: 30-36.
- Moon SU, Na BK, Kang JM, Kim JY, Cho SH, Park YK, Sohn WM, Lin K, Kim TS 2010. Genetic polymorphism and effect of natural selection at domain I of apical membrane antigen-1 (AMA-1) in *Plasmodium vivax* isolates from Myanmar. *Acta Tropica* 114: 71-75.
- Mueller I, Galinski MR, Baird JK, Carlton JM, Kochar DK, Alonso PL, del Portillo HA 2009. Key gaps in the knowledge of *Plasmodium vivax*, a neglected human malaria parasite. *Lancet Infect Dis* 9: 555-566.
- Mueller I, Kaiok J, Reeder JC, Cortés A 2002. The population structure of *Plasmodium falciparum* and *Plasmodium vivax* during an epidemic of malaria in the Eastern Highlands of Papua New Guinea. *Am J Trop Med Hyg* 67: 459-464.
- Nardin EH, Zavala F 1998. Acquired immunity to sporozoites. In IW Sherman, *Malaria: parasite biology, pathogenesis, and protection*, ASM Press, Washington DC, p. 495-511.
- Nogueira PA, Alves FP, Fernandez-Becerra C, Pein O, Santos NR, Pereira da Silva LH, Camargo EP, del Portillo HA 2006. A reduced risk of infection with *Plasmodium vivax* and clinical protection against malaria are associated with antibodies against the N terminus but not the C terminus of merozoite surface protein 1. *Infect Immun* 74: 2726-2733.
- Oliveira-Ferreira J, Lacerda MV, Brasil P, Ladislau JL, Taulil PL, Daniel-Ribeiro CT, 2010. Malaria in Brazil: an overview. *Malar J* 9: 115.
- Ord R, Polley S, Tami A, Sutherland CJ 2005. High sequence diversity and evidence of balancing selection in the *PvMSP3 alpha* gene of *Plasmodium vivax* in the Venezuelan Amazon. *Mol Biochem Parasitol* 144: 86-93.
- Ord RL, Tami A, Sutherland CJ 2008. *ama1* genes of sympatric *Plasmodium vivax* and *P. falciparum* from Venezuela differ significantly in genetic diversity and recombination frequency. *PLoS ONE* 3: e3366.
- Orjuela-Sánchez P, da Silva NS, da Silva-Nunes M, Ferreira MU 2009a. Recurrent parasitemias and population dynamics of *Plasmodium vivax* polymorphisms in rural Amazonia. *Am J Trop Med Hyg* 81: 961-968.
- Orjuela-Sánchez P, da Silva-Nunes M, da Silva NS, Scopel KK, Gonçalves RM, Malafronte RS, Ferreira MU 2009b. Population dynamics of genetically diverse *Plasmodium falciparum* lineages: community-based prospective study in rural Amazonia. *Parasitology* 136: 1097-1105.
- Pacheco MA, Poe AC, Collins WE, Lal AA, Tanabe K, Kariuki SK, Udhayakumar V, Escalante AA 2007. A comparative study of the genetic diversity of the 42 kDa fragment of the merozoite surface protein 1 in *Plasmodium falciparum* and *P. vivax*. *Infect Genet Evol* 7: 180-187.
- Pasay MC, Cheng Q, Rzepczyk C, Saul A 1995. Dimorphism of the C terminus of the *Plasmodium vivax* merozoite surface protein 1. *Mol Biochem Parasitol* 70: 217-219.
- Patil A, Orjuela-Sánchez P, da Silva-Nunes M, Ferreira MU 2010. Evolutionary dynamics of the immunodominant repeats of the *Plasmodium vivax* malaria-vaccine candidate circumsporozoite protein (CSP). *Infect Genet Evol* 10: 298-303.
- Pizarro JC, Vulliez-Le Normand B, Chesne-Seck ML, Collins CR, Withers-Martinez C, Hackett F, Blackman MJ, Faber BW, Remarque EJ, Kocken CH, Thomas AW, Bentley GA 2005. Crystal structure of the malaria vaccine candidate apical membrane antigen 1. *Science* 308: 408-411.
- Polley SD, Weedall GD, Thomas AW, Golightly LM, Conway DJ 2005. Orthologous gene sequences of merozoite surface protein 1 (MSP1) from *Plasmodium reichenowi* and *P. gallinaceum* confirm an ancient divergence of *P. falciparum* alleles. *Mol Biochem Parasitol* 142: 25-31.
- Prajapati SK, Joshi H, Valecha N 2010. *Plasmodium vivax* merozoite surface protein-3 alpha: a high-resolution marker for genetic diversity studies. *J Vec Bor Dis* 472: 85-90.
- Prajapati SK, Verma A, Adak T, Yadav RS, Kumar A, Eapen A, Das MK, Singh N, Sharma SK, Rizvi MA, Dash AP, Joshi H 2006. Allelic dimorphism of *Plasmodium vivax* gam-1 in the Indian subcontinent. *Malar J* 5: 90.
- Premawansa S, Snewin VA, Khouri E, Mendis KN, David PH 1993. *Plasmodium vivax*: recombination between potential allelic types of the merozoite surface protein MSP1 in parasites isolated from patients. *Exp Parasitol* 76: 192-199.

- Price RN, Douglas NM, Anstey NM 2009. New developments in *Plasmodium vivax* malaria: severe disease and the rise of chloroquine resistance. *Curr Opin Infect Dis* 22: 430-435.
- Putaporntip C, Jongwutiwes S, Grynberg P, Cui L, Hughes AL 2009. Nucleotide sequence polymorphism at the apical membrane antigen-1 locus reveals population history of *Plasmodium vivax* in Thailand. *Infect Gen Evol* 9: 1295-1300.
- Putaporntip C, Jongwutiwes S, Iwasaki T, Kanbara H, Hughes AL 2006. Ancient common ancestry of the merozoite surface protein 1 of *Plasmodium vivax* as inferred from its homologue in *Plasmodium knowlesi*. *Mol Biochem Parasitol* 146: 105-108.
- Putaporntip C, Jongwutiwes S, Sakihama N, Ferreira M, Kho W, Kaneko A, Kanbara H, Hattori T, Tanabe K 2002. Mosaic organization and heterogeneity in frequency of allelic recombination of the *Plasmodium vivax* merozoite surface protein-1 locus. *Proc Natl Acad Sci USA* 99: 16348-16353.
- Putaporntip C, Jongwutiwes S, Tanabe K, Thaithong S 1997. Inter-allelic recombination in the merozoite surface protein (MSP-1) gene of *Plasmodium vivax* from Thai isolates. *Mol Biochem Parasitol* 84: 49-56.
- Qari SH, Collins WE, Lobel HO, Taylor F, Lal AA 1994. A study of polymorphism in the circumsporozoite protein of human malaria parasites. *Am J Trop Med Hyg* 50: 45-51.
- Qari SH, Goldman IF, Pova MM, di Santi S, Alpers MP, Lal AA 1992. Polymorphism in the circumsporozoite protein of the human malaria parasite *Plasmodium vivax*. *Mol Biochem Parasitol* 55: 105-113.
- Qari SH, Goldman IF, Pova MM, Oliveira S, Alpers MP, Lal AA 1991. Wide distribution of the variant form of the human malaria parasite *Plasmodium vivax*. *J Biol Chem* 266: 16297-16300.
- Qari SH, Shi YP, Goldman IF, Udhayakumar V, Alpers MP, Collins WE, Lal AA 1993a. Identification of *Plasmodium vivax*-like human malaria parasite. *Lancet* 341: 780-783.
- Qari SH, Shi YP, Pova MM, Alpers MP, Deloron P, Murphy GS, Harjowarno S, Lal AA 1993b. Global occurrence of *Plasmodium vivax*-like human malaria parasite. *J Infect Dis* 168: 1485-1489.
- Rajesh V, Elamaran M, Vidya S, Gowrishankar M, Kochar D, Das A 2007. *Plasmodium vivax*: genetic diversity of the apical membrane antigen-1 (AMA-1) in isolates from India. *Exp Parasitol* 116: 252-256.
- Ranjan A, Chitnis C 1999. Mapping regions containing binding residues within functional domains of *Plasmodium vivax* and *Plasmodium knowlesi* erythrocyte-binding proteins. *Proc Natl Acad Sci USA* 96: 14067-14072.
- Rezende AM, Tarazona-Santos E, Couto AD, Fontes CJ, de Souza JM, Carvalho LH, Brito CF 2009. Analysis of genetic variability of *Plasmodium vivax* isolates from different Brazilian Amazon areas using tandem repeats. *Am J Trop Med Hyg* 80: 729-733.
- Rezende AM, Tarazona-Santos E, Fontes CJ, Souza JM, Couto AD, Carvalho LH, Brito CF 2010. Microsatellite loci: determining the genetic variability of *Plasmodium vivax*. *Trop Med Int Health* 15: 718-726.
- Rich SM, Ferreira MU, Ayala FJ 2000. The origin of antigenic diversity in *Plasmodium falciparum*. *Parasitol Today* 16: 390-396.
- Rich SM, Hudson RR, Ayala FJ 1997. *Plasmodium falciparum* antigenic diversity: evidence of clonal population structure. *Proc Natl Acad Sci USA* 94: 13040-13045.
- Rodrigues MH, Rodrigues KM, Oliveira TR, Cômido AN, Rodrigues MM, Kocken CH, Thomas AW, Soares IS 2005. Antibody response of naturally infected individuals to recombinant *Plasmodium vivax* apical membrane antigen-1. *Int J Parasitol* 35: 185-192.
- Rongnoparut P, Supsamran N, Sattabongkot J, Suwanabun N, Rosenberg R 1995. Phenotype and genotype diversity in the circumsporozoite proteins of *Plasmodium vivax* in Thailand. *Mol Biochem Parasitol* 74: 201-210.
- Rosenberg R, Wirtz RA, Lanar DE, Sattabongkot J, Hall T, Waters AP, Prasittisuk C 1989. Circumsporozoite protein heterogeneity in the human malaria parasite *Plasmodium vivax*. *Science* 245: 973-976.
- Roy SW, Ferreira MU, Hartl DL 2008. Evolution of allelic dimorphism in malarial surface antigens. *Heredity* 100: 103-110.
- Santos-Ciminera PD, Alecrim MG, Roberts DR, Quinnan GV Jr 2007. Molecular epidemiology of *Plasmodium vivax* in the state of Amazonas, Brazil. *Acta Trop* 102: 38-46.
- Schlötterer C 1998. Genome evolution: are microsatellites really simple sequences? *Curr Biol* 8: R132-134.
- Silvie O, Franetich JF, Charrin S, Mueller MS, Siau A, Bodescot M, Rubinstein E, Hannoun L, Charoenvit Y, Kocken CH, Thomas AW, Gemert GJv, Sauerwein RW, Blackman MJ, Anders RF, Pluschke G, Maziera D 2004. Role for apical membrane antigen 1 during invasion of hepatocytes by *Plasmodium falciparum* sporozoites. *J Biol Chem* 279: 9490-9496.
- Singh AP, Ozwara H, Kocken CHM, Puri SK, Thomas AW, Chitnis CE 2005. Targeted deletion of *Plasmodium knowlesi* Duffy binding protein confirms its role in junction formation during invasion. *Mol Microbiol* 55: 1925-1934.
- Snewin VA, Khouri E, Wattavidanage J, Perera L, Premawansa S, Mendis KN, David PH 1995. A new polymorphic marker for PCR typing of *Plasmodium vivax* parasites. *Mol Biochem Parasitol* 71: 135-138.
- Soares IS, Barnwell JW, Ferreira MU, Cunha MG, Laurino JP, Castilho BA, Rodrigues MM 1999. A *Plasmodium vivax* vaccine candidate displays limited allele polymorphism, which does not restrict recognition by antibodies. *Mol Medicine* 5: 459-470.
- Sousa T, Cerávolo I, Fernandes Fontes C, Couto A, Carvalho L, Brito C 2006. The pattern of major polymorphisms in the Duffy binding protein ligand domain among *Plasmodium vivax* isolates from the Brazilian Amazon area. *Mol Biochem Parasitol* 146: 251-254.
- Sousa TN, Tarazona-Santos EM, Wilson DJ, Madureira AP, Falcão PR, Fontes CJ, Gil LH, Ferreira MU, Carvalho LH, Brito CF 2010. Genetic variability and natural selection at the ligand domain of the Duffy binding protein in Brazilian *Plasmodium vivax* populations. *Malar J* 22: 334.
- Storti-Melo LM, de Souza-Neiras WC, Cassiano GC, Joazeiro AC, Fontes CJ, Bonini-Domingos CR, Couto AA, Pova MM, de Mattos LC, Cavasini CE, Rossit AR, Machado RL 2009. *Plasmodium vivax* circumsporozoite variants and Duffy blood group genotypes in the Brazilian Amazon Region. *Trans R Soc Trop Med Hyg* 103: 672-678.
- Suh IB, Hoffman KJ, Kim SH, Song KJ, Song JW, Lee JS, Lim CS 2001. The analysis of *Plasmodium vivax* Duffy receptor binding domain gene sequence from resurgent Korea isolates. *Parasitol Res* 87: 1007-1010.
- Sutton PL, Neyra V, Hernandez JN, Branch OH 2009. *Plasmodium falciparum* and *Plasmodium vivax* infections in the Peruvian Amazon: propagation of complex, multiple allele-type infections without super-infection. *Am J Trop Med Hyg* 816: 950-960.
- Thakur A, Alam MT, Bora H, Kaur P, Sharma YD 2008. *Plasmodium vivax*: sequence polymorphism and effect of natural selection at apical membrane antigen 1 (PvAMA1) among Indian population. *Gene* 419: 35-42.
- Tsuboi T, Kappe SHI, Alyaman F, Prickett MD, Alpers M, Adams JH 1994. Natural variation within the principal adhesion domain of the *Plasmodium vivax* Duffy binding-protein. *Infect Immun* 62: 5581-5586.

- Udagama PV, David PH, Peiris JSM, Ariyaratne YG, Perera KLRL, Mendis KN 1987. Demonstration of antigenic polymorphism in *Plasmodium vivax* malaria with a panel of 30 monoclonal antibodies. *Infect Immun* 55: 2604-2611.
- Udagama PV, Gamage-Mendis AC, David PH, Peiris JS, Perera KLRL, Carter R 1990. Genetic complexity of *Plasmodium vivax* parasites in individual human infections analyzed with monoclonal antibodies against variant epitopes on a single parasite protein. *Am J Trop Med Hyg* 42: 101-110.
- Udomsangpetch R, Kaneko O, Chotivanich K, Sattabongkot J 2008. Cultivation of *Plasmodium vivax*. *Trends Parasitol* 24: 85-88.
- Valderrama-Aguirre A, Zúñiga-Soto E, Mariño-Ramírez L, Moreno LA, Escalante AA, Arévalo-Herrera M, Herrera S 2011. Polymorphism of the Pv200L fragment of merozoite surface protein-1 of *Plasmodium vivax* in clinical isolates from the Pacific coast of Colombia. *Am J Trop Med Hyg* 84: 64-70.
- Van den Eede P, Erhart A, Van der Auwera G, Van Overmeir C, Thang ND, Hung le X, Anné J, D'Alessandro U 2010a. High complexity of *Plasmodium vivax* infections in symptomatic patients from a rural community in central Vietnam detected by microsatellite genotyping. *Am J Trop Med Hyg* 82: 223-227.
- Van den Eede P, Soto-Calle VE, Delgado C, Gamboa D, Grande T, Rodríguez H, Llanos-Cuentas A, Anné J, D'Alessandro U, Erhart A 2011. *Plasmodium vivax* sub-patent infections after radical treatment are common in Peruvian patients: results of a 1-year prospective cohort study. *PLoS ONE* 6: e16257.
- Van den Eede P, Van der Auwera G, Delgado C, Huysse T, Soto-Calle VE, Gamboa D, Grande T, Rodríguez H, Llanos A, Anné J, Erhart A, D'Alessandro U 2010b. Multilocus genotyping reveals high heterogeneity and strong local population structure of the *Plasmodium vivax* population in the Peruvian Amazon. *Malar J* 9: 151.
- VanBuskirk KM, Cole-Tobian JL, Baisor M, Sevova ES, Bockarie M, King CL, Adams JH 2004. Antigenic drift in the ligand domain of *Plasmodium vivax* Duffy binding protein confers resistance to inhibitory antibodies. *J Infect Dis* 190: 1556-1562.
- Veron V, Legrand E, Yrinesi J, Volney B, Simon S, Carme B 2009. Genetic diversity of msp3 and msp1_b5 markers of *Plasmodium vivax* in French Guiana. *Mal J* 8: 40.
- Xainli J, Adams J, King C 2000. The erythrocyte binding motif of *Plasmodium vivax* duffy binding protein is highly polymorphic and functionally conserved in isolates from Papua New Guinea. *Mol Biochem Parasitol* 111: 253-260.
- Yang C, Collins WE, Sullivan JS, Kaslow DC, Xiao L, Lal AA 1999. Partial protection against *Plasmodium vivax* blood-stage infection in Saimiri monkeys by immunization with a recombinant C-terminal fragment of merozoite surface protein 1 in block copolymer adjuvant. *Infect Immun* 67: 342-349.
- Zakeri S, Barjesteh H, Djadid ND 2006. Merozoite surface protein-3 alpha is a reliable marker for population genetic analysis of *Plasmodium vivax*. *Mal J* 5: 53.
- Zakeri S, Raeisi A, Afsharpad M, Kakar Q, Ghasemi F, Atta H, Zamani G, Memon MS, Salehi M, Djadid ND 2010. Molecular characterization of *Plasmodium vivax* clinical isolates in Pakistan and Iran using *PvMSP-1*, *PvMSP-3 alpha* and *pvcsp* genes as molecular markers. *Parasitol Int* 59: 15-21.
- Zeyrek FY, Tachibana S, Yuksel F, Doni N, Palacpac N, Arisue N, Horii T, Coban C, Tanabe K 2010. Limited polymorphism of the *Plasmodium vivax* merozoite surface protein 1 gene in isolates from Turkey. *Am J Trop Med Hyg* 83: 1230-1237.
- Zilversmit M, Hartl DL 2005. Evolutionary history and population genetics of human malaria parasites. In IW Sherman (ed.), *Molecular approaches to malaria*, Press, Washington DC, 542 pp.