

New triterpene and triterpenoid glycosides from *Ilex brevicuspis*

Alexandre T. Cardoso Taketa¹, Manfred Gondorf¹, Eberhard Breitmaier¹, Eloir Paulo Schenkel^{2*}

¹Kekulé-Institut für Organische Chemie und Biochemie der Universität Bonn – Germany, ²Departamento de Ciências Farmacêuticas, Centro de Ciências da Saúde, Universidade Federal de Santa Catarina

***Correspondence:**

E.P. Schenkel
Departamento de Ciências
Farmacêuticas - Centro de Ciências da
Saúde
Universidade Federal de Santa
Catarina -Campus Universitário /
Trindade
88.040-900 – Florianópolis-SC, Brasil
E-mail: schenkel@ccs.ufsc.br

From the leaves of *Ilex brevicuspis* were isolated a new triterpene, 20(S)-3 β ,19 α -dihydroxyurs-12-en-23,28-dioic acid, named here *brevicuspis acid*, and two new triterpenoid glycosides, 3-O- α -L-arabinopyranosyl-20(S)-pomolic acid-28-O- β -D-glucopyranosyl ester, named *brevicuspisaponin 3*, and the 23-sodium salt of (20S)-3 β ,19 α ,24-trihydroxyurs-12-en-23,28-dioic acid-28 β -O- β -D-glucopyranosyl ester, along with the known compound 3-O- α -L-arabinopyranosyl-20(S)-19 α ,24-dihydroxyursolic acid-28-O- β -D-glucopyranosyl ester, already described for *Ilex argentina*. Their structures were established on the basis of chemical and spectroscopic methods.

Uniterms:

- Aquifoliaceae
- *Ilex brevicuspis*
- Triterpenes
- Saponins
- Brevicuspis acid
- Brevicuspisaponins 3 and 4

INTRODUCTION

As previously reported, the leaves of *Ilex brevicuspis* Reissek (Aquifoliaceae) afforded the triterpenes ursolic acid, 23-methylester of 20(S)-rotundioic acid, and the new glycosides *brevicuspisaponin 1* and *2* (Taketa *et al.*, 2000). Continuing the investigation on the leaves of *I. brevicuspis*, as a part of our studies on the adulterants of the genuine maté, *Ilex paraguariensis* (Taketa, Schenkel., 1994; Schenkel *et al.*, 1995; Heinzmann, Schenkel, 1995; Pires *et al.*, 1997; Schenkel *et al.*, 1997; Athayde *et al.*, 1999; Reginatto *et al.*, 1999; Athayde *et al.*, 2001), we report here further new triterpene and saponins from the title plant.

MATERIAL AND METHODS


Plant material

Leaves of *Ilex brevicuspis* Reissek were collected in Osório, State of Rio Grande do Sul, Brazil. A herbarium specimen (leg. Coelho 163) is on deposit in the Herbarium of the Botany Department of the Federal University of Rio

Grande do Sul (Herbarium ICN, Porto Alegre, Brazil).

General experimental procedures

Melting points were obtained in a Kofler melting point apparatus and are uncorrected. IR spectra were recorded in a Perkin-Elmer 881 spectrophotometer. Optical rotations were measured in a Perkin-Elmer 241 polarimeter. EIMS and HRMS spectra were performed in a MS50 spectrometer and FAB-MS spectra on a Concept 1H spectrometer. ¹H and ¹³C NMR spectra were recorded in Bruker AMX and DRX 500 spectrometers. TLC were carried out on silica gel (Merck) GF₂₅₄ and sugars and glycosides were eluted with EtOAc/MeOH/HOAc/H₂O 13:3:4:3. All compounds were visualized using the vanillin-sulfuric acid reagent/100 °C/10 min. Open CC were performed on a normal phase silica gel 40-60 mm using the eluants CHCl₃/MeOH 97:3 and cyclohexane/acetone 1:1 to **1a** and **2a**, CHCl₃/MeOH 9:1 to **1** and CHCl₃/EtOH/H₂O 8:4:0.5 to **2**, **3** and **4**. For **3**, the phase LiChroprep C-18, 40-63 mm, the eluant MeOH/H₂O 3:1 and Sephadex LH-20 in MeOH were used for final purifications.



Extraction and isolation

Air-dried leaves (589 g) were crushed and extracted with ethanol (1.5 L) at room temperature (2 x 7 days). The ethanolic extract (2 L) was evaporated to dryness under reduced pressure and the residue (69 g) was then suspended in water (700 mL) and extracted with chloroform (5 x 500 mL). Between the water and chloroform phases an emulsified phase was formed. The emulsified phase was evaporated to dryness to give the fraction containing the saponins (19 g). Part of this residue was repeatedly chromatographed to give compounds **1** (17 mg), **2** (13 mg) and pure compounds **3** (26 mg) and **4** (17 mg).

Acid hydrolysis

Compounds **2**, **3** and **4** were hydrolyzed on TLC plates in order to identify their sugars, as described by Kartnig and Wegschaidner (1972).

Acetylation of compounds 1, 2 and 4

Compounds **1** (3 mg), **2** (3 mg) and **4** (6 mg) were acetylated with acetic anhydride/pyridine (1:1) at room temperature overnight, affording peracetylated compounds **1a**, **2a** and **4a**.

Acetylated derivative (**1a**):

Acetylated brevicuspis acid [20(S)-3β,19α-dihydroxyurs-12-en-23,28-dioic acid] (1)

White powder, mp 232-235 °C. $[\alpha]_{589}^{20} +33^\circ$, $[\alpha]_{578}^{20}$

$+80^\circ$, $[\alpha]_{546}^{20} +94^\circ$ and $[\alpha]_{436}^{20} +182^\circ$ (MeOH, *c* 0.17). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3425, 2932, 1741, 1698, 1456, 1372, 1237, 1030. FAB-MS (positive-ion mode, mNBA) *m/z*: 567.3 $[\text{M} + \text{Na}]^+$, 467.2, 421.3, 338.3. EIMS *m/z*: 544.4 $[\text{M}]^+$, 426.4, 498.4, 482.4, 438.3, 426.4, 407.3. HRMS: 526.3291 $[\text{M}-\text{H}_2\text{O}]^+$. ¹H NMR (Table I) and ¹³C NMR (Table II).

Brevicuspisaponin 3(2):

[3-O-α-L-arabinopyranosyl-20(S)-pomolic acid-28-O-β-D-glucopyranosyl ester]

FAB-MS (positive-ion mode, mNBA) *m/z*: 789.4 $[\text{M} + \text{Na}]^+$, 703.4, 687.4, 613.1, 599.1, 531.2.

Peracetylated derivative (**2a**):

White powder, mp 153-155 °C. $[\alpha]_{589}^{20} +17^\circ$, $[\alpha]_{578}^{20} +21^\circ$, $[\alpha]_{546}^{20} +23^\circ$, $[\alpha]_{436}^{20} +39^\circ$ and $[\alpha]_{365}^{20} +61^\circ$ (CHCl_3 , *c* 0.21). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3481, 2946, 1752, 1370, 1224, 1056. FAB-MS (positive-ion mode, mNBA) *m/z*: 1083.4 $[\text{M} + \text{Na}]^+$, 904.4, 850.4, 783.3, 683.4, 667.2, 613.3. ¹H NMR (Table I) and ¹³C NMR (Table II).

ILA-1(3):

[3β-O-α-L-arabinopyranosyl-20(S)-19α,24-dihydroxyursolic acid-28-O-β-D-glucopyranosyl ester]

White powder, mp 200-204 °C $[\alpha]_{589}^{20} +12^\circ$, $[\alpha]_{578}^{20} +13^\circ$, $[\alpha]_{546}^{20} +14^\circ$, $[\alpha]_{436}^{20} +24^\circ$ and $[\alpha]_{365}^{20} +37^\circ$ (MeOH, *c* 0.25). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3437, 2932, 1734, 1637, 1382, 1074. FAB-MS (positive mode, mNBA) *m/z*: 805.3 $[\text{M} + \text{Na}]^+$, 642, 482, 411. ¹H NMR (Table I) and ¹³C NMR (Table II).

Brevicuspisaponin 4 (**4**):

[23-sodium salt of (20S)-3 β ,19 α ,24-trihydroxyurs-12-en-23,28-dioic acid-28 β -O- β -D-glucopyranosyl ester]

FABMS (positive-ion mode, thioglycerol) m/z : 725.3 [M+Na]⁺, 703.3 [M+H]⁺, 539.2 [M-C₆H₁₁O₅]. IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 3422, 2930, 1735, 1573, 1457, 1383, 1232, 1073.

Peracetylated derivative (**4a**):

White powder, mp 153-155 °C. FAB-MS (positive-ion mode, mNBA) m/z : 955.4 [M+H]⁺ ¹H NMR (Table I) and ¹³C NMR (Table II).

RESULTS AND DISCUSSION

Triterpene **1** and saponins **2**, **3** and **4** were isolated from the emulsified phase obtained from the leaves of the title plant, using the procedure described in the experimental section. Due to difficulty in purifications, compounds **1**, **2** and **4** were acetylated in order to afford pure compounds **1a**, **2a** and **4a**.

EIMS spectrum of **1a** display a molecular peak at m/z 544 [M⁺] as molecular ion, which together with the HRMS from the fragment ion with m/z 526.3291 [M⁺-H₂O] suggested the molecular formula C₃₂H₄₈O₇. The triterpenoid structure was confirmed by the ¹H broadband-decoupled ¹³C NMR experiment which exhibited the presence of one acetyl group (δ_{C} 170.2 and 21.1) and thirty other carbon atoms. The DEPT subspectra revealed the presence of six methyl, nine methylene, six methane groups. It was possible to recognize the existence of two carboxy groups (δ_{C} 180.7 and 179.2), one double bond (δ_{C} 126.8 and 139.5), one acetylated *sec* alcohol group (δ_{C} 78.5) and one *tert* alcohol group (δ_{C} 73.3). The ¹H NMR spectrum showed the presence of five angular methyl groups (δ_{H} 0.94, 1.05, 1.42, 1.52 and 1.68) and one methyl group attached to CH (δ_{H} 1.10, *d*, ³*J* = 7.0 Hz), suggesting an ursane or oleanane derivative. The multiplicities at δ_{H} 5.76 (3 α -H, *dd*, ³*J* = 12.0 and 3.4 Hz) indicated a 3 β -hydroxy substitution on this skeleton. HMBC experiment allowed to locate 24-CH₃ (δ_{C} 12.8) and 23-COOH (δ_{C} 179.2) at C-4. NOE enhancements were detected for 25 β -CH₃ (δ_{H} 0.94, *s*) and 24-CH₃ (δ_{H} 1.52, *s*) in a ROESY experiment, indicating the 4 α -configuration of the carboxy group.

In the ¹H NMR spectrum, the deshielded methylene hydrogen proton at δ_{H} 3.19 showed characteristic multiplicities (*td*, *J* = 12.9 and 3.8 Hz) that established an axial relative stereochemistry for 16 α -H. It was confirmed through the *W*-type correlation observed between proton signals at δ_{H} 2.02 (equatorial 16 β -H) and δ_{H} 3.26 (equato-

TABLE I - ¹³C-NMR data for the acetylated derivatives **1a**, **2a** and **4a**, and compound **3** (pyridine-d₅)

C	1a*	2a*	3	4a
1	38.2	38.7	38.7	38.8
2	23.5	26.7	27.0	24.0
3	78.5	89.5	89.2	77.9
4	52.1	39.4	44.4	54.9
5	52.0	55.9	56.4	52.6
6	21.3	18.7	19.1	23.1
7	33.2	33.8	33.8	34.2
8	40.5	40.4	40.5	40.6
9	47.8	47.8	47.8	48.1
10	36.7	37.2	36.9	37.0
11	23.9	24.1	24.8	24.2
12	126.8	128.1	127.7	127.7
13	139.5	138.4	139.0	138.4
14	42.1	42.3	42.3	42.1
15	29.2	29.3	29.3	29.2
16	27.0	26.8	26.9	26.6
17	47.9	48.6	48.5	48.6
18	47.4	47.2	47.3	47.1
19	73.3	73.4	73.5	73.3
20	43.2	42.9	43.0	42.9
21	24.9	24.7	24.4	24.7
22	32.4	31.8	32.0	31.7
23	179.2	28.1	23.6	176.5
24	12.8	17.0	63.5	63.9
25	15.8	15.7	15.6	15.5
26	17.1	17.5	17.5	17.1
27	24.3	24.3	24.7	24.1
28	180.7	176.5	177.2	176.4
29	29.8	29.8	30.0	29.6
30	16.1	16.1	16.2	16.0
Ara-1'		104.0	106.7	
Ara-2'		70.5	73.0	
Ara-3'		71.5	74.7	
Ara-4'		69.1	69.6	
Ara-5'		64.1	66.9	
Glc-1''		92.5	96.0	92.4
Glc-2''		71.1	74.3	71.0
Glc-3''		73.6	79.1	73.6
Glc-4''		69.0	71.2	69.0
Glc-5''		73.2	79.5	73.1
Glc-6''		62.3	62.3	62.3

*Values for the acetyl groups are not recorded in the table. Acetyl groups: $\delta_{\text{C}} \cong 20$ (CH₃) and $\delta_{\text{C}} \cong 168$ (C=O).

TABLE II - $^1\text{H-NMR}$ data for the acetylated derivatives **1a**, **2a** and **4a**, and compound **3** (pyridine- d_5) (multiplicities; $J = \text{Hz}$)

H	1a*	2a*	3	4a
1	1.09/ α - 1.53/ β	0.91/ α - 1.54/ β	0.91/ α - 1.52/ β	1.12/ α - 1.62/ β
2	1.71/ β - 1.92/ α	1.84/ β - 2.04/ α	2.03/ β - 2.18 (<i>td</i> ; 13.8, 4.0)/ α	1.12/ β - 1.86/ α
3	5.76 (<i>dd</i> ; 12.0, 3.4)/ α	3.23/ α	3.50 (<i>dd</i> ; 11.6, 4.8)/ α	5.77/ α
5	1.99/ α	0.82/ α	0.95/ α	2.18/ α
6	1.54/ α - 1.65/ β	1.37/ β - 1.55/ α	1.37/ β - 1.61/ α	1.94 - 2.10
7	1.30 - 1.68	1.34/ β - 1.59/ α	1.44/ β - 1.54/ α	1.37/ β - 1.70/ α
9	1.91/ α	1.79/ α	1.77/ α	1.91/ α
11	2.00 - 2.00	2.00 - 2.00	1.93 - 1.98	2.03 - 2.03
12	5.54	5.50	5.49	5.49
15	1.19/ α - 2.22 (<i>td</i> ; 13.5, 3.8)/ β	1.24/ α - 1.87/ β	1.26/ α - 2.47 (<i>td</i> ; 13.6, 4.1)/ β	1.15/ α - 1.64/ β
16	2.02/ β - 3.19 (<i>td</i> ; 12.9, 3.8)/ α	1.87/ β - 3.16 (<i>td</i> ; 11.6, 4.3)/ α	2.09/ β - 3.23 (<i>td</i> ; 13.1, 4.3)/ α	1.82/ β - 3.12 (<i>td</i> ; 11.8, 3.5)/ α
18	3.26 (<i>s</i>)/ β	3.03 (<i>s</i>)/ β	3.18 (<i>s</i>)/ β	3.03 (<i>s</i>)/ β
20	1.99/ α	1.92/ α	1.94/ α	1.90/ α
21	1.30/ β - 2.68 (<i>tt</i> ; 13.5, 3.3)/ α	1.18/ β - 2.57 (<i>tt</i> ; 13.3, 4.2)/ α	1.17/ β - 2.59 (<i>tt</i> ; 13.6, 3.9)/ α	1.17/ β - 2.55 (<i>tt</i> ; 13.2, 3.5)/ α
22	1.93/ α - 2.22 (<i>td</i> ; 13.5, 3.8)/ β	1.80/ α - 1.98/ β	1.90/ α - 2.09/ β	1.78/ α - 1.97/ β
23	-	1.08 (<i>s</i>)	1.50 (<i>s</i>)	-
24	1.52 (<i>s</i>)	0.94 (<i>s</i>)	3.59 (<i>d</i> ; 11.1) - 4.37 (<i>ov.</i>)	4.78 (<i>d</i> ; 11.2) - 5.50 (<i>ov.</i>)
25	0.94 (<i>s</i>)	0.94 (<i>s</i>)	0.84 (<i>s</i>)	1.16 (<i>s</i>)
26	1.05 (<i>s</i>)	0.97 (<i>s</i>)	1.15 (<i>s</i>)	1.00 (<i>s</i>)
27	1.68 (<i>s</i>)	1.70 (<i>s</i>)	1.71 (<i>s</i>)	1.63 (<i>s</i>)
29	1.42 (<i>s</i>)	1.36 (<i>s</i>)	1.37 (<i>s</i>)	1.37 (<i>s</i>)
30	1.10 (<i>d</i> ; 7.0)	0.98 (<i>d</i> ; 6.6)	0.97 (<i>d</i> ; 7.1)	0.96 (<i>d</i> ; 6.8)
Ara-1 $^{\circ}$		4.83 (<i>d</i> ; 7)	4.87 (<i>d</i> ; 6.7)	
Ara-2 $^{\circ}$		5.77 (<i>ov.</i>)	4.42 (<i>ov.</i>)	
Ara-3 $^{\circ}$		5.56 (<i>dd</i> ; 10.0, 3.3)	4.19 (<i>dd</i> ; 8.2, 3.4)	
Ara-4 $^{\circ}$		5.64 (<i>ov.</i>)	4.35 (<i>ov.</i>)	
Ara-5 $^{\circ}$		3.90 (<i>d</i> ; 13.0) - 4.28 (<i>dd</i> ; 13.0, 2.3)	3.85 (<i>d</i> ; 10.9) - 4.36 (<i>ov.</i>)	
Glc-1 $^{\circ}$		6.35 (<i>d</i> ; 8.0)	6.34 (<i>d</i> ; 7.8)	6.36 (<i>d</i> ; 8.0)
Glc-2 $^{\circ}$		5.75 (<i>ov.</i>)	4.23 (<i>t</i> ; 8.5)	5.74 (<i>dd</i> ; 8.6, 8.4)
Glc-3 $^{\circ}$		5.98 (<i>t</i> ; 9.5)	4.31 (<i>t</i> ; 8.5)	5.97 (<i>dd</i> ; 9.5, 8.6)
Glc-4 $^{\circ}$		5.63 (<i>ov.</i>)	4.40 (<i>ov.</i>)	5.62 (<i>dd</i> ; 9.6, 9.5)
Glc-5 $^{\circ}$		4.41 (<i>ddd</i> ; 9.5, 4.2, 2.2)	4.05 (<i>ddd</i> ; 9.7, 3.8, 3.0)	4.41 (<i>ddd</i> ; 9.2, 4.1, 2.0)
Glc-6 $^{\circ}$		4.33 (<i>dd</i> ; 12.1, 2.2) 4.63 (<i>dd</i> ; 12.3, 4.2)	4.38 (<i>ov.</i>) 4.43 (<i>ov.</i>)	4.32 (<i>d</i> ; 12.1) 4.60 (<i>dd</i> ; 12.6, 4.3)

*Values for the acetyl groups are not recorded in the table.

rial 18β -H) in the HH COSY experiment. Further, evidence for an equatorial 18β -H was provided by heteronuclear long-range correlation *via* three bonds couplings, ${}^3J({}^{13}\text{C}, {}^1\text{H})$, detected by the HMBC experiment. This proton displayed cross-signals with carbons C-14 (δ_{C} 42.1) and C-16 (δ_{C} 27.0), revealing a dihedral angle close to 180° , providing evidence for the *cis*-fusion between the rings D/E.

A triplet of triplets (${}^3J=13.3$ and 3.3 Hz) observed for 21α -H (δ_{H} 2.68, axial) revealed the β -configuration of the C-30 methyl group (δ_{H} 16.1), corresponding to an ursane derivative with the $20(S)$ -configuration that could be confirmed by means of the ROESY experiment, indicating spatial correlation between the 18β -H (δ_{H} 3.26) and hydrogens 29β -CH₃ (δ_{H} 1.42) and 30β -CH₃ (δ_{H} 1.10). Compared with ursolic acid (Tkachev *et al.*, 1994), compound **1a** presented two important γ -effects in the ring E. The first one was caused by the axial 30β -CH₃,

shielding the carbons C-18 and C-22 by 5 and 4 ppm, respectively, in the ${}^{13}\text{C}$ NMR. The second effect was observed due to the presence of the axial 19α -hydroxy group, that shielded C-21 by about 6 ppm. Thus, **1a** differs from the structure of rotundioic acid (Nakatni *et al.*, 1989) by a different configuration of C-20 and turns out to be the peracetylated derivative of $20(S)$ - $3\beta, 19\alpha$ -dihydroxyurs- $12\text{-en-}23,28$ -dioic acid, named brevicuspisic acid.

Fast atom bombardment mass spectroscopy (FAB-MS, positive mode) of **2a** generated a fragment at m/z 1083 $[\text{M}+\text{Na}]^+$ as a pseudo molecular ion, in accordance with seven acetylated OH groups and with the spectrum obtained for the non-peracetylated compound **2**, showing a pseudo molecular ion at m/z = 789 $[\text{M}+\text{Na}]^+$. ${}^{13}\text{C}$ NMR of **2a** data revealed the occurrence of signals from one carboxy carbon (δ_{C} 176.5), seven acetate groups (δ_{C} 169 - 170 and δ_{C} 20.7 - 21.1), one double bond (δ_{C} 128.1 and




FIGURE 1 - Ring D from **1a**: additional evidence for the relative configuration by the ${}^1\text{H}$ NMR coupling constants (a), *W*-type correlation deduced from the HH COSY diagram (b) and long-range correlation ${}^3J({}^{13}\text{C}, {}^1\text{H})$ in the HMBC experiments (c).




FIGURE 2 - Main ROESY correlation observed for compound **4a** in pyridine- d_5 .

138.4), two anomeric carbons (δ_C 104.0 and 92.5), one glycosylated *sec* alcohol (δ_C 89.5) and one *tert* alcohol (δ_C 73.4). ^1H NMR confirmed the presence of six angular methyl groups (δ_H 2x 0.94, 0.97, 1.08, 1.36 and 1.70), one methyl group attached to CH (δ_H 0.98, d , $^3J = 6.6$ Hz), one olefinic proton (δ_H 5.50) and two anomeric sugar protons (δ_H 4.83, d , $^3J = 8.0$ Hz and δ_H 6.35, d , $^3J = 8.0$ Hz). Both sugars presented antiperiplanar configuration of 1'-H and 2'-H ($^3J = 8.0$ Hz), reflecting the α -configuration of the L-arabinopyranose and the β -configuration of the D-glucopyranose. The glycosidic linkages were established using the HMBC techniques. It was possible to observe correlation between the anomeric proton of α -L-arabinose (δ_H 4.83) and C-3 (δ_C 89.5), and also between the anomeric proton of β -D-glucose (δ_H 6.35) and C-28 (δ_C 176.5).

ROESY experiments were used to establish the α -configuration for 1'-H (δ_H 4.83), 3-H (δ_H 3.23) and 23-CH₃ (δ_H 1.08, *s*). Moreover, the β -configuration of 29-CH₃ (δ_H 1.36, *s*) attached to C-19 (δ_C 73.4) indicated the presence of a 19 α -*tert* alcohol group. In the same way, the α -configuration of 20-H (δ_H 1.92) defined the configuration 30 β -CH₃ (δ_H 0.98, d , $^3J = 6.6$ Hz) at C-20. These data indicated the aglycone to be the 20(*S*)-isomer of pomolic acid (Brieskorn *et al.*, 1967) and compound **2a** was elucidated as the peracetylated derivative obtained from the 3-*O*- α -L-arabinopyranosyl-20(*S*)-pomolic acid-28-*O*- β -D-glucopyranosyl ester, named brevicuspisaponin 3.

Examination of compound **3** allowed its characterization as ILA-1, a saponin already isolated from the leaves of *Ilex argentina* (Schenkel *et al.*, 1995). One- and two-dimensional NMR spectroscopic data confirmed **3** as 3-*O*- α -L-arabinopyranosyl-20(*S*)-19 α ,24-dihydroxyursolic acid-28-*O*- β -D-glucopyranosyl ester.

Compound **4** was characterized as a sodium salt by IR that shows a symmetric stretching band at 1573 cm⁻¹, a diagnostic frequency for a carboxylic acid salt. The presence of the sodium was determined by the flame ionization, employing a polarized filter for the D-sodium line, and compounds **3** as negative control. Acidic hydrolysis on TLC plates (Kartning, Wegschaidler, 1972) indicated the presence of glucose. The FABMS of this genuine saponin (positive-ion mode) displayed peaks at $m/z = 725.3$ [M+Na]⁺ and 703.3 [M+H]⁺. Compound **4a** showed the pseudo-molecular ion by FABMS (positive-ion mode) at $m/z = 955.4$ [M+H]⁺.

^{13}C NMR spectra of compound **4a** revealed the presence of signals of two carboxy carbons (δ_C 176.4 and 176.5), one double bond (δ_C 127.7 and 138.4), one anomeric sugar carbon (δ_C 92.4), one *sec*-hydroxyl (δ_C 77.9), one *tert*-hydroxyl functions (δ_C 73.3) and two

hydroxymethyl groups (δ_C 62.3 and 63.9). The ^1H NMR spectrum showed the presence of four angular methyl groups (δ_H 1.00, 1.16, 1.37 and 1.63), one methyl group attached to CH (δ_H 0.96, d , $^3J_{HH} = 6.8$ Hz), one olefinic proton (δ_H 5.49) and one anomeric sugar proton (δ_H 6.36, d , $^3J_{HH} = 8.0$ Hz). Its NMR data comparison with compound **2a** (table 2) indicated that the sugar β -D-glucose was esterified with the C-28 carboxy function in the aglycone.

The location of the carboxylate group at C-4 was derived from the HMBC correlation signals δ_C 176.5 \leftrightarrow δ_H 4.78 (24_A-H) and 5.50 (24_B-H). Moreover, the carbon resonance of this hydroxymethyl group presented long-range correlation with the signals at δ_H 5.77 (3 α -H) and 2.18 (5 α -H). The occurrence of a triplet of triplets at δ_H 2.55 ($J_{HH} = 13.2$ and 3.5 Hz) indicated an axial hydrogen 21 α -H and thus, the (20*S*)-configuration. It was corroborated through spatial correlation observed between 18 β -H (δ_H 3.03) and hydrogens 29 β -CH₃ (δ_H 1.37) and 30 β -CH₃ (δ_H 0.96; d , $^3J_{HH} = 6.8$ Hz) in the ROESY diagram (Figure 2). Furthermore, ROESY correlation between signals at δ_H 4.78 (24_A β -H) and δ_H 1.16 (25 β -CH₃) also established the configuration on C-4 and located the sodium carboxylate group at C-23 (4 α -configuration). This experiment also showed the correlation between 3 α -H (δ_H 5.77) and 5 α -H (δ_H 2.18). Thus, compound **4a** is a new saponin and was elucidated as the peracetylated derivative from of the 23-sodium salt of (20*S*)-3 β ,19 α ,24-trihydroxyurs-12-en-23,28-dioic acid-28 β -*O*- β -D-glucopyranosyl ester, named brevicuspisaponin 4.

Considering these and the previous published results (Taketa *et al.*, 2000), the saponin profile of *I. brevicuspis* leaves is markedly different from those found in *I. paraguariensis* leaves. The latter presents saponins derived from the oleanolic and ursolic acid without oxygenated functions at C-19, C-23 or C-24, as demonstrated to *I. brevicuspis*. These are important features that may be useful to develop methodologies for the quality control of maté products based on the characterization of the free triterpenes or the saponins.

ACKNOWLEDGMENTS

We are grateful to the Botanist Geraldo C. Coelho from the Departamento de Biologia e Química of Universidade de Ijuí, Ijuí/RS for locating, identifying and helping to collect the plant material, to Dr. G. Eckhardt and to Mrs. U. Dahmen (Bonn University), for recording the mass and IR spectra. This work was supported by research stipends from DAAD (Germany), and CNPq (Brazil).

RESUMO

Novos triterpenos e glicosídeos triterpenóides de *Ilex brevicuspis*

Das folhas de *Ilex brevicuspis* foram isolados e identificados um novo triterpeno, ácido 20(S)-3 β ,19 α -diidroxiurs-12-en-23,28-dióico, denominado ácido brevicúspico, e dois novos glicosídeos, éster 28-O- β -D-glicopiranosil do ácido 3 β -O- α -L-arabinopiranosil-20(S)-pomólico e o sal sódico em C-23 do éster 28 β -O- β -D-glicopiranosil do ácido (20S)-3 β ,19 α ,24-triidroxiurs-12-en-23,28-dióico. Foi isolado ainda o éster 28-O- β -D-glicopiranosil do ácido 3 β -O- α -L-arabinopiranosil-20(S)-19 α ,24-diidroxiursólico, já descrito anteriormente para a espécie *Ilex argentina*. As estruturas foram estabelecidas por métodos espectroscópicos e químicos.

UNITERMOS: Aquifoliaceae. *Ilex brevicuspis*. Triterpenos. Saponinas. Ácido brevicúspico. *Brevicuspisaponinas 3 e 4*.

REFERENCES

- ATHAYDE, M. L., SCHENKEL, E. P., GOSMANN, G., GUILLAUME, D. Triterpenoids from the leaves of *Ilex theezans* Martius ex Reiss. *Acta Farm. Bonaerense*, v. 18, p. 49-52, 1999.
- ATHAYDE, M. L., SCHENKEL, E.P., GNOATO, S., GOSMANN, G., GIBERTI, G.C., GUILLAUME D. Triterpenes and saponins from *Ilex argentina* leaves. *Acta Farm. Bonaerense*, v. 20, p.13-16, 2001.
- BRIESKORN, C. H., WUNDERER, H. Pomol- und Pomolsäure. *Chem. Ber.*, v. 100, p. 1252-1253, 1967.
- HEINZMANN, B.M., SCHENKEL, E. P. Saponins from *Ilex dumosa*. *J. Nat. Prod.*, v. 58, p. 1419-1422, 1995.
- KARTING, T., WEGSCHAIDER, O. Zur Kenntnis der Saponine aus *Herniaria glabra*. *Planta Med.*, v. 21, p.144-150, 1972.
- NAKATANI, M., MIYAZAKI, Y., IWASHITA, T., NAOKI, H., HASE, T. Triterpenes from *Ilex rotunda* fruits. *Phytochemistry*, v. 28, p.1479-1482, 1989.
- PIRES, V. S., GUILLAUME, D., GOSMANN, G., SCHENKEL, E. P. "Erva-Mate" adulterating plants: saponins from *Ilex dumosa*. *J. Agric. Food Chem.*, v. 45, p.1027-1031, 1997.
- REGINATTO, F. H., ATHAYDE, M. L., GOSMANN, G., SCHENKEL, E. P. Methylxanthines accumulation in *Ilex* species - Caffeine and theobromine in erva-mate (*Ilex paraguariensis*) and other *Ilex* species. *J. Braz. Chem. Soc.* v. 10, p.443-446, 1999.
- SCHENKEL, E. P., ATHAYDE, M. L., GIBERTI, G. C., GUILLAUME, D. A new saponin from *Ilex argentina*. *Acta Farm. Bonaerense*, v.14, p.5-10, 1995.
- SCHENKEL, E.P., GOSMANN, G., MONTANHA, J.A., HEINZMANN, B.M., ATHAYDE, M.L., TAKETA, A.T.C., PIRES, V., GUILLAUME, D. Saponins from maté *Ilex paraguariensis* and other South American *Ilex* species: Ten years research on *Ilex* saponins. *Ciênc. Cult.*, v. 49, p.315-320, 1997.
- TAKETA, A. T. C., SCHENKEL, E. P. Saponins from *Ilex pseudobuxus*. *Acta Farm. Bonaerense*, v.13, p.159-164, 1994.
- TAKETA, A. T. C., SCHMITTMANN-SCHLAGER, T., GUILLAUME, D., GOSMANN, G., SCHENKEL, E.P. Triterpenoid glycosides and a triterpene from *Ilex brevicuspis*. *Phytochemistry*, v. 53, p. 901-904, 2000.
- TKACHEV, A. V., DENISOV, A. Y., GATILOV, Y. V., BAGRYANSKAYA, I. Y., SHEVTSOV, S. A., RYBALOVA, T. V. Stereochemistry of hydrogen peroxide-acetic acid oxidation of ursolic acid and related compounds. *Tetrahedron*, v. 50, p.11459-11462, 1994.

Recebido para publicação em 20/12/01.