

Mowat-Wilson Syndrome

Síndrome de Mowat-Wilson

Carlos Eduardo Steiner

Departamento de Genética Médica,
Faculdade de Ciências Médicas,
Universidade Estadual de
Campinas, Campinas SP, Brazil.

Correspondence:

Carlos Eduardo Steiner; Rua
Tessália Vieira de Camargo, 126;
13083-887 Campinas SP; Brasil;
E-mail: steiner@fcm.unicamp.br

Conflict of interest:

There is no conflict of interest to
declare.

Received 11 November 2014

Accepted 01 December 2014

As medical specialties, Neurology, Psychiatry, and Clinical Genetics share many affinities, not only because 80% of the human genome is expressed in the central nervous system, but also due to parallels in their history. Congenital abnormalities and traumatic injuries causing neurobehavioral deficits allowed drawing a map of the brain functional structure, beginning in the modern medicine with the notorious report¹ of Phineas Gage casualty in 1848 and the first publication of the cortical cartography represented as a *homunculus* by Penfield and Boldrey² in 1937.

In a similar way, translocations and unbalanced abnormalities seen in karyotype of individuals with single gene disorders also allowed locus identification for type 1 neurofibromatosis (17q11.2), Duchenne muscular dystrophy (Xp21), Cornelia de Lange syndrome (5p13.2), and many others. A first map of the human genome was published³ in 2001 and refined in the following years.

Besides, several chromosomal microdeletion and microduplication present with characteristic neurobehavioral features, such as friendly loquacious personality in Williams syndrome, obsession with eating, food related behavior, and unusual skill with jigsaw puzzle in Prader-Willi syndrome, or polyembolokoilamania, onychotillomania, sleep disturbance, and self-destructive behavior in Smith-Magenis syndrome. The 22q11 deletions in velocardiofacial syndrome are frequently associated with schizophrenia⁴. Patients presenting with these conditions, also denominated contiguous gene disorders, are frequently seen by neurologists, psychiatrists, and clinical geneticists who independently or in interdisciplinary teams helped delineating the neurobehavioral phenotype and consensus diagnostic criteria.

Mowat-Wilson syndrome (MWS) is another example of a condition reflecting these medical specialties overlap. It was initially reported on six children, one of them presenting with a chromosome 2q deletion⁵. In fact, MWS was later associated to heterozygous mutations in the *ZEB2* gene located in 2q22.3 which cause abnormalities in the neural crest development resulting in deviant facial features, functional and structural defects of the central nervous system (development delay, seizures, microcephaly, hypoplasia of the corpus callosum), heart defects, and intestinal aganglionosis leading to Hirschsprung disease⁶.

As in most developmental delay syndromes, MWS has been associated to a range of non specific behavioral symptoms like hypotonia, marked delay in the motor milestones, learning problems, moderate to severe intellectual deficiency, epilepsy, attention deficit disorder, and autistic features, but a characteristic neuropsychological profile has also been proposed⁶. Performance in MWS included sociable demeanor and happy affect with frequent laughs, a high rate of oral behaviors (bruxism, chewing or mouthing objects or body parts), an increased rate of repetitive behaviors, under-reaction to pain, and severely impaired or absent speech with contrasting better receptive language⁷.

The 15 years following its original description revealed remarkable discoveries regarding MWS physiopathology, molecular basis, inheritance, and phenotypic spectrum, but diagnostic criteria and a precise incidence have not been defined yet. Half of the literature on this condition was published in the last five years and the rate of publications on it increases every year. Thus, the diagnosis of MWS must be considered by professionals that evaluate patients with developmental disorders.

In this issue, Paz et al.⁸ present the first series of Brazilian patients diagnosed with MWS, adding new data to the international literature. Seven individuals were diagnosed by mutation analysis of the *ZEB2* gene. Detailed information on

dysmorphic and neurological examination, epileptic history, electroencephalographic findings, Denver II scale evaluation, and brain magnetic resonance imaging are discussed.

References

1. Van Horn JD, Irimia A, Torgerson CM, et al. Mapping connectivity damage in the case of Phineas Gage. *PLoS ONE*. 2012;7(5):e37454.
2. Schott GD. Penfield's homunculus: a note on cerebral cartography. *J Neurol Neurosurg Psychiatry*. 1993;56:329-33.
3. The International Human Genome Mapping Consortium. A physical map of the human genome. *Nature*. 2001;409:934-41.
4. Watson CT, Marques-Bonet T, Scharp AJ, Mefford HC. The genetics of microdeletion and microduplication syndromes: an update. *Annu Rev Genomics Hum Genet*. 2014;31:215-44. <http://dx.doi.org/10.1146/annurev-genom-091212-153408>
5. Mowat DR, Croaker GDH, Cass DT, Kerr BA, Chaitow J, Adès LC et al. Hirschsprung disease, microcephaly, mental retardation, and characteristic facial features: delineation of a new syndrome and identification of a locus at chromosome 2q22-q23. *J Med Genet*. 1998;35(8):617-23. <http://dx.doi.org/10.1136/jmg.35.8.617>
6. Van de Putte T, Francis A, Nelles L, Grunsven LA, Huylebroeck D. Neural crest-specific removal of *Zfx1b* in mouse leads to a wide range of neurocristopathies reminiscent of Mowat-Wilson syndrome. *Hum Mol Genet*. 2007;16(12):1423-36. <http://dx.doi.org/10.1093/hmg/ddm093>
7. Evans E, Einfeld S, Mowat D, et al. The behavioral phenotype of Mowat-Wilson syndrome. *Am J Med Genet Part A*. 2012;158A:358-66. <http://dx.doi.org/10.1002/ajmg.a.34405>
8. Paz JA, Kim CA, Goossens M, Giurgea I, Marques-Dias MJ. Mowat-Wilson syndrome: neurological and molecular study in seven patients. *Arq Neuropsiquiatr* 2015;73(1):12-17.