

Spinal cord compression due to extramedullary hematopoiesis in beta-thalassemia

Compressão da medula espinhal por hematopoese extramedular na beta-talassemia

Renan Ramon Souza LOPES¹, Larissa Soares CARDOSO², Franz ONISHI¹

This study reports the case of a 35-year-old man with beta-thalassemia intermedia, with irregular treatment, who presented progressive paresthesia mainly affecting the left side and strength loss in the lower limbs for five months. Besides, he developed difficulty to ambulate and urinate. MRI (Figures 1, 2, 3 and 4) showed vertebral canal stenosis with spinal cord compression. This

finding illustrates a rare complication of thalassemia, an extramedullary hematopoietic center due to ineffective erythropoiesis, causing vertebral bone hyperplasia^{1,4,5}. Given the substantial number of segments affected, surgical procedures were contraindicated based upon hemorrhage risk. Radiotherapy was performed with subsequent strength recovery^{2,3}.




Figure 1. T1-weighted axial MRI with contrast showing moderate contrast enhancement of the epidural masses. Ribs and the vertebral bodies are also enhanced.




Figure 2. T2-weighted axial MRI showing severe spinal cord compression (the black arrows indicate the ribs and the white one indicates the vertebral bone).

¹Universidade Federal de São Paulo, Escola Paulista de Medicina, São Paulo SP, Brazil.

²Universidade Federal de Roraima, Boa Vista RR, Brazil.

Renan Ramon Souza LOPES  <https://orcid.org/0000-0002-5512-4612>; Larissa Soares CARDOSO  <https://orcid.org/0000-0002-3495-4473>; Franz ONISHI  <https://orcid.org/0000-0002-5641-5104>

Correspondence: Franz Jooji Onishi; E-mail: franzoneishi@gmail.com

Conflict of interest: There is no conflict of interest to declare.

Authors' contributions: RRSL: work design, writing and preparation, and final revision of the text content and suitability to the journal format; LSC: work design, writing and preparation, and final revision of the text content and suitability to the journal format; FO: work design, writing and preparation, and final revision of the text content and suitability to the journal format.

Received on March 23, 2020; Received in its final form on May 05, 2020; Accepted on May 08, 2020.



Figure 3. T1-weighted sagittal MRI showing moderate and heterogeneous enhancement of epidural masses.



Figure 4. T2-weighted sagittal MRI showing spinal cord compression caused by an extensive epidural lesion.

References

1. Duque SG, Jureschke FR, Leal RG. Reversible paraparesis secondary to spinal cord compression in a patient with β -thalassaemia. *Neurologia*. 2019 May;34(4): 270-2.
2. Fareed S, Soliman AT, De Sanctis V, Kohla S, Soliman D, Khirfan D, et al. Spinal cord compression secondary to extramedullary hematopoiesis: A rareness in a young adult with thalassemia major. *Acta Biomed*. 2017 Aug;88(2):237-42. <https://doi.org/10.23750/abm.v88i2.6221>
3. Moncef B, Hafedh J. Management of spinal cord compression caused by extramedullary hematopoiesis in beta-thalassemia. *Intern Med*. 2008;47(12):1125-8. <https://doi.org/10.2169/internalmedicine.47.0890>
4. Ruo Redda MG, Allis S, Reali A, Bartoncini S, Roggero S, Anglesio SM, et al. Complete recovery from paraparesis in spinal cord compression due to extramedullary haemopoiesis in beta-thalassaemia by emergency radiation therapy. *Intern Med J*. 2014 Apr;44(4):409-12. <https://doi.org/10.1111/imj.12386>
5. Yathiraj PH, Singh A, Vidyasagar S, Varma M, Mamidipudi V. Excellent and durable response to radiotherapy in a rare case of spinal cord compression due to extra-medullary hematopoiesis in β -thalassemia intermedia: case report and clinicoradiological correlation. *Ann Palliat Med*. 2017 Apr;6(2):195-9. <https://doi.org/10.21037/apm.2016.12.05>