

Methicillin-resistant *Staphylococcus* spp. isolated from curd cheese “requeijão” and “especialidade láctea type requeijão” sold in Brazil

Jamile de Oliveira Hachiya¹ Gabriel Augusto Marques Rossi^{1*} Laryssa Freitas Ribeiro¹
Rafael Akira Sato¹ Higor Oliveira Silva¹ Ana Maria Centola Vidal² Luiz Augusto do Amaral¹

¹Departamento de Medicina Veterinária Preventiva e Reprodução Animal, Faculdade de Ciências Agrárias e Veterinárias (FCAV), Universidade Estadual Paulista (UNESP), Via de Acesso Paulo Castellane, s/n, 14884-900, Jaboticabal, SP, Brasil. E-mail: gabrielrossiveterinario@gmail.com. *Corresponding author.

²Departamento de Medicina Veterinária, Universidade de São Paulo (UNESP), Faculdade de Zootecnia e Engenharia de Alimentos (FZEA), Pirassununga, SP, Brasil.

ABSTRACT: This study focused on counting *Staphylococcus* spp. in curd cheeses “requeijão” and “especialidade láctea type requeijão” sold in Brazil, assessing the presence of *mecA* gene in obtained isolates and establishing antimicrobial resistance profile of the *mecA* gene positive isolates. To this, a set of 200 samples of these dairy products were evaluated. Low counts of *Staphylococcus* spp. were observed in these dairy products. All the isolates were determined as coagulase-negative strains using coagulase test and PCR. However, two isolates (3.70%) were carriers of *mecA* gene and they can be considered as risk for public health. These isolates presented resistance to penicillin, oxacillin and erythromycin. In conclusion, low counts of *Staphylococcus* were detected in curd cheese “requeijão” and “especialidade láctea type requeijão” sold in Brazil. However, coagulase-negative methicillin-resistant *Staphylococcus* spp. was detected in these dairy products. This is the first report of the detection of methicillin-resistant *Staphylococcus* spp. in heat-treated dairy products in Brazil. Results served as a warning to public sanitary authorities to control multidrug-resistant strains in veterinary and human medicine.

Key words: antimicrobial resistance, cheese, food safety, MRS, *Staphylococcus* spp.

Isolamento de *Staphylococcus* spp. resistente à meticilina de queijos fundidos requeijão e especialidade láctea tipo requeijão comercializados no Brasil

RESUMO: Este trabalho objetivou realizar a contagem de *Staphylococcus* spp. em queijos fundidos “requeijão” e “especialidade láctea tipo requeijão” comercializados no Brasil, verificar a presença do gene *mecA* nos isolados obtidos e estabelecer o perfil de resistência antimicrobiana dos isolados positivos para tal gene. Para isso, 200 amostras desses produtos lácteos foram avaliadas. Baixas contagens de *Staphylococcus* spp. foram observadas nas amostras. Todos os isolados foram considerados como coagulase-negativos através do teste da coagulase e através da PCR. Entretanto, em dois isolados (3,70%) foi possível detectar o gene *mecA* e representam potencial risco à saúde pública. Esses isolados apresentaram resistência a penicilina, oxacilina e eritromicina. Conclui-se que baixas contagens de *Staphylococcus* foram detectadas em queijos fundidos “requeijão” e “especialidade láctea type requeijão”. Entretanto, cepas de *Staphylococcus* spp. coagulase-negativas e resistentes à meticilina foram detectadas nesses derivados lácteos. Esse é o primeiro relato da ocorrência de cepas de *Staphylococcus* spp. resistentes à meticilina em produtos lácteos termicamente tratados comercializados no Brasil. Os resultados servem como um alerta para as autoridades sanitárias públicas nacionais para o controle de cepas multirresistentes em medicina veterinária e humana.

Palavras-chave: resistência antimicrobiana, segurança alimentar, MRS, queijos, *Staphylococcus* spp.

INTRODUCTION

“Requeijão” is the third most sold cheese in Brazil (ZACARCHENCO et al., 2012). “Requeijão” is defined as “the product obtained using acid or enzymatic milk coagulation followed by the curd mass fusion (80°C for 5 seconds), whey removal and washing according to Brazilian legislation (BRAZIL, 1997). Also, addition of milk cream, butter or butter oil in its composition is allowed. Another curd cheese routinely consumed in Brazil is “especialidade láctea type requeijão”, a similar cheese produced using the same process but the

addition of vegetable ingredients such as corn starch and vegetable oil is allowed in composition.

The genus *Staphylococcus* is among the most important pathogenic bacteria reported in dairy products. This genus is divided into two groups: coagulase-positive and coagulase-negative strains. Among the first one, *S. aureus* is known for its clinical importance and as a foodborne pathogen. The coagulase-negative group is comprised by several species involved in opportunistic infections in immunocompromised patients (MARTINS & CUNHA, 2007).

Staphylococcus spp. are also frequently associated to mastitis (PITKALA et al., 2004), biofilm formation (OTTO, 2009) and are closely related to improper food handling of food handlers (ARGUDÍN et al., 2012), leading contamination of milk and other dairy products, posing public health into risk. Consequently, *Staphylococcus* spp. antimicrobial resistance is a concern for public health (KOKSAL et al., 2009).

Methicillin resistance is an increasing global problem (MARTINS & CUNHA, 2007), even for domestic animals (CUNY et al., 2010). The presence of these isolates in food and animals highlight the potential risk for human transmission (WEESE & DUJIKEREN, 2010). Some researchers have reported strains of methicillin resistant *Staphylococcus* spp. (MRS) causing mastitis (MOON et al., 2007), in bulk tank milk (VIRGIN et al., 2009), in bovine milk, pecorino and mozzarella cheeses (NORMANO et al., 2009) and in raw meats (BOER et al., 2009). Based on all these information, this study focused on counting *Staphylococcus* spp. in curd cheeses “requeijão” and “especialidade láctea type requeijão” sold in Brazil, assessing the presence of *mecA* gene in obtained isolates and establishing antimicrobial resistance profile of the *mecA* gene positive isolates.

MATERIALS AND METHODS

This research was performed using samples of “requeijão”, “requeijão light”, “especialidade láctea type requeijão” and “especialidade láctea type requeijão light”, obtained from supermarkets located in the municipality of Jaboticabal, state of São Paulo, Brazil, during the year of 2015. Samples were kept refrigerated in cooler boxes with ice, during the transport to laboratory. Each kind of dairy product (“requeijão”, “requeijão light”, “especialidade láctea type requeijão”) had 50 samples bought, from 5 different commercial brands, each one with 2 different production lots (each lot composed by 5 samples). For the “especialidade láctea type requeijão light”, the samples were from three different commercial brands and 10 different production lots (each lot composed by 5 samples). In total, 200 samples were used for this study.

For each sample, 25g was placed into a Scott Bottle containing 225mL of 0.1% peptone water (Difco, Detroit, USA). Out of this first dilution (10^{-1}), other serial dilutions were performed, until 10^{-3} . Selective plating was done using 0.1mL of these dilutions in Baird-Parker agar (Difco, Detroit,

USA). Plates were incubated for 45-48 hours under 35°C. Whenever black colonies grew in the agar, it was transferred to a tube with Tryptic Soy Agar (TSA) (Difco, Detroit, USA) medium, and incubated for 24 hours under 35°C (APHA, 2011). Moreover, the colonies were transferred to tubes containing Brain Heart Infusion broth (BHI) (Difco, Detroit, USA) and incubated for 24 hours under 37°C. The coagulase test was performed using the isolates from BHI broth and rabbit plasma, in order to assess the presence of coagulase-positive strains (APHA, 2011).

The DNA from the 57 isolates was extracted according to the method described by KESKIMAKI et al. (2001). Presence of *Staphylococcus* spp. and *S. aureus* DNA molecules was detected using primers 16SrRNA (Pereira et al., 2009) and Sa442-1/2 (MOROT-BIZOT et al., 2004). A PCR also was performed to identify coagulase-positive *Staphylococcus* spp. (*S. intermedius*, *S. schleiferi* subsp. *coagulans*, *S. hyicus*, *S. lutrae*, *S. delphini*, and *S. pseudintermedius* according to SASAKI et al. (2010) (Table 1). In order to detect methicillin-resistant isolates (*mecA*), a PCR was performed using the primers as described by GORTEL et al. (1999): 5' AAA ATC GAT GGT AAA GGT TGG C 3' e 5' AGT TCT GCA GTA CCG GAT TTG C 3', with a 533bp product and 55°C annealing temperature.

PCR amplifications were carried out with mixtures containing PCR buffer 1x [100mM Tris-HCl pH 8.8; 500mMKCl; 0.8% (v/v) Nonidet P40]; MgCl₂ 2mM; dNTP's 0.2mM, 1.5U de Taq DNA polymerase, 5pmol of each primer, 60ng genomic DNA and sterile water until reaching the volume of 20μL. Cycling conditions consisted of an initial step at 95°C for 5min, followed by 35 cycles at 94°C for 1min, annealing at the specific temperature of each primer for 1min and 72°C for 1.5min. The final extension was at 72°C for 10min. The amplified products were visualized in a 1% agarose gel electrophoresis stained with 0.5μg mL⁻¹ ethidium bromide (Vetec, São Paulo, Brazil). A previously known molecular weight marker was used as a reference (GeneRuler 1kb DNA Ladder – Fermentas, Thermo Fish Scientific, São Paulo, Brazil).

MRS positive isolates were examined for resistance to 11 antimicrobials by disk-diffusion (Kirby-Bauer) method. The following disks (Laborclin™) were used: oxacillin (1μg), cefepime (30μg), rifampicin (5μg), chloramphenicol (30μg), clindamycin (2μg), erythromycin (15μg), penicillin G (10un), sulfatrim (23,75/1,25μg), tetracycline (30μg), gentamicin (10μg) and ciprofloxacin (5μg).

Table 1 - Primer, nucleotides sequence, amplicon size (bp), annealing temperature, positive control and reference of each primer used for the detection of *Staphylococcus* spp.

	Primer	Nucleotides Sequence	Size (bp)	Annealing temperature (°C)	Positive control	References
<i>Staphylococcus</i> spp.	16srRNA	GTA GGT GGC AAG CGTTAT CC CGC ACA TCA GCG TCA G	228	55	ATCC 25983	Pereira et al., 2009
<i>S. aureus</i>	Sa442	AATCTTTGTCGGTACACGATATT CTTCACG CGTAATGAGATTTTCAGTAGATAA TACAACA	102	55	ATCC 25983	Morot-Bizot et al., 2004
<i>S. aureus</i>	au-F3 au-nucR	TCGCTTGCTATGATTGTGG GCCAATGTTCTACCATAGC	359		ATCC 25983	
<i>S. intermedius</i>	in-F in-R3	CATGTCATATTATTG CGAATGA AGGACCATCACCATT GACATATTGA AACC	430			
<i>S. schleiferi</i> subspecie coagulans	sch-F sch-R	AATGGCTACAATGAT AATCACTAA CATATCTGTCTTTTCG GCGCG	526			
<i>S. hyicus</i>	hy-F1 hy-R1	CATTATATGATTTGA ACGTG GAATCAATATCGTAA AGTTGC	793			Sasaki et al., 2010
<i>S. deplhini</i> grupo A	dea-F dea-R	TGAAGGCATATTGTA GAACAA CGRTACTTTTCGTGA GGTCG	661			
<i>S. deplhini</i> grupo B	debF debR4	GGAAGRITTCGTTTTTCCTAGAC TATGCGATTCAAGAA CTGA	1135			
<i>S. pseudintermedius</i>	pseF2 pseR5	TRGGCAGTAGGATT CGTTAA CTTTTGTGCTYCMIT TTGG	926	56		

The significance between the *Staphylococcus* spp. counting of the groups was assessed through a variance analysis and the Kruskal-Wallis test with 5% of significance level. Data regarding the “presence or absence” of *Staphylococcus* spp. in the samples group (comparison among light and normal products; and among “requeijão” and “especialidade láctea” products) was performed using the chi-square test with 5% of significance level. The analyses were performed using Software R®.

RESULTS AND DISCUSSION

Table 2 presents *Staphylococcus* spp. counts of the dairy product samples analyzed in

this study. Results of *Staphylococcus* spp. count did not meet the necessary requirements to perform the variance analysis. Moreover, when using the nonparametric Kruskal-Wallis test with 5% of significance level, no statistical significance ($P > 0.05$) was observed among these products. The chi-square test performed to evaluate the difference among the presence or absence of these microorganisms in the products also presented no significance ($P > 0.05$).

Staphylococcus spp. count in the samples was low and all the strains were considered as coagulase-negative strains on coagulase-test. *S. aureus* is considered as the most important foodborne pathogen of this group and it is considered as coagulase-positive specie. Based on such results,

Table 2 - Average *Staphylococcus* spp. count in “requeijão”, “requeijão light”, “especialidade láctea type requeijão” and “especialidade láctea type “requeijão light” sold in Jaboticabal municipality, state of São Paulo, Brazil during 2015.

Product	Brand	Lot	<i>Staphylococcus</i> spp. count (CFU g ⁻¹)	Product	Brand	Lot	<i>Staphylococcus</i> spp. count (CFU g ⁻¹)		
“Requeijão”	A	1	<0.1 x 10 ⁰	“Especialidade láctea”	A	1	8.6 x 10 ⁰		
		2	<0.1 x 10 ⁰			2	<0.1 x 10 ⁰		
	B	1	<0.1 x 10 ⁰		B	1	<0.1 x 10 ⁰		
		2	0.3 x 10 ⁰			2	0.1 x 10 ⁰		
	C	1	<0.5 x 10 ⁰		C	1	<0.1 x 10 ⁰		
		2	<0.5 x 10 ⁰			2	<0.1 x 10 ⁰		
	D	1	0.8 x 10 ⁰		D	1	<0.1 x 10 ⁰		
		2	<0.1 x 10 ⁰			2	<0.1 x 10 ⁰		
	E	1	<0.1 x 10 ⁰		E	1	<0.1 x 10 ⁰		
		2	0.4 x 10 ⁰			2	19.7 x 10 ⁰		
	“Requeijão Light”	A	1		<0.1 x 10 ⁰	“Especialidade láctea” light	A	1	<0.1 x 10 ⁰
			2		<0.1 x 10 ⁰			2	<0.1 x 10 ⁰
B		1	<0.1 x 10 ⁰	B	3		<0.1 x 10 ⁰		
		2	<0.1 x 10 ⁰		4		0.1 x 10 ⁰		
C		1	<0.1 x 10 ⁰	C	1		0.2 x 10 ⁰		
		2	0.1 x 10 ⁰		2		<0.1 x 10 ⁰		
D		1	<0.1 x 10 ⁰	D	3		<0.1 x 10 ⁰		
		2	1.9 x 10 ⁰		4		<0.1 x 10 ⁰		
E		1	<0.1 x 10 ⁰	c	1		<0.1 x 10 ⁰		
		2	<0.1 x 10 ⁰		2		0.1 x 10 ⁰		

it is possible to state that the consumption of these dairy products does not directly pose risk to the Brazilian population, with regard to foodborne illness occurrence. Furthermore, the low number of the pathogen observed in the samples could be explained by the high temperature processes used in the production flowchart (BRAZIL, 1997) and good manufacturing practices implementation in these industries (SANTANA et al., 2009).

A set of 57 isolates were subjected to a PCR in order to confirm if they were *Staphylococcus* spp. (PEREIRA et al., 2009), *S. aureus* (MOROT-BIZOT et al., 2004) or other coagulase-positive *Staphylococcus* spp. (*S. intermedius*, *S. schleiferi* subsp. *coagulans*, *S. hyicus*, *S. lutrae*, *S. delphini*, and *S. pseudintermedius* (SASAKI et al., 2010). In

total 54 isolates were confirmed as *Staphylococcus* spp. with amplification of the corresponding product. There were no amplification products using primers for *Staphylococcus* spp. coagulase-positive, agreeing with the coagulase-test results and allowing us to confirm that all isolates were coagulase-negative strains.

Using PCR to detect the presence of *mecA* gene, one isolate obtained from “requeijão light” (brand C and lot 2) and another isolate obtained from “especialidade láctea type requeijão light” (brand C and lot 2) were considered as positive. The prevalence of methicillin-resistant coagulase-negative *Staphylococcus* spp. in evaluated samples was 3.70% among the 54 isolates. This two isolates showed resistance to oxacillin, penicillin G and

erythromycin but did not to cefepime, rifampicin, chloramphenicol, clindamycin, sulfatrim, tetracycline, gentamicin and ciprofloxacin antimicrobials. Resistance to certain non-beta-lactam (as erythromycin) in *Staphylococcus* spp. isolates carrying *mecA* gene was already reported (HUBER et al., 2011).

NORMANNO et al. (2007) established 3.75% as the prevalence of methicillin-resistant *Staphylococcus* spp. in milk and dairy products in Italy highlighting the role of dairy products in transmission of antimicrobial-resistant bacteria to humans. Furthermore, some authors isolated strains of *Staphylococcus* spp. methicillin-resistant from some food such as: cow's milk, pecorino and mozzarella cheeses (NORMANO et al., 2009) and raw meat (BOER et al., 2009). However, the occurrence of methicillin-resistant *Staphylococcus* spp. in “requeijão” and “especialidade láctea”, or even in any other heat-treated dairy product, was never reported in Brazil, until this paper.

“Requeijão” is one of the most consumed cheeses in Brazil and it is usually consumed raw (ZACARCHENCO et al., 2012). Presence of coagulase-negative methicillin-resistant *Staphylococcus* spp. strains in these products present a risk to the country's public health. Thus, considering the potential implications in food production chain, there is a need to be constantly monitoring the employers' health and hygiene, once humans can be colonized by contact during food processing or consumption of contaminated food (KLUYTMANS, 2009).

Moreover, nowadays coagulase-negative *Staphylococcus* spp. is considered as able on transferring antimicrobial resistance genes to other *Staphylococcus* spp. strains (TULINSKI et al., 2012). Human infections caused by methicillin-resistant *Staphylococcus* spp. are increasing (MARTINS & CUNHA, 2007), even in animals (CUNY et al., 2010), being considered an emergent problem in veterinary medicine and public health (HUBER et al., 2011). Animals usually are infected and colonized during contact with humans and vice-versa (GRAVELAND et al., 2010). Consequently, prophylactic measures to avoid methicillin-resistant *Staphylococcus* spp. contamination in foods and infections in humans and animals are required in order to reduce risk in Brazil.

CONCLUSION

In conclusion, processing technology ensures the safety of these products categories; however, they may be carriers of methicillin resistant

Staphylococcus spp. strains. Results served as a warning to Brazilian public sanitary authorities to control multidrug-resistant strains in veterinary and human medicine.

ACKNOWLEDGMENTS

The authors acknowledge the financial support of this research by the Brazilian research funding agency Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (Process no. 14/21534-6).

REFERENCES

- APHA (AMERICAN PUBLIC HEALTH ASSOCIATION). Committee on Microbiological for Foods. **Compendium of methods for the microbiological examination of foods**. 4.ed. Washington, 2011. 676p.
- ARGUDÍN, M.A et al. Genotypes, exotoxin gene content and antimicrobial resistance in *Staphylococcus aureus* recovered from food and food-handlers. **Applied and Environmental Microbiology**, v.78, n.8, p.2930-2935, 2012. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3318794/>>. Accessed: Jan. 04, 2017. doi: 10.1128/AEM.07487-11.
- BOER, E. et al. Prevalence of methicillin-resistant *Staphylococcus aureus* in meat. **International Journal of Food Microbiology**, v.134, n.1-2, p.52-56, 2009. Available from: <<http://www.sciencedirect.com/science/article/pii/S0168160508006466>>. Accessed: Dec. 29, 2016. doi: 10.1016/j.ijfoodmicro.2008.12.007.
- BRAZIL. Ministry of Supply, Agriculture and Livestock. Portaria nº 359, de 4 de setembro de 1997. **Official Gazzete**, Brasília, DF, Sept. 08, 1997. Section 1, p.19690-19691.
- CUNY C. et al. Emergence of methicillin-resistant *Staphylococcus aureus* (MRSA) in different animal species. **International Journal of Medical Microbiology**, v.300, n.2-3, p.109-117, 2010. Available from: <<http://www.sciencedirect.com/science/article/pii/S1438422109001313>>. Accessed: Dec. 28, 2017. doi: 10.1016/j.ijmm.2009.11.002.
- GORTEL, K. et al. Methicillin resistance among staphylococci isolated from dogs. **American Journal of Veterinary Research**, v.60, n.12, p.1526-1530, 1999. Available from: <<https://www.ncbi.nlm.nih.gov/pubmed/10622162>>. Accessed: Jan. 04, 2017.
- GRAVELAND, H. et al. Methicillin resistant *Staphylococcus aureus* ST398 in veal calf farming: human MRSA carriage related with animal antimicrobial usage and farm hygiene. **Plos One**, v.5, n.6, p.e10990, 2010. Available from: <<http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0010990>>. Accessed: Jan. 03, 2017. doi: 10.1371/journal.pone.0010990.
- HUBER, H. et al. Prevalence and characteristics of methicillin resistant coagulase-negative staphylococci from livestock, chicken carcasses, bulk tank milk, minced meat, and contact persons. **BMC Veterinary Research**, v.7, n.6, p.1-7, 2011. Available from: <<http://bmcvetres.biomedcentral.com/articles/10.1186/1746-6148-7-6>>. Accessed: Dec. 06, 2016. doi: 10.1186/1746-6148-7-6.
- KESKIMAKI, M. et al. EPEC, EAEC and STEC in stool specimens: prevalence and molecular epidemiology of isolates.

- Diagnostic Microbiology & Infectious Disease**, v.40, n.4, p.151-156, 2001. Available from: <[http://www.dmidjournal.com/article/S0732-8893\(01\)00265-6/abstract](http://www.dmidjournal.com/article/S0732-8893(01)00265-6/abstract)>. Accessed: Jan. 04, 2017. doi: 10.1016/S0732-8893(01)00265-6.
- KLUYTMANS, J.A.J.W. Methicillin-resistant *Staphylococcus aureus* in food products: cause for concern or case for complacency? **Clinical Microbiology & Infection**, v.16, n.1, p.11-15, 2009. Available from: <[http://www.clinicalmicrobiologyandinfection.com/article/S1198-743X\(14\)617778/abstract](http://www.clinicalmicrobiologyandinfection.com/article/S1198-743X(14)617778/abstract)>. Accessed: Jan. 01, 2017. doi: 10.1111/j.1469-0691.2009.03110.x.
- KOKSAL, F. et al. Antibiotic resistance patterns of coagulase-negative *Staphylococcus* strains isolated from blood cultures of septicemic patients in Turkey. **Microbiological Research**, v.164, n.4, p.404-410, 2009. Available from: <<http://www.sciencedirect.com/science/article/pii/S0944501307000560>>. Accessed: Jan. 03, 2017. doi: 10.1016/j.micres.2007.03.004.
- MARTINS, A.; CUNHA, M.L.R.S. Methicillin resistance in *Staphylococcus aureus* and Coagulase-Negative Staphylococci: epidemiological and molecular aspects. **Microbiology & Immunology**, v.51, n.9, p.787-795, 2007. Available from: <<https://www.ncbi.nlm.nih.gov/pubmed/17895595>>. Accessed: Dec. 28, 2017. doi: 10.1111/j.1348-0421.2007.tb03968.x.
- MOON, J.S. et al. Phenotypic and genetic antibiogram of Methicillin-resistant Staphylococci isolated from bovine mastitis in Korea. **Journal of Dairy Science**, v.90, n.3, p.1176-1185, 2007. Available from: <<https://www.ncbi.nlm.nih.gov/pubmed/17297092>>. Accessed: Jan. 04, 2017. doi: 10.3168/jds.S0022-0302(07)71604-1.
- MOROT-BIZOT, S.C. et al. Development of a multiplex PCR for the identification of *Staphylococcus* genus and four staphylococcal species isolated from food. **Journal of Applied Microbiology**, v.97, n.5, p.1087-1094, 2004. Available from: <<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2672.2004.02399.x/abstract>>. Accessed: Jan. 04, 2017. doi: 10.1111/j.1365-2672.2004.02399.x.
- NORMANNO, G. et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) in foods of animal origin product in Italy. **International Journal of Food Microbiology**, v.117, n.2, p.219-222, 2007. Available from: <<http://www.sciencedirect.com/science/article/pii/S0168160507002565>>. Accessed: Dec. 22, 2016. doi: 10.1016/j.ijfoodmicro.2007.04.006.
- OTTO, M. Staphylococcal biofilms. **Current Topics in Microbiology and Immunology**, v.322, p.207-228, 2009. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2777538/>>. Accessed: Jan. 04, 2017.
- PEREIRA, V. et al. Characterization for enterotoxin production, virulence factors, and antibiotic susceptibility of *Staphylococcus aureus* isolates from various food in Portugal. **Food Microbiology**, v.26, n.3, p.278-282, 2009. Available from: <<https://www.ncbi.nlm.nih.gov/pubmed/19269569>>. Accessed: Jan. 04, 2017. doi: 10.1016/j.fm.2008.12.008.
- PITKALA, A. et al. Bovine mastitis in Finland 2001 - Prevalence, distribution of bacteria, and antimicrobial resistance. **Journal of Dairy Science**, v.87, n.8, p.2433-2441, 2004. Available from: <<https://www.ncbi.nlm.nih.gov/pubmed/15328265>>. Accessed: Jan. 02, 2017. doi: 10.3168/jds.S0022-0302(04)73366-4.
- SANTANA, N.G. et al. Microbiological quality and safety of meals served to children and adoption of good manufacturing practices in public school catering in Brazil. **Food Control** v.20, n.3, p.255-261, 2009. Available from: <<http://www.sciencedirect.com/science/article/pii/S0956713508001217>>. Accessed: Jan. 03, 2017. doi: 10.1016/j.foodcont.2008.05.004.
- SASAKI, T. et al. Multiplex-PCR method for species identification of coagulase-positive Staphylococci. **Journal of Clinical Microbiology**, v.48, n.3, p.765-769, 2010. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2832457/>>. Accessed: Dec. 29, 2016. doi: 10.1128/JCM.01232-09.
- TULINSKI, P. et al. Methicillin-resistant coagulase-negative Staphylococci on pig farms as a reservoir of heterogeneous Staphylococcal cassette chromosome *mec* elements. **Applied and Environmental Microbiology**, v.78, n.2, p.299-304, 2012. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3255757/>>. Accessed: Jan. 04, 2017. doi: 10.1128/AEM.05594-11.
- VIRGIN, J.E. et al. Short communication: Methicillin-resistant *Staphylococcus aureus* detection in US bulk tank milk. **Journal of Dairy Science**, v.92, n.10, p.4988-4991, 2009. Available from: <<http://www.sciencedirect.com/science/article/pii/S0022030209708306>>. Accessed: Jan. 03, 2017. doi: 10.3168/jds.2009-2290.
- WEESE J.S.; DUJIKEREN E. Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine. **Veterinary Microbiology**, v.140, n.3-4, p.418-429, 2010. Available from: <<https://www.ncbi.nlm.nih.gov/pubmed/19246166>>. Accessed: Dec. 29, 2016. doi: 10.1016/j.vetmic.2009.01.039.
- ZACARCHENCO, P.B et al. "Requeijão" culinário: Historic, market and technological aspects. **Revista das Indústrias de Laticínios**, v.18, p.70-74, 2012. Available from: <<http://revistalaticinios.com.br/wp-content/uploads/2011/09/56-76-Fazer-Melhor.pdf>>. Accessed: Dec. 23, 2016.