

Episodic memory improvement in illiterate adults attending late-life education irrespective of low socioeconomic status: insights from the PROAME study

Emma Patrice Ruppert^{1,2}, João Victor de Faria Rocha^{3,4}, Aída Lourandes da Silva¹,
Kelle Luisa dos Santos Tomaz³, Clarisse Vasconcelos Friedlaender^{3,4},
Joanna de Castro Magalhães Assenção³, Luciana Paula Rincon³, Norton Gray Ferreira Ribeiro³,
Dulce Constantina de Souza Santos⁵, Ana Paula Zacarias Lima⁶, Isabel Elaine Allen⁷, Paulo Caramelli³,
Lea Tenenholz Grinberg^{7,8}, Francisca Izabel Pereira Maciel³, Elisa de Paula França Resende^{1,3,7}

ABSTRACT. The majority of people with dementia live in low or middle-income countries (LMICs) where resources that play a crucial role in brain health, such as quality education, are still not widely available. In Brazil, illiteracy remains a prevalent issue, especially in communities with lower socioeconomic status (SES). The PROAME study set out to explore basic education in illiterate adults as a means to improve cognitive reserve. **Objective:** This manuscript aims to explore the relationship between SES and learning, as well as cognitive outcomes, in an older illiterate population. **Methods:** This six-month clinical trial (NCT04473235) involved 108 participants, of which 77 concluded all assessments, enrolled in late-life basic education. SES assessments included Quality of Urban Living Index, Municipal Human Development Index and Household SES calculated for each participant. Cognitive assessments encompassed the Free and Cued Selective Reminding Test (FCSRT), a word list to assess reading, and the Beta III matrix. **Results:** The sample consisted primarily of women, with a mean age of 58.5. Participants improved their reading ($p=0.01$) and their FCSRT ($p=0.003$). Regarding episodic memory, women outperformed men ($p=0.007$) and younger participants improved more than their older counterparts ($p=0.001$). There was no association observed between SES and cognitive outcomes. **Conclusion:** Irrespective of SES, participants demonstrated positive outcomes after attending basic education. These findings highlight that late life education could be an important non-pharmacologic preventative measure, especially in LMICs.

Keywords: Socioeconomic Factors; Cognitive Reserve; Memory, Episodic; Dementia; Literacy; Cognitive Aging.

Melhoria da memória episódica em adultos analfabetos que frequentam educação tardia, independentemente do baixo nível socioeconômico: considerações do estudo PROAME

RESUMO. A maioria das pessoas com demência vive em países de baixa/média renda, onde recursos essenciais para a saúde cerebral, como educação de qualidade, ainda não são amplamente acessíveis. No Brasil, o analfabetismo ainda é frequente, especialmente em comunidades de baixo nível socioeconômico. O estudo PROAME teve como objetivo explorar a educação básica tardia em pessoas analfabetas como ferramenta para o aumento da reserva cognitiva. **Objetivo:** Investigar a relação

This study was conducted by the Behavioral and Cognitive Neurology Unit, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil.

¹Faculdade de Ciências Médicas de Minas Gerais, Belo Horizonte MG, Brazil.

²University of Pittsburgh, Pascoal Lab, Pittsburgh PA, USA.

³Universidade Federal de Minas Gerais, Faculdade de Medicina, Grupo de Pesquisa Neurologia Cognitiva e do Comportamento, Belo Horizonte MG, Brazil.

⁴Universidade Federal de Minas Gerais, Faculdade de Medicina, Programa de Pós-Graduação em Ciências Aplicadas à Saúde do Adulto, Belo Horizonte MG, Brazil.

⁵Escola Municipal Acadêmico Vivaldi, Belo Horizonte MG, Brazil.

⁶Escola Municipal Dr. Júlio Soares, Belo Horizonte MG, Brazil.

⁷University of California San Francisco, Global Brain Health Institute, San Francisco CA, USA.

⁸University of California, Memory and Aging Center, UCSF Weill Institute for Neurosciences, San Francisco CA, USA.

Correspondence: Emma Ruppert; Email: ruppertemmap@gmail.com.

Disclosure: The authors report no conflicts of interest.

Funding: none.

Received on October 06, 2023; Received in its final form on January 29, 2024; Accepted on March 03, 2024.



entre nível socioeconômico com aprendizado e com desempenho em testes cognitivos, em adultos analfabetos. **Métodos:** Este estudo clínico de seis meses (NCT04473235) contou com 108 participantes inscritos no projeto Educação para Jovens e Adultos (EJA), dos quais 77 completaram os testes. O nível socioeconômico de cada participante foi medido usando-se: o Índice de Qualidade de Vida Urbana, o Índice de Desenvolvimento Humano Municipal e o nível socioeconômico doméstico. Avaliações cognitivas incluíram: o Teste de Recordação Seletiva Livre e Guiada (TRSLG), uma lista de palavras para avaliar leitura e a matriz Beta III. **Resultados:** A amostra era predominantemente feminina, com idade média de 58,5. Os participantes melhoraram a leitura ($p=0,01$) e o TRSLG ($p=0,003$). Com relação à memória episódica, as mulheres tiveram resultados superiores aos dos homens ($p=0,007$) e participantes mais jovens melhoraram mais que seus colegas mais velhos ($p=0,001$). Não foi observada nenhuma relação entre o nível socioeconômico e o desempenho cognitivo. **Conclusão:** Independentemente do nível socioeconômico, participantes obtiveram resultados positivos após frequentar a educação básica. Isso sugere que a educação tardia pode ser uma medida preventiva não farmacológica importante, especialmente em países de baixa/média renda.

Palavras-chave: Fatores Socioeconômicos; Reserva Cognitiva; Memória Episódica; Demência; Alfabetização; Envelhecimento Cognitivo.

INTRODUCTION

Approximately 57 million individuals are currently living with dementia, and projections indicate that this figure will triple by 2050¹⁻⁴. Dementia is the primary cause of disability in high-income countries (HICs), but the majority of those affected reside in low or middle-income countries (LMICs)⁴. This discrepancy is primarily attributed to inequalities in access to resources that play a crucial role in brain health, such as quality education^{1,2}. Notably, dementia can manifest up to a decade earlier in LMICs due to these inequalities⁵. Moreover, people with dementia who have low education manifest a more widespread cognitive impairment in comparison to those with higher schooling⁶. To mitigate the impact of these imbalances on brain health, it is imperative that they be addressed through targeted strategies, tailored to nations with lower economic means.

Approximately one third of Alzheimer's disease cases can be attributed to underlying modifiable risk factors⁴. Control of the following factors can prevent as much as 40% of dementia cases: low education, hearing loss, hypertension, obesity, diabetes, alcohol abuse, traumatic brain injury (TBI), physical inactivity, depression, smoking, social isolation, and air pollution^{2,3,6,7}. In Brazil, these factors account for up to 50.5% of dementia cases, especially low educational attainment and hearing loss^{2,3}.

Due to the lifestyle-related nature of many preventable and protective factors, their impact is disproportionately felt in socioeconomically deprived regions⁵. Access to resources, including healthy diet, safety, education, healthcare services, physical activity, and even social factors like isolation and exclusion can pose significant challenges in communities with lower SES^{2,4,8,9}. Paradoxically, much of the research into the role of these factors in dementia prevention has been conducted in HICs that do not face these same challenges^{2,4,8,9}.

In 2022, 9.6 million Brazilians aged over 15 were illiterate, with 16% of the elderly population being

illiterate¹⁰. Importantly, older adults with reduced literacy skills tend to have poorer health outcomes and an elevated risk of mortality¹¹. Investigating prevention strategies tailored to individuals with low education levels is crucial for reducing the incidence and impact of dementia, especially in LMICs.

This work is part of a larger research project, the PROAME study, that aims to investigate the significance of acquiring basic education in later stages of life as a means to enhance cognitive reserve. This manuscript explores the relationship between socioeconomic status and learning in an older illiterate vulnerable population.

METHODS

Participants

This longitudinal clinical trial involved the screening of 130 adults who were actively participating in basic education classes offered by the Brazilian government under the program "Education for Teens and Adults" (*Educação de Jovens e Adultos*). The inclusion criteria mandated participants to be illiterate (Test of Functional Health Literacy in Adults — TOFHLA score <53), and available to take part in the study for a minimum of 12 months¹²⁻¹⁴. Exclusion criteria included: significant cognitive complaints, unmanaged psychiatric conditions, ongoing substance abuse, evident cognitive impairment (defined as a score of ≤ 2 standard deviations for age and education on the Mini-Mental State Examination and/or a score of ≤ 6 on the delayed recall task from the Brief Cognitive Screening Battery), past medical history of dyslexia, contraindications to formal MRI procedures and MRI-detected structural brain lesions¹⁵⁻¹⁹.

Following these criteria, a total of 108 participants were enrolled in the study. Data collection occurred at baseline and after six months of participation in basic education classes. At baseline, participants also answered the Cognitive Reserve Index questionnaire

(CRIq)²⁰. Figure 1 illustrates the flow of participants through the study, ultimately showing that 77 participants successfully completed all required clinical assessments and interviews both at baseline and after the educational intervention.

Ethics

This study is part of a larger research program titled Better Memory with Literacy Acquisition Later in Life (NCT04473235) and was approved by the Ethics Committee of the Universidade Federal de Minas Gerais. Informed consent was obtained prior to data collection from all participants.

Socioeconomic status assessment (SES)

Quality of Urban Living Index (QULI)

Created by the city of Belo Horizonte, this index serves as a comprehensive measure encompassing access to basic resources, cultural amenities, educational opportunities, sports facilities, housing conditions, urban infrastructure, environmental factors, healthcare services, public transportation and other urban services and security/safety within specific microregions of the city. These microregions make up 80 different areas. The index is available on the city hall's official website and is graded on a scale from 0 (lowest) to 1 (highest). The most recent version is from 2016²¹. This assessment offers a valuable insight into the region. It is important to note that due to evolving social inequalities there is the possibility of misrepresentation, particularly for communities with lower SES situated within regions containing more affluent areas. This index was obtained from the 92 participants with verifiable addresses in

Belo Horizonte. It revealed that these individuals were distributed across 20 different microregions within the city. This piece of data was missing for 16 participants.

The participants had an average score of 0.67, which is similar to the city-wide average of 0.63²¹. During the statistical analysis, the QULI score was treated as a continuous variable. However, for the purpose of discussion, we classified the measure using the median value of 0.67. This categorization resulted in a dichotomous distinction, for a more practical analysis — scores equal to or greater than the median were considered “high,” while scores lower than the median were considered “low”.

Municipal Human Development Index (MHDI)

A collaboration between the United Nations Development Program, the Brazilian Institute of Applied Economic Research, and the João Pinheiro Foundation, this index is made accessible through the interactive platform Atlas Brasil. It utilizes the Human Development Index (HDI) criteria: education, life expectancy, and gross national income per capita. This measure was collected for 92 participants. For statistical purposes this index was used as a continuous variable. During the discussion, we further divided the participants according to the HDI classification system as: very low (0–0.49), low (0.50–0.59), medium (0.60–0.69), high (0.70–0.79), to very high (0.80–1).²²

A distinct advantage of this measure lies in its international recognition. However, the data used for its calculation is based on the last national census conducted in 2010, potentially rendering it less reflective of the current conditions.

Household Socioeconomic Status (SES)

The SES of each participant's household was determined using Brazil's Economic Classification Criteria. These criteria are nationally recognized and use a range of factors from a single household to estimate SES. These factors include quality of living conditions (such as number of bathrooms, electronic devices, automobiles, and household appliances), educational level of the primary provider, and availability of essential public services (access to treated water and paved roads)²³. Each of these factors is assigned a point. These points are then tallied to yield a cumulative score that corresponds to a socioeconomic level — A, B1, B2, C1, C2, or D-E. Each level is associated with an approximate monthly income, in US dollars, of: \$4,559 for level A, \$2,092.54 for B1, \$1,093.57 for B2, \$610.53 for C1, \$362.39 for C2, and \$163.26 for D-E²³. This data was missing for 19 participants, that did not complete the full assessment.




Figure 1. Flowchart depicting participants included in this study.

This tool offers the advantage of facilitating a tailored analysis of income and economic circumstances, providing a personalized insight into participants' SES. However, this approach does not account for the way in which the area a person lives in can influence their access to essential services, safety, and overall socioeconomic standing.

Cognitive assessments

Cognitive assessments were performed at baseline and after six months of education. The Free and Cued Selective Reminding Test (FCSRT), also referred to as delayed-recall episodic memory scores, was used for episodic memory^{24,25}.

Reading skill evaluation was conducted by presenting the participants with a list of words devised by Rodrigues et al.²⁶. This list consisted of 48 words to be read by the participants. Correctly pronouncing a word awarded a point, while an incorrect pronunciation resulted in zero points. Lastly, the Beta III matrix was employed to gauge non-verbal reasoning skills among the participants²⁷.

Intervention

The participants were categorized into two distinct groups based on their school enrollment. The control group (n=50) engaged in regular basic education classes, dedicating three hours each day for four days a week over a period of six months. The subjects covered in these classes included geography, mathematics, and history; however, these classes did not target literacy training. In contrast, the intervention group (n=58) attended, in addition to these basic education classes, intensive literacy instruction facilitated by a skilled educator, which was focused on how to read and write. Further characterization of these groups can be found in the Supplementary Material (<https://www.demneuropsy.com.br/wp-content/uploads/2024/03/DN-2023.0098-Supplementary-Material.docx>). This paper will not delve into the specifics of these subgroups, and this matter will be explored in another publication.

Statistical analysis

The statistical analyses were carried out using the open software Jamovi Version 2.3.21.0. Descriptive and continuous variables were summarized using measures such as the mean, median, and interquartile ranges (IQR). Categorical variables were assessed in terms of frequency. During the analytical phase, QULI and MHDI were used as continuous variables. However, to provide further context, in the discussion, these SES measures were categorized into groups.

To examine the relationship between SES and TBI, simple linear regression analyses were conducted for QULI and MHDI, as well as Fisher's exact test to determine if there was a significant association between Household SES and TBI. The exploration of potential associations between SES measures and improvements in episodic memory and reading skills was conducted using a pair of mixed linear models. The first model used the sum of attempts of FCSRT as the dependent variable. The model incorporated several covariates — QULI, MHDI, Household SES, Beta III, years of school attended as a child, age, sex, and timeline. The second model utilized the number of words read correctly and mirrored the covariates of the first model.

RESULTS

The sample consisted of 108 participants, of which 77 successfully completed the cognitive assessments. The average age was 58.5 (standard deviation — SD=9.64). The majority were women (74.1%). In terms of race, most participants identified as black/brown, as shown in Table 1.

In terms of education, most participants (61.8%) reported attending school as a child and the average number of years of schooling, including during childhood and adulthood, was 1.9 years (SD=2.1). When questioned about their illiteracy, the primary explanations given by most participants were childhood labor (31.5%), followed by growing up in rural communities with limited school access (19.4%), learning difficulties (13.0%) and lack of parental encouragement (12.0%).

Regarding the CRIq scores in our sample, the mean was 72.5 (SD=6.8). When considering the different categories, the mean scores were as follows: education 76.6 (SD=10.6), work 87.1 (SD=6.4), and leisure 74.0 (SD=9.8). These scores are all in the medium-low category for cognitive reserve²⁰.

Regarding SES, the participants' QULI scores closely resembled the city-wide mean (M=0.67, SD=0.06, IQR=0.09), while the mean MHDI was categorized as high (M=0.73, SD=0.08, IQR=0.12), the Household SES revealed that most participants had a low income, as seen in Table 1.

Twenty-nine individuals reported having experienced a previous TBI, with an average of 27.9 years since the trauma occurred (SD=20.4, [0.16,60]). Information about previous TBI was missing for 19 participants. Approximately 41.4% of these individuals indicated seeking medical attention at a hospital

Table 1. Characteristics of the sample.

Characteristics	Frequency (%)
Sex	
Women	80 (74.1)
Men	28 (25.9)
Race*	
Black/brown	78 (88.6)
White	9 (10.2)
Indigenous	1 (1.1)
Mental health diagnosis	
Anxiety	33 (37.1)
Depression	18 (20.2)
Tobacco[†]	
Current use	6 (6.7)
Past use	40 (44.9)
Reported health conditions[†]	
Hypertension	46 (59.7)
Diabetes mellitus	22 (28.57)
Hearing difficulties	21 (27.27)
Attended school as a child[†]	
Yes	55 (61.8)
No	34 (38.2)
QULI[‡]	
Low	46 (50.0)
High	46 (50.0)
MHDI[‡]	
Medium	37 (40.2)
High	33 (35.9)
Very high	22 (23.9)
Personal SES[‡]	
B	8 (9.00)
C1	14 (15.7)
C2	34 (38.2)
D-E	33 (37.1)

Abbreviations: QULI, Quality of Urban Living Index; MHDI, Municipal Human Development Index; SES, Social Economic Status.

Notes: *missing = 20; [†]missing = 19; [‡]missing = 16.

following the injury, and twelve reported losing consciousness during the event. The results of linear regression analyses revealed that traumatic brain injury exhibited an association with lower QULI ($\beta=0.024$, $p=0.026$, 95%CI 0.003–0.045) and a lower MHDI ($\beta=0.03$, $p=0.018$, 95%CI 0.006–0.06). Fisher's exact test showed an association between TBI and Household SES ($p=0.046$).

In respect to cognitive outcomes, after study completion participants exhibited a significant improvement in their episodic memory performance compared to baseline. However, this improvement was not different between control and intervention subgroups, which will be explored in another publication.

Notably, the factors of sex and age had significant effects on this improvement. Men demonstrated a lesser degree of improvement compared to women. Similarly, older participants displayed less improvement compared to their younger counterparts. In contrast, Beta III, years of school attended as a child, QULI, MHDI, Household SES did not have a significant impact on episodic memory improvement, as indicated in Table 2.

Turning to participants' reading abilities, an overall enhancement was observed after attending classes compared to initial testing. No other factors showed significant impact, as depicted in Table 3.

Table 2. Results of mixed linear regression on delayed-recall episodic memory scores.

	β	p-value	95%CI
Timeline	3.122	0.003*	1.17; 5.08
Sex	-4.327	0.007*	-7.35; -1.30
Age	-0.178	<0.001*	-0.33; -0.03
Years of school attended as a child	0.096	0.740	-0.47; 0.66
Beta III	-0.154	0.645	-0.80; 0.49
QULI	7.673	0.463	-12.71; 28.05
MHDI	1.465	0.860	-14.80; 17.73
Household SES level			
B1–C2	-2.216	0.554	-9.52; 5.08
B2–C2	3.507	0.152	-1.26; 8.47
C1–C2	-0.126	0.947	-3.85; 3.59
D–E–C2	-0.402	0.786	-3.29; 2.48

Abbreviations: QULI, Quality of Urban Living Index; MHDI, Municipal Human Development Index; SES, Social Economic Status.

Note: * $p<0.05$.

DISCUSSION

Disparities in the accessibility of preventive measures should contribute to the surge in dementia cases over the next three decades, particularly in LMICs^{2,3,5}. While the protective influence of childhood education against dementia is well-documented, the potential impact of acquiring basic education later in life remains largely unexplored. This paper explored the role

of SES in relation to learning in a sample of illiterate adults (Figure 2).

Our study findings showed significant improvement in both episodic memory and reading skills through late-life basic education. Secondly, age and sex were factors that affected episodic memory improvement. Lastly, our findings highlight that learning as an adult is an effective tool irrespective of low SES.

While our study primarily examined the impact of basic education on the brain health of illiterate adults, the findings are complemented by existing research on the role of continued education in late life. Notably, the Tasmanian Healthy Brain Project, a longitudinal study on cognitively unimpaired adults, demonstrated that individuals engaged in university-level schooling experienced greater improvements in verbal memory, verbal episodic memory, and episodic memory over a span of seven years compared to those who did not engage in schooling²⁸. The novelty of our study lies in demonstrating similar benefits from basic education, suggesting that literacy programs tailored for illiterate adults can enhance brain health. This underscores the importance of offering public literacy programs for adults, especially in LMICs where illiteracy is still prevalent.

As expected, women had a greater improvement in episodic memory. Although the exact reason remains unclear, it is widely acknowledged that there exists a female advantage in verbal episodic memory^{29,30}. The observed influence of age on episodic memory improvement also aligns with expectation, as age leads to a decline in fluid cognitive abilities. The decline typically commences around the age of 60 and accelerates in the

Table 3. Results of mixed linear regression on reading.

	B	p-value	95%CI
Timeline	4.277	0.010*	1.14; 7.41
Sex	-7.45	0.184	-18.30; 3.40
Age	0.259	0.725	-0.27; 0.78
Beta III	1.621	0.179	-0.72; 3.96
QULI	-13.060	0.725	-85.51; 59.34
MHDI	51.120	0.088	-6.66; 108.90
Years of school attended as a child	-0.550	0.597	-2.58; 1.48
Household SES level			
B1–C2	8.686	0.164	-3.37; 20.75
B2–C2	6.686	0.110	-1.37; 14.74
C1–C2	2.69	0.415	-3.71; 9.09
D–E–C2	1.757	0.514	-3.71; 9.09

Abbreviations: QULI, Quality of Urban Living Index; MHDI, Municipal Human Development Index; SES, Social Economic Status.

Note: *p<0.05.




Figure 2. Visual abstract summarizing the study findings.

subsequent decade³¹. Age might exert an even more pronounced impact than evident in our study, since the majority of participants were under the age of 70. This could indicate that public education programs for illiterate adults may be further optimized if started at an earlier age.

In regard to SES, our sample exhibited a seemingly high SES based on area of residency, such as QULI and MHDI, yet a low SES with regard to household income. Most participants fell within economic classes C1, C2 and D-E, corresponding to monthly incomes below the national income average of USD \$632.21. This highlights that, despite residing in areas with available resources, participants had low personal SES, which could affect their utilization of these resources. Our findings indicate that education can benefit people from low and medium socioeconomic backgrounds alike. Additionally, our results showed an association between lower SES and previous TBI. And while the relationship between race and insurance status on TBI outcomes has been described, there is still little research exploring previous TBI and current SES, especially in Latin America^{32,33}.

It is possible that the improvement seen in the participants was not solely due to education, but also due to the reduction in loneliness and in depression symptoms. While the role of loneliness and depression has been well documented as a risk factor for cognitive decline and dementia, there is still little research on how decreasing these factors could lead to better cognition^{2-4,7}.

This study had a few limitations: a small sample size, potential for floor effects, as well as a relatively short time frame. Given that illiterate individuals often have lower overall SES, the sample may have been relatively homogeneous. This could limit the observed impact of SES on educational outcomes. Furthermore, due to the predominantly female composition of the sample, generalizing gender effects on a larger scale remains challenging. A notable challenge encountered was that some participants did not have a consistent attendance

rate, which explains the difference in sample sizes depending on the day the data was collected.

This study offers valuable insights into the potential of late-life basic education as a public health strategy for addressing the global dementia burden, particularly in LMICs where illiteracy remains a significant concern. Importantly, it demonstrates that individuals from diverse socioeconomic backgrounds can benefit from learning interventions, especially if started at a younger age. These findings highlight an important non-pharmacologic preventative measure that could be very useful in LMICs. To strengthen these findings and deepen our understanding of the preventive role of late-life education in dementia, further investigations, specifically focusing on illiterate older adults, are warranted.

AUTHORS' CONTRIBUTIONS

EPR: data curation, formal analysis, investigation, visualization, writing – original draft, writing – review & editing. JVFR: data curation, formal analysis, investigation, project administration, writing – review & editing. ALS: formal analysis, investigation, writing – review & editing. KLST: investigation, project administration, writing – review & editing. CVF: investigation, methodology, validation, writing – review & editing. JCMA: investigation, writing – review & editing. LPR: investigation, writing – review & editing. NGFR: investigation, writing – review & editing. DCSS: investigation, resources. APZL: investigation, writing – review & editing. IEA: formal analysis, validation. PC: conceptualization, investigation, methodology, resources, supervision, writing – review & editing. LTG: conceptualization, supervision, writing – review & editing. FIPM: investigation, methodology, resources, supervision, writing – review & editing. EPFR: conceptualization, data curation, formal analysis, investigation, methodology, project administration, resources, supervision, validation, writing – review & editing.

REFERENCES

- Vassilaki M, Petersen RC, Vemuri P. Area deprivation index as a surrogate of resilience in aging and dementia. *Front Psychol*. 2022;13:930415. <https://doi.org/10.3389/fpsyg.2022.930415>
- Suemoto CK, Mukadam N, Brucki SMD, Caramelli P, Nitrini R, Laks J, et al. Risk factors for dementia in Brazil: differences by region and race. *Alzheimers Dement*. 2023;19(5):1849-57. <https://doi.org/10.1002/alz.12820>
- Borelli WV, Leotti VB, Strelow MZ, Chaves MLF, Castilhos RM. Preventable risk factors of dementia: population attributable fractions in a Brazilian population-based study. *Lancet Reg Health Am*. 2022;11:100256. <https://doi.org/10.1016/j.lana.2022.100256>
- Zhang XX, Tian Y, Wang ZT, Ma YH, Tan L, Yu JT. The epidemiology of Alzheimer's disease modifiable risk factors and prevention. *J Prev Alzheimers Dis*. 2021;8(3):313-21. <https://doi.org/10.14283/jpad.2021.15>
- Resende EPF, Guerra JLL, Miller BL. Health and socioeconomic inequities as contributors to brain health. *JAMA Neurol*. 2019;76(6):633-4. <https://doi.org/10.1001/jamaneurol.2019.0362>
- Caramelli P, Poissant A, Gauthier S, Bellavance A, Gauvreau D, Lecours AR, et al. Educational level and neuropsychological heterogeneity in dementia of the Alzheimer type. *Alzheimer Dis Assoc Disord*. 1997;11(1):9-15. <https://doi.org/10.1097/00002093-199703000-00003>

7. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. 2020;396(10248):413-46. [https://doi.org/10.1016/S0140-6736\(20\)30367-6](https://doi.org/10.1016/S0140-6736(20)30367-6)
8. Santamaria-Garcia H, Sainz-Ballesteros A, Hernandez H, Moguilner S, Maito M, Ochoa-Rosales C, et al. Factors associated with healthy aging in Latin American populations. *Nat Med*. 2023;29(9):2248-58. <https://doi.org/10.1038/s41591-023-02495-1>
9. Nitrini R, Barbosa MT, Brucki SMD, Yassuda MS, Caramelli P. Current trends and challenges on dementia management and research in Latin America. *J Glob Health*. 2020;10(1):010362. <https://doi.org/10.7189/jogh.10.010362>
10. Instituto Brasileiro de Geografia e Estatística. Pesquisa Nacional por Amostra de Domicílios Contínua. Educação 2022 [Internet]. [cited on Aug 30, 2023] Available from: https://biblioteca.ibge.gov.br/visualizacao/livros/liv102002_informativo.pdf
11. Kaup AR, Simonsick EM, Harris TB, Satterfield S, Metti AL, Ayonayon HN, et al. Older adults with limited literacy are at increased risk for likely dementia. *J Gerontol A Biol Sci Med Sci*. 2014;69(7):900-6. <https://doi.org/10.1093/geron/glt176>
12. Apolinario D, Mansur LL, Carthery-Goulart MT, Brucki SM, Nitrini R. Cognitive predictors of limited health literacy in adults with heterogeneous socioeconomic backgrounds. *J Health Psychol*. 2015;20(12):1613-25. <https://doi.org/10.1177/1359105313520337>
13. Maragno CAD, Mengue SS, Moraes CG, Rebelo MVD, Guimarães AMM, Dal Pizzol TS. Test of health Literacy for Portuguese-speaking adults. *Rev Bras Epidemiol*. 2019;22:e190025. <https://doi.org/10.1590/1980-549720190025>
14. Parker RM, Baker DW, Williams MV, Nurss JR. The test of functional health literacy in adults: a new instrument for measuring patients' literacy skills. *J Gen Intern Med*. 1995;10(10):537-41. <https://doi.org/10.1007/BF02640361>
15. Brucki SMD, Nitrini R, Caramelli P, Bertolucci PHF, Okamoto IH. Sugestões para o uso do mini-exame do estado mental no Brasil. *Arq Neuropsiquiatr*. 2003;61(3B):777-81. <https://doi.org/10.1590/s0004-282x2003000500014>
16. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189-98. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
17. Nitrini R, Lefèvre BH, Mathias SC, Caramelli P, Carrilho PEM, Sauaia N, et al. Testes neuropsicológicos de aplicação simples para o diagnóstico de demência. *Arq Neuropsiquiatr*. 1994;52(4):457-65. <https://doi.org/10.1590/s0004-282x1994000400001>
18. Nitrini R, Caramelli P, Herrera Júnior E, Porto CS, Charchat-Fichman H, Carthery MT, et al. Performance of illiterate and literate nondemented elderly subjects in two tests of long-term memory. *J Int Neuropsychol Soc*. 2004;10(4):634-8. <https://doi.org/10.1017/S1355617704104062>
19. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry*. 1998;59. Suppl 20:22-33; quiz 34-57. PMID: 9881538.
20. Nucci M, Mapelli D, Mondini S. Cognitive Reserve Index questionnaire (CRIq): a new instrument for measuring cognitive reserve. *Aging Clin Exp Res*. 2012;24(3):218-26. <https://doi.org/10.3275/7800>
21. Prefeitura de Belo Horizonte Índice de Qualidade de Vida Urbana (IQ-VU-BH) [Internet]. 2016 [cited on Sep 28, 2023]. Available from: <https://prefeitura.pbh.gov.br/estatisticas-e-indicadores/indice-de-qualidade-de-vida-urbana>.
22. Atlas do Desenvolvimento Humano no Brasil. Como está o desenvolvimento humano no Brasil? [Internet]. Atlas Brasil; 2021 [cited on July 06, 2023]. Available from: <http://www.atlasbrasil.org.br>
23. Associação Brasileira de Empresas de Pesquisa. Critério de Classificação Econômica Brasil. Alterações na aplicação do Critério Brasil, válidas a partir de 01/09/2020 [Internet] 2019. [cited on Aug 30, 2023]. Available from: https://www.abep.org/criterioBr/01_cceb_2020.pdf
24. Grober E, Sanders AE, Hall C, Lipton RB. Free and cued selective reminding identifies very mild dementia in primary care. *Alzheimer Dis Assoc Disord*. 2010;24(3):284-90. <https://doi.org/10.1097/WAD.0b013e-3181cfc78b>
25. Zibetti MR, Bordignon S, Trentini CM. Memória e aprendizagem no procedimento de recordação seletiva livre e com pistas. *Temas Psicol*. 2014;22(4):771-82. <https://doi.org/10.9788/TP2014.4-08>
26. Rodrigues JC, Nobre AP, Gauer G, Salles JF. Construção da tarefa de leitura de palavras e pseudopalavras (TLPP) e desempenho de leitores proficientes. *Temas Psicol*. 2015;23(2):413-29. <https://doi.org/10.9788/TP2015.2-13>
27. Rabelo IS, Pacanaro SV, Leme IFAS, Ambiel RAM, Alves, GDS. Teste não verbal de inteligência geral-BETA III-subtestes raciocínio matricial e códigos. São Paulo: Casa do Psicólogo; 2011.
28. Bindoff AD, Summers MJ, Hill E, Alty J, Vickers JC. Studying at university in later life slows cognitive decline: a long-term prospective study. *Alzheimers Dement (NY)*. 2021;7(1):e12207. <https://doi.org/10.1002/trc2.12207>
29. Asperholm M, van Leuven L, Herlitz A. Sex differences in episodic memory variance. *Front Psychol*. 2020;11:613. <https://doi.org/10.3389/fpsyg.2020.00613>
30. PLOS ONE Staff. Correction: The magnitude of sex differences in verbal episodic memory increases with social progress: data from 54 countries across 40 years. *PLoS One*. 2019;14(5):e0217033. <https://doi.org/10.1371/journal.pone.0217033>
31. Thow ME, Summers MJ, Saunders NL, Summers JJ, Ritchie K, Vickers JC. Further education improves cognitive reserve and triggers improvement in selective cognitive functions in older adults: The Tasmanian Healthy Brain Project. *Alzheimers Dement (Amst)*. 2017;10:22-30. <https://doi.org/10.1016/j.dadm.2017.08.004>
32. Magalhães ALG, Barros JLVM, Cardoso MGF, Rocha NP, Faleiro RM, Souza LC, et al. Traumatic brain injury in Brazil: an epidemiological study and systematic review of the literature. *Arq Neuropsiquiatr*. 2022;80(4):410-23. <https://doi.org/10.1590/0004-282X-ANP-2021-0035>
33. Haines KL, Nguyen BP, Vatsaas C, Alger A, Brooks K, Agarwal SK. Socioeconomic status affects outcomes after severity-stratified traumatic brain injury. *J Surg. Res*. 2019;235:131-40. <https://doi.org/10.1016/j.jss.2018.09.072>