

PATHOTYPE DIVERSITY OF *Pyricularia grisea* FROM IMPROVED UPLAND RICE CULTIVARS IN EXPERIMENTAL PLOTS

ANNE S. PRABHU¹, MARTA C. FILIPPI¹ & LEILA G. ARAUJO^{2*}

¹Embrapa Arroz e Feijão, Cx. Postal 179, CEP 75375-000, Santo Antônio de Goiás, GO, Brazil, Fax (062) 533-2172, e-mail: prabhu@cnpaf.embrapa.br; ²Escola de Agronomia, Universidade Federal de Goiás, Cx. Postal 131, CEP 74001-970, Goiânia, GO, e-mail: leilag@cnpaf.embrapa.br

(Accepted for publication 12/08/2002)

Corresponding author: Anne Sitarama Prabhu

PRABHU, A.S., FILIPPI, M.C. & ARAÚJO, L.G. Pathotype diversity of *Pyricularia grisea* from improved upland rice cultivars in experimental plots. *Fitopatologia Brasileira* 27:468-473. 2002.

ABSTRACT

A study was undertaken to examine the pathogenic diversity of *Pyricularia grisea* isolates retrieved from 14 upland rice (*Oryza sativa*) cultivars in experimental plots during a period of five years. Inoculations were performed on 32 genotypes with 85 monoconidial isolates under controlled greenhouse conditions. Based on the reaction pattern of eight international differentials, eleven pathotypes of *P. grisea* were identified. The predominant international races or pathotypes were IB-9 (56.4%), IB-1 (16.4%) and IB-41 (11.8%). A set of eight commercial upland rice cultivars ('Carajás', 'Confiança', 'Maravilha', 'Primavera', 'Progresso', 'Caiapó', 'IAC-47', 'IAC-201') was utilized as additional differentials for describing the virulence pattern of *P. grisea*. Twenty-six Brazilian pathotypes were identified on the basis of disease reaction on these differentials, in contrast to the 11 international pathotypes. The most predominant

Brazilian pathotypes, BB-21 and BB-41 were represented by 28.2% and 17.6% of the isolates tested, respectively. Isolates virulent and avirulent to cultivar 'Primavera' were encountered within the pathotype IB-1. Utilizing Brazilian cultivars as differentials, the 14 isolates of the pathotype IB-1 could be further classified into eight local pathotypes, BB-41, BB-13, BB-21, BB-9, BB-29, BB-61, BD-9 and BG-1. Virulence to improved rice cultivars 'Canastra', 'Confiança', 'Carisma', 'Maravilha', 'Primavera' and 'Bonança' was frequent in pathogen population. Some of the Brazilian pathotypes that showed differential reaction on commercial rice cultivars could be utilized for incorporating resistance genes in susceptible cultivars improved for grain quality, by conventional breeding methods.

Additional key words: *Oryza sativa*, *Magnaporthe grisea*, physiologic races, differential varieties, resistance, virulence.

RESUMO

Diversidade de patótipos de *Pyricularia grisea* provenientes de cultivares de arroz de terras altas em parcelas experimentais

Foi estudada a diversidade patogênica de isolados de *Pyricularia grisea* coletados de 14 cultivares de arroz (*Oryza sativa*) de terras altas, em campos experimentais durante um período de cinco anos no Brasil. Foram realizadas inoculações em 32 genótipos com 85 isolados monospóricos em condições de casa de vegetação. Com base nas reações das oito diferenciadoras internacionais, onze patótipos de *P. grisea* foram identificados. Os patótipos internacionais predominantes foram IB-9 (56.4%), IB-1 (16.4%) e IB-41 (11.8%). Oito cultivares locais ('Carajás', 'Confiança', 'Maravilha', 'Primavera', 'Progresso', 'Caiapó', 'IAC-47' e 'IAC-201') foram utilizadas como diferenciadoras para descrever o padrão de virulência de isolados de *P. grisea*. Foram identificados 26 patótipos locais com base no tipo de reação nestas diferenciadoras, em contraste com os

11 patótipos internacionais. Os patótipos brasileiros mais predominantes, BB-21 e BB-41, representaram 28,2% e 17,6%, dos isolados testados, respectivamente. Isolados virulentos e avirulentos com a cultivar 'Primavera' foram encontrados dentro do patótipo IB-1. Os 14 isolados do patótipo IB-1 foram subdivididos em oito patótipos locais, BB-41, BB-13, BB-21, BB-9, BB-29, BB-61, BD-9 e BG-1. A virulência para as cultivares melhoradas como 'Canastra', 'Confiança', 'Carisma', 'Maravilha', 'Primavera' e 'Bonança' foi freqüente na população do patógeno. Alguns patótipos brasileiros que apresentaram reações diferenciais em cultivares comerciais poderão ser utilizados para incorporação de genes de resistência nas cultivares suscetíveis melhoradas para qualidade de grãos pelo método de melhoramento convencional.

INTRODUCTION

Rice blast caused by *Pyricularia grisea* (Cooke) Saccardo = *P. oryzae* Cavara [Tel.: *Magnaporthe grisea* (Hebert) Barr.] continues to be one of the major rice (*Oryza*

sativa) diseases in different parts of the world including Brazil, in spite of the extensive research efforts on breeding varieties for blast resistance, over the years. The disease assumes greater dimension in yield losses under upland conditions in the west central and northern states of Brazil. Embrapa Rice & Beans in collaboration with other state research organizations has developed several upland rice cultivars with superior grain

*Bolsista do CNPq

quality and different degrees of blast resistance for cultivation. Blast epidemics have been reported to be of common occurrence in the newly released cultivars such as 'Primavera', 'Maravilha' and 'Confiança' during the past three years. The limited durability of blast resistance of the improved rice cultivars could be attributed to the highly complex and variable nature of the pathogen. The blast fungus has been known to be composed of races or pathotypes (Latterell *et al.*, 1954).

Latterell *et al.* (1954) first reported the occurrence of pathogenic races in *P. grisea* in the USA. Since then several races have been reported in different countries using different rice cultivars as differentials. The number of races has increased as more isolates were tested (Ou, 1980). An international set of eight cultivars was selected for identifying races of *P. grisea* on the basis of reaction on these differentials. These races were called international races and were designated as IA, to IH, followed by numerals to indicate race numbers (Atkins *et al.*, 1967; Ling & Ou, 1969). International races representing all nine groups of races were recovered from 15 rice cultivars in Colombia (Correa-Victoria & Zeigler, 1993; Levy *et al.*, 1993). In Brazil, the occurrence of physiologic races has been reported by several investigators (Amaral *et al.*, 1979; Bedendo *et al.*, 1979; Ribeiro, 1980; Ribeiro & Terres, 1987; Prabhu & Filippi, 1989; Urashima & Isogawa, 1990; Malavolta *et al.*, 1992; Filippi *et al.*, 1999).

In Japan, a set of nine differentials, each with a known resistance gene, was proposed (Yamada *et al.*, 1976). Since these differentials are not isogenic lines, other unknown resistance genes may be present in cultivars that caused a failure to describe adequately the pathotype diversity in the tropics (Ou, 1980). Recently, Mackill & Bonman (1992) developed an improved differential set of five near isogenic lines (NIL's) of cultivar CO39, with single major blast resistance genes, for characterizing pathogen population. However, their value became limited because only a few known genes are involved to describe the entire virulence spectrum of pathotypes.

In a study utilizing International, Japanese and Korean differential sets, Bonman *et al.* (1986) have shown that none of the agriculturally important Philippine isolates of *P. grisea* could be differentiated. The conventional differential sets and the expression of results as race numbers did not yield useful information. According to these authors, for practical purpose in studies on the diversity of agriculturally important isolates of *P. grisea*, the use of widely grown local commercial cultivars a differential set is preferable to the conventional set of eight international differentials. Detailed investigations on pathogenic variability of *P. grisea* at the Santa Rosa breeding site in Colombia have also shown that several races could be further differentiated into different pathotypes when local commercial cultivars were used as differentials (Correa-Victoria & Zeigler, 1993).

The knowledge on the pathotypes of *P. grisea* that affect the rice cultivars in the experimental plots under field conditions in Brazil is important for resistance breeding. The present paper reports the pathotypic diversity among the

isolates retrieved from improved upland rice cultivars under natural conditions of infection in field plots, the cultivar differences in relation to the pathotypes that affect them and virulence characterization of Brazilian pathogen population of *P. grisea* using an additional set of eight local upland rice cultivars as differentials.

MATERIALS AND METHODS

Single conidial isolates of *P. grisea* were obtained from sporulating lesions on leaves of 13 improved upland rice cultivars ('Primavera', 'Maravilha', 'Canastra', 'Caiapó', 'Carajás', 'Progresso', 'Rio Paranaíba', 'Araguaia', 'Guarani', 'IAC-201', 'Bonança' and 'Carisma'). Collections were made in experimental plots located in the Embrapa Rice and Beans, Goiânia, during five consecutive years (1996-2000).

Five hundred and twenty isolates were established and from them, 85 were selected for evaluation according to the year and cultivar from which they were retrieved. The virulence frequency of the selected isolates was tested under controlled greenhouse conditions, utilizing 32 genotypes including 14 upland rice cultivars ('Araguaia', 'Bonança', 'Caiapó', 'Canastra', 'Carajás', 'Carisma', 'Confiança', 'Guarani', 'IAC-47', 'IAC-201', 'Maravilha', 'Primavera', 'Progresso' and 'Rio Paranaíba'), a check for partial resistance evaluation ('Moroberekan'), a somaclone of cultivar 'Araguaia' (SC09) with a known gene for vertical resistance, besides eight standard international differentials ('Dular', 'Kanto 51', 'NP125', 'Raminad Str 3', 'Usen', 'Zenith', 'Caloro' and 'Sha-tiao-tsao'), five near isogenic lines of 'CO 39' ('C 101 LAC', 'C101A 51', 'C104 PKT', 'C101 PKT' and 'C101-TTP-4L-23'), a recently released improved upland rice cultivar ('Aimoré'), and two introduced lines of *Oryza glaberrima* Steud (OG 217 and OG 218). The test material was planted in plastic trays (15 x 30 cm) containing 3 kg of soil fertilized with NPK (5 g of 5-30-15 + Zn and 3 g of ammonium sulfate per 3 kg of soil). Thirty two cultivars, 16 per tray, were sown (ten to 12 seeds/cultivar) in 5 cm long rows.

The inoculation procedure described earlier by Filippi *et al.* (1999) was utilized. The physiologic races were identified based on the reaction of eight standard international differentials (Atkins *et al.*, 1967; Ling & Ou, 1969). Leaf blast reaction was assessed seven to nine days after inoculation taking into consideration only two types of reaction of the host, compatible or susceptible and incompatible or resistant reaction, according to Ling & Ou (1969). The infection types 0 to 3 were considered as resistant and 4 to 9 as susceptible in the disease evaluation scale of 0-9 (International Rice Research Institute, 1988). In case of ambiguous or intermediate reaction, inoculation tests were repeated whenever necessary and the ones that gave consistent and uniform reaction were utilized for analysis. A tray containing international and Brazilian differential cultivars as a non-inoculated control was maintained to ensure that no contamination occurred during the inoculation procedure

RESULTS AND DISCUSSION

The Brazilian pathotypes were differentiated on the basis of reaction in eight commercial upland rice cultivars. They were selected taking into consideration the virulence frequency of the isolates, genetic base, and agricultural importance. Care was taken to avoid cultivars with 'IAC-47' as one of the parents such as 'Rio Paranaíba', 'Guarani', 'Cuiabana', etc. The selected eight cultivars, henceforth, can be used as standard set of differentials for identifying Brazilian pathotypes of *P. grisea* collected from upland rice cultivars. The international differentials were substituted (Table 1).

The Brazilian pathotypes were prefixed by the letter "B" instead of "I". The groups were designated as BA, BB, BC, BD, BE, BF, BG, BH and BI. The isolates of *P. grisea* virulent to 'Carajás' belong to BA group (128 reaction patterns) similar to the international race group IA. This group separation was based on susceptible reaction of the international rice differential 'Raminad Str.3' (Atkins *et al.*, 1967). The isolates, which induced susceptible reaction on the second differential 'Zenith' and resistant reaction on 'Raminad Str' were grouped under IB group. In a similar manner the isolates showing susceptible reaction on second Brazilian differential 'Primavera' pertain to BB group (64 reaction patterns). The isolates with susceptible reaction on 'Maravilha' but resistant reaction on 'Carajás' and 'Primavera' were designated as BC group (32 patterns). The isolates causing susceptible reaction on 'Confiança', 'Progresso', 'Caiapó', 'IAC-47' and 'IAC-201' but resistant reaction to the first three differentials taken in a sequential order, were designated as BD, BE, BF, BG, BH groups with 16, 8, 4, 2, and 1 reaction patterns, respectively. The race group BI was proposed for the isolates which show resistant reaction on all eight differentials as in the case of international race group II. The numeral following the group letters indicates the pathotype number similar to the numbering of international races and the same key with reaction patterns of international differentials (Ling & Ou, 1969) was used for designating Brazilian pathotypes.

Based on the reaction pattern of eight international differentials, 11 pathotypes (IB-1, IB-9, IB-41, IB-45, IB-33, IB-37, IC-1, IC-9, IC-25, ID-9, and IG-1) were identified in a sample population of 85 *P. grisea* isolates collected from 14 upland rice cultivars (Table 2). The predominant international races or pathotypes were IB-9 (56.4%), IB-1 (16.4%) and IB-41 (11.8%). The pathotype IB-9 was recovered from all 14 rice cultivars. All seven isolates of *P. grisea* collected from the cultivar 'Rio Paranaíba' during a four-year period from 1997 to 2000 was classified as the pathotype IB-9. The pathotypes IB-1, IB-41 and IB-45 were recovered from eight, seven and three of the 14 rice cultivars, respectively. The predominance of the pathotypes IB-9, IB-1 and IB-41 was also reported in tests conducted during 1986-88, among isolates retrieved from ten upland rice cultivars (Prabhu & Filippi, 1989). These results showed that there was no change in relation to the frequency of the predominant pathotypes despite the change in varietal pattern. The three pathotypes representing group IC, two of IC-9, and one each of IC-1 and IC-25 obtained from 'Primavera' were in low frequency.

Twenty-six Brazilian pathotypes were identified utilizing eight commercial upland rice cultivars in contrast to 11 international pathotypes (Table 2). They represented race groups BB, BA, BC, BD, BG, BF and BI. The most predominant Brazilian pathotypes BB-21 and BB-41 were represented by 28.2% and 17.6%, respectively of the 85 test isolates. A relatively high frequency of pathotype BB-21 was obtained from 'IAC-47', 'Canastra' and 'Bonança'. All isolates recovered from 'Bonança' were classified as BB-21 whereas with international differentials two pathotypes IB-9 and IB-33 were identified. Similarly, all isolates collected from 'Rio Paranaíba' were classified as the pathotype IB-9 but with the Brazilian differentials it was subdivided into four different pathotypes such as BB-41, BB-49, BC-1 and BD-9. Twenty-four of the 28 pathotypes were not very frequent and were recovered at random from different cultivars. The utility of the pathotypes of *P. grisea* is enhanced if the designation indicates both international and Brazilian groups followed by its respective number. For example, pathotype of

TABLE 1 - Pathotype group of *Pyricularia grisea*, standard international differentials, Brazilian differentials and their parents

Pathotype group ¹	International differentials	Brazilian differentials	Parents of Brazilian differentials
A	Raminad Str 3	Carajás	IREM 293-B /IAC 81-176
B	Zenith	Confiança	IAC 164/IRAT 216
C	NP-125	Maravilha	TOX1010-49-1/IRAT121// (COL1xM312A
D	Usen	Primavera	IRAT10/LS85-158
E	Dular	Progresso	(COL.1x M312A)IRAT124//RHS107-2-1-2TB-1JM
F	Kanto	Caiapó	IRAT13/B.CAMPO//CNA _x 104/PEROLA
G	Sha-tiao-tsaio	IAC-47	IAC 1246/IAC 1391
H	Caloro	IAC-201	IAC 165/LABELLE

¹The isolates which produce resistant reaction on all eight differentials pertain to the pathotype group I.

TABLE 2 - Pathotypes of *Pyricularia grisea* determined based on standard international differentials, additional set of Brazilian upland rice (*Oryza sativa*) cultivars and virulence spectrum

Cultivar ¹	Isolate ²	IP ³	BP ⁴	CV ⁵	Cultivar	Isolate	IP	BP	CV	
Carisma	Py2294-00	IB-41	BB-21	0.65	Progresso	Py2396-00	IB-33	BB-41	0.75	
	Py2288-00	IB-1	BB-29	0.65		Py2390-00	IB-41	BB-21	0.65	
	Py1983-99	IB-9	BB-29	0.71		Py1579-98	IB-1	BB-41	0.62	
Maravilha	Py1981-99	IB45	BA-21	0.56		Py1577-98	IB-1	BB-61	0.62	
	Py2436-00	IB-41	BB-21	0.68		Py1365-97	IB-9	BB-29	0.62	
	Py2433-00	IB-1	BB-21	0.75		Py964-96	IB-9	BB-61	0.56	
	Py2004-99	IB-9	BB-21	0.68		Py962-96	IB-1	BB-21	0.68	
	Py2001-99	IB-9	BB-45	0.62	IAC-47	Py2254-00	IB-41	BB-21	0.71	
	Py1598-98	IB-41	BB-21	0.62		Py2253-00	IB-9	BB-21	0.68	
	Py1387-97	IB-9	BB-29	0.59		Py1980-99	IB-9	BB-41	0.62	
IAC-201	Py1389-97	IB-9	BB-49	0.68		Py1593-98	IB-41	BB-21	0.65	
	Py2248-00	ID-9	BD-16	0.31		Py1347-97	IB-9	BB-21	0.71	
	Py2249-00	IB-9	BB-41	0.71		Py723-96	IB-9	BA-21	0.71	
	Py1355-97	IB-1	BB-41	0.68	Confiança	Py2362-00	IG-1	BI-1	0.12	
	Py948-96	IB-9	BB-57	0.59		Py2356-00	IB-9	BB-41	0.62	
Rio Paranaíba	Py940-96	IB-9	BB-61	0.59		Py1322-97	IB-45	BB-21	0.65	
	Py733-96	IB-9	BB-25	0.75		Py1317-97	IB-9	BB-13	0.53	
	Py2407-00	IB-9	BB-41	0.56	Carajás	Py1311-97	IB-41	BB-41	0.53	
	Py2403-00	IB-9	BD-9	0.59		Py2309-00	IB-9	BA-29	0.59	
	Py2012-99	IB-9	BC-1	0.59		Py2307-00	IB-9	BA-29	0.65	
	Caiapó	Py2010-99	IB-9	BB-41	0.62		Py2000-99	IB-1	BB-13	0.62
		Py1589-98	IB-9	BC-49	0.65		Py1997-97	IB-37	BA-125	0.62
		Py1276-97	IB-9	BB-41	0.62		Py1421-97	IB-45	BA-61	0.50
		Py1271-97	IB-9	BB-41	0.59	Guarani	Py1415-97	IB-45	BA-53	0.65
		Py2347-00	IB-9	BB-33	0.71		Py2425-00	IB-41	BB-21	0.68
Py2345-00		IB-1	BB-9	0.81	Py2422-00		IB-1	BB-29	0.53	
Primavera		Py1995-99	IB-9	BB-41	0.62		Py1971-99	IB-1	BD-9	0.50
		Py1991-99	IB-1	BB-41	0.68		Py1967-99	IB-1	BB-61	0.68
		Py1596-98	IB-9	BD-9	0.62		Py1586-98	IB-1	BB-41	0.65
		Py1594-97	IB-9	BB-1	0.75		Py1581-98	IB-9	BD-9	0.68
		Py1402-97	IB-9	BG-1	0.59	Araguaia	Py1393-97	IB-9	BA-9	0.71
		Py2376-00	IC-25	BD-16	0.25		Py2265-00	IB-1	BB-13	0.78
	Py2008-99	IC-9	BD-16	0.21	Py2267-00		IB-9	BA-29	0.71	
	Canastra	Py2007-99	IC-1	BD-16	0.37		Py1359-97	IB-9	BG-1	0.56
		Py1303-97	IC-9	BD-16	0.37		Py1363-97	IB-9	BB-45	0.59
		Py1300-97	IB-9	BB-41	0.59		Py646-96	IB-9	BF-1	0.56
Py2323-00		IB-9	BB-21	0.71	Bonança	Py651-96	IB-9	BG-1	0.53	
Py2329-00		IB-41	BB-21	0.68		Py2453-00	IB-9	BB-21	0.71	
Py1989-99		IB-9	BB-21	0.71		Py2449-00	IB-33	BB-21	0.75	
Py1988-99		IB-9	BB-21	0.68		Py1975-99	IB-9	BB-21	0.71	
		Py1375-97	IB-9	BC-29	0.62		Py1977-99	IB-9	BB-21	0.75
	Py1379-97	IB-9	BB-21	0.71		Py1978-99	IB-9	BB-21	0.62	
	Py1382-97	IB-41	BB-21	0.65						

¹Cultivar from which the isolate was collected; ²Accession number of *P. grisea* culture collection of Embrapa Rice & Beans, last two digits indicate the year of collection; ³International pathotype; ⁴Brazilian pathotype; ⁵Coefficient of virulence was calculated as the proportion of the total number of compatible (susceptible) reactions in relation to 32 genotypes.

the isolate collected from 'Primavera' has to be designated as IB-9/BD-17. These results are in accordance with those obtained in other countries, where the standard international set does not permit identification of all pathotype diversity that occurs in several rice growing regions (Ou, 1980; Bonman *et al.*, 1986; Correa-Victoria & Zeigler, 1993).

Distinct differences in virulence pattern of the isolates on 32 genotypes were observed (Table 2). The coefficient of virulence ranged from 0.12 for the pathotype IG-1/BI-1 from 'Confiança' to 0.81 for the pathotype IB-1/BB-9 from 'Caiapó'. The cultivar 'Primavera' showed low coefficients

of virulence varying from 0.21 to 0.59.

The virulence pattern of 14 different isolates of pathotype IB-1 on Brazilian rice differentials is presented in Table 3. The eight international differentials showed one reaction pattern i.e. the isolates induced resistant reaction on 'Raminad Str 3' and susceptible reaction on all other differentials and hence was classified as pathotype IB-1. When the Brazilian commercial cultivars were utilized as differentials they could be further divided into eight pathotypes: BB-41, BB-13, BB-21, BB-9, BB-29, BB-61, BD-9 and BG-1 (Table 3). Even though the four isolates of

TABLE 3 - Virulence pattern of isolates classified as pathotype IB-1 of *Pyricularia grisea* on Brazilian rice (*Oryza sativa*) differentials

Isolate ¹	A Carajás	B Confiança	C Maravilha	D Primavera	E Progresso	F Caiapó	G IAC-47	H IAC-201	Pathotype ³
Py2433-00	R ²	S ²	S	R	S	R	S	S	BB-21
Py2345-00	R	S	S	S	R	S	S	S	BB-9
Py1991-99	R	S	R	S	R	S	S	S	BB-41
Py2265-00	R	S	S	S	R	R	S	S	BB-13
Py1967-99	R	S	R	R	R	R	S	S	BB-61
Py2288-00	R	S	S	R	R	R	S	S	BB-29
Py1355-97	R	S	R	S	R	S	S	S	BB-41
Py1579-98	R	S	R	S	R	S	S	S	BB-41
Py1577-98	R	S	R	R	R	R	S	S	BB-61
Py962-96	R	S	S	R	R	R	S	S	BB-29
Py1971-99	R	R	R	S	R	S	S	S	BD-9
Py1359-97	R	R	R	R	R	R	S	S	BG-1
Py2000-99	R	S	S	S	R	R	S	S	BB-13
Py1586-98	R	S	R	S	R	S	S	S	BB-41

¹Accession number of *P. grisea* culture collection of Embrapa Rice & Beans, last two digits indicate the year of collection; ²R= resistant reaction; S= susceptible reaction; ³Pathotypes were identified using eight commercial upland rice cultivars as additional differentials

pathotype BB-41 were recovered from four different cultivars, (Py 1991-99 from 'Caiapó'; Py 1355-97 from 'IAC-201'; Py 1579-98 from 'Progresso'; Py 1586-98 from 'Guarani') in different years, they showed similar virulence pattern. They exhibited resistant reaction on 'Carajás', 'Maravilha' and 'Progresso' and susceptible reaction on the other five differentials and thus classified as BB-41.

Both virulent and avirulent isolates were found in the population of *P. grisea* to the known genes in NIL's. Of 85 isolates, 82 and 79 were virulent for genes *Pi-3* and *Pi-4a*, respectively. These results suggest that the virulence on *Pi-3* and *Pi-4* pre-existed abundantly in populations of *P. grisea* even though these genes have never been utilized in upland rice breeding. The virulence frequencies were relatively lower in descending order on *Pi-4b*, *Pi-2*, and *Pi-1*. The host from which they were collected seems to condition the virulence pattern. For example, six of the seven isolates from 'Maravilha' were virulent to *Pi-2* and avirulent to *Pi-1*. On the other hand, four of the six isolates originated from 'IAC-201' were virulent to *Pi-1* and avirulent to *Pi-2*.

Cross inoculations of *P. grisea* isolates on newly released cultivars 'Primavera', 'Maravilha', 'Bonança' and 'Carisma', from which they were collected, exhibited differential interaction between isolates and cultivars (Table 4). The five isolates, which were virulent to 'Primavera', were avirulent to 'Maravilha' whereas of the seven isolates of 'Maravilha' six were avirulent to 'Primavera'. The pathotype IB9/BB45 from 'Maravilha' was not virulent to 'Maravilha', but virulent to 'Primavera' (Table 4). Studies conducted in Philippines and Colombia with widely prevalent isolates of *P. grisea* also showed that some isolates were not found to be virulent to the cultivar of their origin. This was attributed to variation in virulence pattern of conidia produced in culture

TABLE 4 - Pathotype x cultivar interaction of *Pyricularia grisea* in relation to leaf blast reaction in inoculations tests

Pathotype ¹	Cultivar			
	Primavera	Maravilha	Bonança	Carisma
Primavera				
IC-25/BD-16	9 ²	0	0	0
IC-9/BD-16	7	0	0	0
IC-1/BD-16	9	0	0	0
IC-9/BD-16	4	0	0	0
IB-9/BB-41	5	0	0	0
Maravilha				
IB-41/BB-21	0	7	7	7
IB-1/BB-21	0	5	7	7
IB-9/BB-21	3	7	7	7
IB-9/BB-45	4	3	0	4
IB-41/BB-21	0	7	7	7
IB-9/BB-29	0	7	0	7
IB-9/BB-49	0	5	4	7
Bonança				
IB-9/BB-21	0	9	7	9
IB-33/BB-21	0	7	7	7
IB-9/BB-21	0	7	5	4
IB-9/BB-21	0	7	5	7
IB-9/BB-21	0	4	5	7
Carisma				
IB-41/BB-21	0	5	5	5
IB-1/BB-29	0	5	0	7
IB-9/BB-29	0	5	5	7
IB-45/BA-21	0	7	0	7

¹International and Brazilian pathotypes of *P. grisea* were identified from isolates collected from commercial rice cultivars indicated in the column.

²Disease scores 0 to 3 and 4 to 9 correspond to resistant and susceptible reactions, respectively.

media and the resistance spectrum of the cultivar (Ou, 1980; Correa-Victoria & Zeigler, 1993). The isolates from 'Primavera' were virulent to 'Primavera' and avirulent to 'Bonaça' and *vice versa*. The isolates of 'Carisma' showed differences in their reaction on 'Bonaça'. This indicates that the international and the Brazilian pathotypes may not give required information in relation to cultivar specific isolates. Pathotype may not necessarily be the appropriate phenotypic unit for breeding purpose because 71 distinct virulence patterns were observed on the 21 cultivars tested (Zeigler *et al.*, 1995). The isolate cultivar interaction could be more useful for incorporating resistance genes against a specific isolate pertaining to a given pathotype. The advantages of pathotype analysis are to broadly group the pathogen population into a reduced number of units. For example, the 85 isolates in this study could be grouped into 11 international or 21 Brazilian pathotypes.

Individual differences in phenotypic virulence are common and the virulence pattern of the *P. grisea* in test plots must be monitored constantly. Virulence in the field also preexists for genes that have never been utilized but in low frequencies. The sample of isolates collected over a period of five years in this study showed variation between individuals with distinct differences in reaction on commercial rice cultivars. Some of the Brazilian pathotypes could be utilized for incorporating resistance genes in susceptible cultivars improved for grain quality by conventional breeding methods.

LITERATURE CITED

- AMARAL, R.E.M., SOUZA, D.M., MALAVOLTA, U.M.A. & ISSA, E. Raças fisiológicas de *Pyricularia grisea* Cav. no Estado de São Paulo. *O Biológico* 45:205-208. 1979.
- ATKINS, J.G., ROBERT, A.L., ADAAIR, C.R., GOTO, K., KOZAKA, T., YANAGIDA, R., YAMADA, M. & MATSUMOTO, S. An international set of rice varieties for differentiating races of *Pyricularia oryzae*. *Phytopathology* 57:297-301. 1967.
- BEDENDO, I.P., RIBEIRO, A.S. & CARDOSO, C.N. Variabilidade do fungo de *Pyricularia oryzae*, agente causal da brusone no arroz. *Summa Phytopathologica* 5:106-109. 1979.
- BONMAN, J.M., DE DIOS, V.T.I., & KHIN, M.M. Physiologic specialization of *Pyricularia oryzae* in the Philippines. *Plant Disease* 70:767-769. 1986.
- CORREA-VICTORIA, F.J. & ZEIGLER, R.S. Pathogenic variability in *Pyricularia grisea* at a rice blast "hot-spot" breeding site in Eastern Colombia. *Plant Disease* 77:1029-1035. 1993.
- FILIPPI, M.C., PRABHU, A.S. & LEVY, E.M. Differential compatibility of *Pyricularia grisea* isolates with some Brazilian irrigated rice cultivars. *Fitopatologia Brasileira* 24:447-450. 1999.
- INTERNATIONAL RICE RESEARCH INSTITUTE. Standard evaluation system for rice. International rice testing program. 3rd ed. Los Banos. International Rice Research Institute. 1988.
- LATTERELL, F.M., TULLIS, E.C., OTTEN, R.J. & GUBERNICK, A. Physiologic races of *Pyricularia oryzae*. *Phytopathology* 44:495-496. 1954 (Abstract).
- LEVY, M., CORREA, F.J., ZEIGLER, R.S., XU, S. & HAMER, J.E. Genetic diversity of the rice blast fungus in a disease nursery in Colombia. *Phytopathology* 83:1427-1433. 1993.
- LING, K.C. & OU, S.H. Standardization of the international race numbers of *Pyricularia oryzae*. *Phytopathology* 59:339-342. 1969.
- MACKILL, D.J. & BONMAN, J.M. Inheritance of blast resistance in near-isogenic lines of rice. *Phytopathology* 83:746-749. 1992.
- MALAVOLTA, V.M.A. & SOUZA, T.M.W. Variabilidade de *Pyricularia oryzae* no Estado de São Paulo. *Summa Phytopathologica* 18:287-290. 1992.
- OU, S.H. Pathogen variability and host resistance of the rice blast fungus, *Pyricularia oryzae* Cav. *Annual Review of Phytopathology* 18:167-187. 1980.
- PRABHU, A.S. & FILIPPI, M.C. As raças fisiológicas de *Pyricularia oryzae* virulentas nas cultivares melhoradas de arroz de sequeiro. *Fitopatologia Brasileira* 14:140. 1989. (Resumo).
- RIBEIRO, A.S. Prevalência de raças de *Pyricularia oryzae* Cav. No Rio Grande de Sul. *Pesquisa Agropecuária Brasileira* 15:175-182. 1980.
- RIBEIRO, A.S. & TERRES, A.N.L.S. Variabilidade do fungo *Pyricularia oryzae* e sua relação com cultivares resistentes à brusone. *Fitopatologia Brasileira* 12:316-321. 1987.
- URASHIMA, A.S. & ISOGAWA, Y. Identification of races of *Pyricularia oryzae* causing blast disease in variety IAC 4440 in Paraíba Valley-SP. *Summa Phytopathologica* 16:243-247. 1990.
- YAMADA, M., KIYOSAWA, S., YAMAGUCHI, T., HIRANO, T., KOBAYASHI, K. & WATANABI, S. Proposal for a new method for differentiating races of *Pyricularia oryzae* Cavra in Japan. *Annual. Phytopathological Society of Japan*. 42:216-219. 1976.
- ZEIGLER, R.S., CUOC, L.X., SCOTT, R.P., BERNARDO, M.A., CHEN, D.H., VALENT, B. & NELSON, R.J. The relationship between lineage and virulence in *Pyricularia grisea* in the Philippines. *Phytopathology* 85:443-451. 1995.