



## Underrated *Staphylococcus* species and their role in antimicrobial resistance spreading

Ciro César Rossi<sup>1</sup> , Monalessa Fábila Pereira<sup>2</sup>  and Marcia Giambiagi-deMarval<sup>1</sup> 

<sup>1</sup>Universidade Federal do Rio de Janeiro, Instituto de Microbiologia Paulo de Góes, Laboratório de Microbiologia Molecular, Rio de Janeiro, RJ, Brazil.

<sup>2</sup>Universidade Federal do Espírito Santo, Departamento de Patologia, Vitória, ES, Brazil.

### Abstract

The increasing threat of antimicrobial resistance has shed light on the interconnection between humans, animals, the environment, and their roles in the exchange and spreading of resistance genes. In this review, we present evidences that show that *Staphylococcus* species, usually referred to as harmless or opportunistic pathogens, represent a threat to human and animal health for acting as reservoirs of antimicrobial resistance genes. The capacity of genetic exchange between isolates of different sources and species of the *Staphylococcus* genus is discussed with emphasis on mobile genetic elements, the contribution of biofilm formation, and evidences obtained either experimentally or through genome analyses. We also discuss the involvement of CRISPR-Cas systems in the limitation of horizontal gene transfer and its suitability as a molecular clock to describe the history of genetic exchange between staphylococci.

**Keywords:** Horizontal gene transfer, antimicrobial resistance, coagulase negative staphylococcus, biofilm, CRISPR, One Health.

Received: February 26, 2019; Accepted: May 06, 2019.

### Introduction

Every year, hundreds of thousands of deaths around the world are attributed to the ever increasing problem of antimicrobial resistance (Laxminarayan *et al.*, 2016). It is estimated that, if the issue is not properly addressed, by the year of 2050, more than 10 million annual deaths will be caused by antimicrobial-resistant microorganisms, surpassing deaths by cancer (O'Neill, 2014). Although the accuracy of this frightening estimate is questioned by some authors, since the future scenario of disease treatment may be considerably different from the current one, the clinical, economical, and public health burden associated with antimicrobial resistance is undeniable (Kraker *et al.*, 2016; Robinson *et al.*, 2016).

In the context of resistance dissemination, bacteria of the *Staphylococcus* genus, residents of the normal microbiota of human beings and animals, play a central role. *Staphylococcus aureus* is the main pathogen of the group, responsible for a variety of clinical infections in humans and a leading cause of bacteremia, endocarditis, and many infections related to invasive medical devices (Tong *et al.*, 2015). Meanwhile, coagulase negative staphylococci (CoNS), especially *S. epidermidis* and *S. haemolyticus*,

have emerged as recurrent causative agents of nosocomial infections, mainly those related to indwelling devices (Becker *et al.*, 2014). They are a serious threat in the twilight of the multidrug resistance era, for actively participating in the horizontal transmission of resistance (Becker *et al.*, 2014; Czekaj *et al.*, 2015).

Genomic analyses indicate that many of the genetic determinants of resistance may have been exchanged between several staphylococcal species from different environments and hosts (Rolo *et al.*, 2017; Rossi *et al.*, 2017a; Kohler *et al.*, 2018;). This includes species that are understudied and/or underestimated, either for having been recently discovered, rarely involved with infectious diseases, or for lacking canonical staphylococcal virulence factors. However, recent evidence shows that these underrated species, despite not being usual pathogens, may have an important role in the exchange of antimicrobial resistance genes, by acting as gene-reservoirs for more pathogenic species, such as *S. aureus* (Otto, 2013; Rossi *et al.*, 2017a, 2019).

Given the increasing recognition of the importance of previously overlooked *Staphylococcus* species, the goal of this review was to present evidences that put these bacteria in the front row of resistance dissemination and highlight their potential threat to human and animal health.

Send correspondence to Marcia Giambiagi-deMarval. Instituto de Microbiologia Paulo de Góes, Departamento de Microbiologia Médica, CCS-Centro de Ciências da Saúde-Bloco I-sala I2-07. Avenida Carlos Chagas Filho, 373-Cidade Universitária, 21941-902, Rio de Janeiro, RJ, Brazil. E-mail: [marciagm@micro.ufrj.br](mailto:marciagm@micro.ufrj.br)

## Antimicrobial resistance increase and the interconnectedness of its spreading

In the past decade, consumption of antibiotic drugs increased by 35%, with 76% of this growth concentrated in Brazil, Russia, India, China, and South Africa (Van Boeckel *et al.*, 2014), with a projection of a rise in consumption in the next 15 years of 67% (Van Boeckel *et al.*, 2015). The disturbing situation of antimicrobial resistance led the World Health Organization to elaborate the “Global action plan on antimicrobial resistance”, with goals that include: (i) improving the understanding of the problem through communication, education and training, (ii) increasing surveillance and research, (iii) advancing in preventive actions to reduce the incidence of infections, and (iv) optimizing the use of antimicrobial drugs in human and veterinary health (WHO, 2017). Tied to this worldwide concern to restrain the dispersion of resistance, the emerging engagement of scientists and other professionals with the “One Health” agenda increases the acknowledgement of the need for global approaches as the only possible way to keep the predicted disaster of 2050 of more than 10 million annual deaths caused by resistant microorganisms from actually happening.

Considered as a “new professional imperative”, One Health is a collaborative and multidisciplinary effort to achieve optimal health for people, animals and our environment from local to global scales (King *et al.*, 2008). It recognizes that the welfare of these three domains is interconnected and this link, ignored for so long, is crucial for the spreading of antimicrobial resistance. The use of antimicrobial drugs in agriculture, for example, is the largest worldwide, thus being a major driver of resistance for several reasons, like exposure of bacteria to sub-therapeutic doses of the antibiotics and the exposure of human and animals to those drugs and microorganisms, either via consumption of products or environmental release (Silbergeld *et al.*, 2008). Studies also point out that the use of antimicrobial drugs, particularly to maintain health, productivity, and promoting growth of food animals, contribute to the dispersion of resistant bacteria in livestock and human beings (Marshall and Levy, 2011; Van Boeckel *et al.*, 2015). In aquaculture, it is estimated that around 80% of antimicrobials used reach the environment with their activity intact, thereby expanding the surroundings where selection of resistant microorganisms will take place (Cabello *et al.*, 2013). Even insects commonly associated with food animals, like houseflies and cockroaches, are presumably vehicles of microorganisms from the farms to urban centers (and vice versa), as evidenced by multidrug-resistant clonal lineages carried by them, that were also isolated from different environments (Zurek and Ghosh, 2014).

Pet animals have been shown to act as reservoirs of resistant bacteria, which in turn act as reservoirs of mobile genetic elements that carry antimicrobial resistance genes (Guardabassi *et al.*, 2004; Rossi *et al.*, 2017a). In fact, the relationship between the animals and their owners signifi-

cantly shapes the microbiota of both counterparts (Song *et al.*, 2013). For that reason, the indiscriminate use of antimicrobials in veterinary practice represents a direct threat to human beings. More aggravating is the fact that some drugs that are either not recommended to be used in humans, or those that are considered as last resources, are heavily used to treat animals. Some examples include the polymixins and chloramphenicol and its derivatives, the latter presenting several adverse effects that limit its employment (Cabello *et al.*, 2013; Poirel *et al.*, 2017).

Consistently, multidrug resistant strains, like methicillin-resistant *Staphylococcus aureus* (MRSA) are ubiquitous, being isolated from humans, pets, food, other animals and the environment (Vanderhaeghen *et al.*, 2010; Kock *et al.*, 2013; Rossi *et al.*, 2017b). Due to local variations in control practices and specific characteristics of circulating clones, the overall geographic distribution of MRSA, for example, can range from 1 to 5% of isolates in northeastern Europe to more than 50% in certain Latin American countries (Brazil, Uruguay, Venezuela, Bolivia, Peru, and Chile) and in Japan (Lee *et al.*, 2018).

## Variety and clinical significance of *Staphylococcus* spp.

Although *S. aureus* is the major bacterium of its genus, more than 50 *Staphylococcus* species are registered in the *List of Prokaryotic Names with Standing in Nomenclature* database, available at <http://www.bacterio.net> (Parte, 2018). However, DNA sequencing of complete genomes or housekeeping genes, phylogenetic analyses, DNA–DNA hybridization, protein profiles, and genotyping techniques have constantly led to reclassifications or proposals of new species and subspecies (Sasaki *et al.*, 2007; Fitzgerald, 2009; Taponen *et al.*, 2012). As these techniques advance and new sources of *Staphylococcus* are explored, especially in different animals, new species are also discovered, such as *S. nepalensis*, isolated from goats (Spergser *et al.*, 2003), *S. stepanovicii*, isolated from wild small animals (Hauschild *et al.*, 2010), *S. pseudintermedius*, isolated from several domestic animals, such as dogs, cats, horses and parrots (Devriese *et al.*, 2005), and *S. agnetis*, isolated from bovines with subclinical and clinical mastitis (Taponen *et al.*, 2012).

In general, staphylococci are natural inhabitants of skin and mucous membranes of human beings and animals, while the prevalence of species widely varies according to the host. *S. felis*, for example, is typically isolated from feline, either healthy or presenting signs of lower urinary tract disease, otitis externa, and ocular disease (Rossi *et al.*, 2017b; Worthing *et al.*, 2018); *S. pseudintermedius* is prevalent in domestic dogs, healthy or related to diseases like pyoderma and otitis externa (Ruscher *et al.*, 2009; Rossi *et al.*, 2018); *S. caprae* is involved with intramammary infections in dairy goats (Moroni *et al.*, 2005), among others. Regardless of their source, infections caused by unusual human pathogens are sporadically reported (Morfin-Otero

*et al.*, 2012; Novakova *et al.*, 2006a,b), with special emphasis on those caused by *S. pseudintermedius*, with most cases indicating the contact of domestic dogs with their owners as the probable source of infection (Van Hoovels *et al.*, 2006; Riegel *et al.*, 2011; Lozano *et al.*, 2017). Given their great adaptability to unfavorable conditions (Uribe-Alvarez *et al.*, 2016; Rossi *et al.*, 2017c), staphylococci isolated from the surrounding environment are also responsible for human acquisition of relevant pathogens, including MRSA (Hardy *et al.*, 2006; Sexton *et al.*, 2006).

The production of coagulase and its plasma-clotting activity is a central diagnostic feature to distinguish staphylococcal strains in coagulase-positive staphylococci (CoPS) and coagulase-negative staphylococci (CoNS) (Becker *et al.*, 2014). In addition to being key to diagnostics and group differentiation of *Staphylococcus* in CoPS and CoNS, coagulase is a virulence factor that leads to the cleavage of soluble fibrinogen to produce a fibrin coat in the surface of the bacteria, thus protecting it from phagocytosis and other host defenses (Powers and Wardenburg, 2014). Moreover, the polymorphisms of the coagulase-coding gene, *coa*, allows it to be explored in molecular typing techniques (Salehzadeh *et al.*, 2016; Javid *et al.*, 2018). However, because some populations of CoPS may not have the *coa* gene, while some CoNS present this gene, the applications of these methods are limited (Almeida *et al.*, 2018). These coagulase-variable strains are more frequently found in some species than others, but their misdiagnosis may lead to unsuitable treatment of infections and control measures. This is especially significant when the detection of the pathogenic *S. aureus* relies on coagulase production and strains of these species do not produce coagulase, leading to isolate misidentification (Sundareshan *et al.*, 2017). Strains of the CoNS *S. chromogenes*, *S. xylosus*, *S. cohnii* and *S. agnetis* have been reported to clot plasma, leading to misidentification of the pathogens causing mastitis in dairy animals (Taponen *et al.*, 2012; Santos *et al.*, 2016; Almeida *et al.*, 2018). For that reason, researchers have relied on more accurate identification methods, including the sequencing of the 16S rRNA and the housekeeping genes *rpoB*, encoding the  $\beta$  subunit of RNA polymerase, as well as *tuf*, encoding the EF-tu elongation factor (Ghebremedhin *et al.*, 2008; Li *et al.*, 2012). Given its simplicity to perform and its cost effectiveness, matrix assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF) is emerging as a potential tool for microbial identification and diagnosis (Singhal *et al.*, 2015). MALDI-TOF identification of staphylococci shows a good correlation with sequencing results, although the lack of standards for uncommon and recently identified species is still a bottleneck (Rossi *et al.*, 2017b).

CoPS, with a special emphasis on *S. aureus*, responsible for several clinical infections, from those in skin and soft tissues to systemic disease processes like sepsis (Tong *et al.*, 2015), are considered to be more pathogenic than CoNS. A plethora of virulence factors have been described

and extensively reviewed for *S. aureus*, comprising molecules involved in tissue adhesion, immune evasion and host cell injury (Powers and Wardenburg, 2014). Proteins covalently anchored to the cell wall peptidoglycan may participate in not only biotic and abiotic surface adhesion, but also in biofilm formation and iron acquisition, among other functions (Foster *et al.*, 2014). Moreover, a wide variety of toxins can be secreted, aiming to evade the defense mechanisms of the host (Otto, 2014). Other relevant pathogenic CoPS belong in the *Staphylococcus intermedium* group (SIG). This group includes zoonotic pathogens typically associated with dog bites, i.e., *S. intermedium*, *S. pseudintermedius*, the recently described *S. cornubiensis* (Murray *et al.*, 2018), and *S. delphini*, first isolated from skin lesions of dolphins (Varaldo *et al.*, 1988).

The increasing recognition of the importance of *S. pseudintermedius* as a zoonotic pathogen has boosted investigations on virulence factors involving infections caused by this bacterium. Among them, pore-forming toxins seem to play a pivotal role in the characteristic skin infections (Abouelkhair *et al.*, 2018; Maali *et al.*, 2018). Because many of these virulence factors are encoded in mobile genetic elements (MGE) with extensive variation in gene content, different strains strongly vary in their virulence arsenal (Otto, 2014; Moon *et al.*, 2015).

The CoNS constitute the vast majority of staphylococci, comprising more than 80% of the species described to date (Becker *et al.*, 2014; Parte, 2018). Historically considered as harmless inhabitants of the human and animal microbiota, in the past two decades CoNS have emerged as the major nosocomial pathogens, mostly associated with invasive medical devices (Becker *et al.*, 2014), being particularly threatening to immunocompromised individuals (Morfin-Otero *et al.*, 2012). Among these opportunistic pathogens, *S. epidermidis* is the most frequent cause of nosocomial infections (Gomes *et al.*, 2014), followed by *S. haemolyticus* (Czekaj *et al.*, 2015). The infections caused by these species are particularly important because they are difficult to treat, since device colonization is usually related to biofilm formation, which can lead to complications, including sepsis, endocarditis, and a wide variety of local infections derived from the bloodstream spreading of bacteria (Chang *et al.*, 2018; Oliveira *et al.*, 2018).

### Commensal and opportunistic staphylococci acting as gene reservoirs

*Staphylococcus* species that are usually considered as harmless inhabitants of the microbiota of animals and humans beings, like most CoNS, lack nearly all of the virulence factors described for *S. aureus* and do not encode many known specific factors, apart from a limited amount of toxins and exoenzymes (Zhang *et al.*, 2003). Their emergent threat comes from the fact that these bacteria may carry a huge amount of antimicrobial resistance genes located in MGEs (Gomes *et al.*, 2014; Czekaj *et al.*, 2015; Hosseinkhani *et al.*, 2018).

Multidrug-resistant CoNS have been increasingly linked to infections outbreaks in healthcare units (Chang *et al.*, 2018; Li *et al.*, 2018), including strains that are not only isolated from patients, but also from healthcare workers, and the environment (Widerstrom *et al.*, 2016). Microbiome studies reveal clonal staphylococcal strains wide ability to colonize diverse hosts and surrounding environments (Song *et al.*, 2013; Lax *et al.*, 2014). Their widespreadness can lead to infections caused by multidrug resistant strains in both humans and animals by species that are atypical pathogens in one of the hosts. Dutta *et al.* (2018), for example, have recently isolated *Staphylococcus pettenkoferi* strains causing peritonitis from a domestic cat. This species was discovered in 2002 in various human patients showing multiple clinical manifestations, but had never been isolated from animals before. Likewise, as aforementioned, the typical canine staphylococcal species *S. pseudintermedius* can occasionally cause disease in human beings as well (Riegel *et al.*, 2011; Lozano *et al.*, 2017).

This long overlooked exchange of microorganisms and antimicrobial resistance between different hosts and environments is now clear, and strains isolated from humans and domestic animals carry several resistance genes in common (Schwarz *et al.*, 2018). Even though some antimicrobials have their use restricted to treat infections in animals, many multidrug resistance genes in staphylococci isolated from them confer resistance to antimicrobial agents that are highly important in human medicine (Wendlandt *et al.*, 2015). Then, even if one staphylococcal species is not a common pathogen, it can be a potential threat, because it may be capable of transferring antimicrobial resistance genes to more pathogenic species, such as *S. aureus*, thereby enhancing its capacity to resist drug therapy (Haaber *et al.*, 2017). For that reason, some CoNS have been suggested to act as antimicrobial genes reservoirs within the *Staphylococcus* genus (Cafini *et al.*, 2016; Rossi *et al.*, 2017a).

The acquisition of antimicrobial resistance genes is mainly credited to the occurrence of conjugation and bacteriophage transduction and the presence of dozens of insertion sequences in staphylococcal genomes, whose rearrangements contribute to genome plasticity and strains' phenotypic diversification (Takeuchi *et al.*, 2005; Haaber *et al.*, 2017). For a while, bacteriophage transduction was perceived as the primary route of horizontal gene transfer between staphylococci, while conjugation was believed to play only a minor role in the evolution of this genus, given the scarcity of mobilization and conjugation loci in staphylococcal plasmids (Ramsay *et al.*, 2016; Haaber *et al.*, 2017). In fact, only 5% of sequenced staphylococcal plasmids harbor genes required for autonomous transfer by conjugation, which is contrasting with the abundant evidences for horizontal transfer of plasmids between different lineages and species of *Staphylococcus* (Ramsay *et al.*, 2016).

However, new mechanisms of transference of those types of plasmids have been discovered and probably explain the extensive plasmid transfer within the genus. These mechanisms include conjugation mediated by integrative and conjugative elements (ICEs), also referred to as conjugative transposons (Lee *et al.*, 2012), *in trans* recognition of multiple variants of the canonic origin of transfer (*oriT*) by some conjugative plasmids (O'Brien *et al.*, 2015b), and the activity of novel conjugative plasmids described in community isolates of *S. aureus* that are capable of mobilizing unrelated non-conjugative plasmids (O'Brien *et al.*, 2015a).

The Staphylococcal Cassette Chromosome *mec* (SCC*mec*), a genomic island that encodes resistance to methicillin and nearly all other beta-lactam antibiotics, is also a protagonist in the emergence of resistant strains. Analyses of numerous SCC*mec* sequences indicate that this mobile genetic element evolved by recombination and assembly events involving an ancestral SCC*mec* III cassette between strains of the *S. sciuri* group and the species *S. vitulinus* and *S. fleurettii*, which were then transferred to *S. aureus* and other species (Rolo *et al.*, 2017), with CoNS acting as their central reservoirs (Saber *et al.*, 2017).

Our group has demonstrated the transference of high molecular weight plasmids carrying the *mupA* gene for mupirocin resistance from *S. epidermidis*, *S. aureus*, and *S. haemolyticus* strains, from either human or canine origin, to another *S. aureus* strain (Bastos *et al.*, 1999; Rossi *et al.*, 2016, 2018). Mupirocin resistance spreading is alarming, since this drug, which is used as an intranasal ointment by healthcare workers, can significantly reduce the occurrence of nosocomial infections caused by MRSA (Patel *et al.*, 2009). Similarly, Cafini *et al.* (2016) demonstrated the transfer of linezolid resistance mediated by the *cfr* gene through plasmids, between *S. epidermidis*, *S. aureus*, and *Enterococcus* spp. isolates from Japan. *Enterococcus* spp. is also a protagonist of nosocomial infections and has been pointed out to participate in plasmid exchange with MRSA, through which resistance to the last-resource antibiotic vancomycin appears to be acquired (Kohler *et al.*, 2018). A study performed by Meric *et al.* (2015) with hundreds of genomes of *S. aureus* and *S. epidermidis* showed that these species share only about half of their pan genome, but there was a considerable sharing of mobile genetic elements between the two species, in particular genes associated with pathogenic islands and the SCC*mec*.

Biofilm formation, a characteristic feature of many CoNS (Barros *et al.*, 2015; Buttner *et al.*, 2015), seems to provide the ideal environment for the occurrence of horizontal gene transfer (Madsen *et al.*, 2012).

### Biofilms as the perfect place for horizontal gene transfer among staphylococci

Bacterial biofilms are dense surface-associated cellular communities embedded in a protective self-produced matrix of exopolysaccharides, whose development confers

new properties to its inhabitants (Black and Costerton, 2010, Flemming *et al.*, 2016). Biofilms feature a social cooperation between bacteria that enhances nutrient uptake and distribution and, given its protective matrix, hinders the exposure to antimicrobials and host defenses, thus increasing survival (Flemming *et al.*, 2016). Hence, biofilm formation plays a fundamental role in virulence.

Although the biofilm mechanisms involved in resistance to host defenses are not entirely understood, they include spatially limiting the access of leukocytes and their products to the target cells, suppression of leukocyte effector functions and cell-cell communication to increase resistance (Leid, 2009). Reducing membrane permeability also contributes to limit the entrance of antimicrobials (Liu *et al.*, 2000). However, the presence of antibiotics that affect the *Staphylococcus* cell wall have been shown to modulate natural transformation in a process that seems to be dependent of the alternative sigma factor H, SigH (Thi *et al.*, 2016). In fact, the expression of SigH-controlled genes makes *S. aureus* cells competent for transformation by plasmids or chromosomal DNA (Morikawa *et al.*, 2012), which in turn increases the probability of plasmid exchange between the cells within the biofilm. Liu *et al.* (2017) demonstrated that the presence of many antibiotics induces the expression of the *ccrCI* gene, involved with the excision of the *SCCmec* from the bacterial chromosome, thus triggering its transfer.

Overall, the high cellular density of biofilms, increased concentration of exogenous DNA, enhanced cell competence, stabilization of cell-cell contact by the matrix and even the biofilm architecture, that facilitates the dispersion of MGEs, may contribute to horizontal gene transfer (Madsen *et al.*, 2012; Savage *et al.*, 2013). The fact that more transconjugants are produced after filter-matings, when compared with planktonic cells mating, indicates the importance of biofilms for the transfer of MGEs (Madsen *et al.*, 2012). It has been shown that *S. aureus* biofilms can drastically increase the rates of conjugation and transformation of plasmids containing resistance to multiple drugs (Savage *et al.*, 2013).

As observed for many bacteria, during biofilm maturation *S. aureus* can suffer lysis and release its genomic DNA, producing the so-called external DNA (eDNA), which is a major component of the matrix of biofilms established by this bacterium (Sugimoto *et al.*, 2018). The eDNA adsorbs to the surface of a single cell in long loop structures, which act as an adhesive substance that facilitates cell attachment, in addition to influence the hydrophobicity of the bacterial cell surface (Okshevsky and Meyer, 2015). Furthermore, in *S. epidermidis*, eDNA production reduces the depth of vancomycin penetration, as the matrix-embed and negatively charged DNA interacts with positive charges of the antimicrobials (Doroshenko *et al.*, 2014). Moreover, the eDNA released by the cells to constitute the biofilm matrix is also available for horizontal gene transfer by transformation to competent cells in the community (Hannan *et al.*, 2010; Vorkapic *et al.*, 2016).

The dispersion of MGEs and eDNA is likely facilitated by the empty spaces within biofilms, which also function as channels that allow the flowing of fluids and the consequent dispersion of nutrients and oxygen to all cells. Studies indicate that this biofilm architecture is influenced by the production of biosurfactant compounds by the adherent bacteria (Davey *et al.*, 2003). Consistent with this, we have recently shown that *S. haemolyticus* strains are capable of producing biosurfactants that affect biofilm formation on abiotic surfaces (Rossi *et al.*, 2016). Likewise, surfactant peptides produced by *S. epidermidis* control biofilm maturation and detachment from colonized catheters (Wang *et al.*, 2011).


While, as abovementioned, CoNS lack most of the *S. aureus* virulence factors, biofilm formation is a major feature of *S. epidermidis* and *S. haemolyticus*, as well as of other CoNS, such as *S. saprophyticus*, *S. hominis* and *S. cohnii*. This phenotype is positively correlated with widespread antimicrobial resistance (Allori *et al.*, 2006; Czekaj *et al.*, 2015).

### The CRISPR paradox in gene-reservoirs' staphylococci

Clustered regularly interspaced short palindromic repeats (CRISPRs), allegedly present in the genomes of close to 90% of archaea and 50% of bacteria, are a prokaryotic evolutionary defense response to the preponderance of bacteriophages in the biosphere, acting as an interference system against foreign nucleic acids (Horvath and Barrangou, 2010). The CRISPR locus is composed of direct repeats of palindromic sequences interspersed with small sequences called spacers, which are fragments of exogenous sequences (Figure 1A). A functional CRISPR system has adjacent proteins (CRISPR-associated proteins, Cas), responsible for the process of recognition of foreign nucleic acids and the interference against invasive genetic elements (Horvath and Barrangou, 2010).

CRISPR interference begins with the entrance of an exogenous DNA or RNA in the bacterial cell, which is processed into small fragments by repair proteins, like those of the RecBCD complex (Amitai and Sorek, 2016), and later recognized by a complex of universal Cas1-2 proteins, which incorporates a fragment of the invader nucleic acid in the CRISPR locus as a new spacer (Figure 1B). The CRISPR locus is expressed as a long transcript that is processed, either by Cas proteins or intrinsic RNases, into small RNAs referred to as CRISPR-RNAs (crRNAs). Each crRNA is made of a fragment of the original direct repeat and the spacer, whose complementarity to foreign mobile genetic elements is the basis of the interference process, fulfilled with the activity of another Cas protein (like Cas9), or Cas complexes (van der Oost *et al.*, 2014).

Marraffini and Sontheimer (2008) described a *S. epidermidis* CRISPR containing a spacer whose sequence was homologous to a region of the *nes* gene, coding a nickase found in virtually every staphylococcal conjugative plas-



**Figure 1** - Features of CRISPR systems. (A) Main structure of a CRISPR system. For simplicity, a type II system, containing only the Cas9 protein as the interference effector, is displayed. (B) Steps of a CRISPR system activity, from foreign nucleic acid recognition and spacer incorporation (a step called adaptation) to interference.

mid sequenced. Consistently, this CRISPR prevented conjugation and transformation from happening, while the removal of the spacer allowed plasmid uptake. Later, Hatoum-Aslan *et al.* (2014) constructed mutants for all the nine *cas/csm* genes of the type III-A system in *Staphylococcus epidermidis* RP62a and showed that many mutations affected interference by impacting the interference complex formation or the crRNA biogenesis. At least 4% of the spacers within staphylococcal CRISPRs are identical to publicly available plasmid sequences, thus demonstrating their antiplasmid activity (Rossi *et al.*, 2017a). This percentage may be underrated due to under-sampling of mobile genetic elements and, as a consequence, their relative scarcity in public databases (Mojica *et al.*, 2005).

Since CRISPR-Cas systems were believed to exist in roughly 50% of bacteria, and given its antiplasmid activity, it is supposed to highly impact horizontal gene transfer among staphylococci. However, a study performed by Cao *et al.* (2016) with hundreds of *S. aureus* clinical isolates from China revealed that less than 1% of them harbored a complete arrangement of *cas* genes. Later, our group analyzed the genomes of dozens of CoNS of 15 species and also found that less than 15% of them carried CRISPRs and *cas* genes (Rossi *et al.*, 2017a). Regardless of how abundant CRISPRs really are among *Staphylococcus* isolates, they are clearly rarer than previously thought, which is consistent with the role of staphylococcal as reservoirs of antimicrobial resistance genes.

Because spacers are incorporated in the CRISPR locus following an organized and thus chronological order (van der Oost *et al.*, 2014), their sequences can be explored as molecular clocks to reveal the history of genetic invasion of a given isolate, and even to study the epidemiologic connection between different strains carrying a CRISPR of a common origin. With that prerogative, we explored the spacer sequences of CRISPRs located within *SCCmec* elements from different *Staphylococcus* species. As indicated by the high sequence identity between the *SCCmec*s, the identical sequence and organization of spacers evidenced that the mobile genetic element had been transferred between strains of canine *S. pseudintermedius* and *S. schleiferi*, and then to strains of *S. capitis* and *S. aureus* isolated from humans (Rossi *et al.*, 2017a; Rossi *et al.*, 2019).

### Concluding remarks

Most *Staphylococcus* species lack many of the *S. aureus* virulence factors and, because they are less frequently isolated from infectious processes, their impact in pathogenesis is usually overlooked. However, sequence and experimental evidences of horizontal transfer of antimicrobial resistance genes between them, favored by their capacity of forming biofilms and the scarcity of restrictive CRISPR-systems, show that these bacteria participate actively in the process of drug resistance spreading. Moreover, the fact that we exchange microbiota with our surroundings, makes it imperative that epidemiologic stud-

ies and strategies to control the advance of antimicrobial resistance consider the integrated nature of the relationship between human beings, animals, and the environment.

## Acknowledgments

The authors thank the financial support from Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ, grants E-26/203.037/2016, E-26/010.101056/2018 and E-26/201.451/2014), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, grant 305940/2018-0) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior Programa de Excelência Acadêmica – Finance Code 001 (CAPES ProEx, grant 23038.002486/2018-26).

## Conflict of Interest

The authors declare no conflict of interests.

## Author Contributions

CCR and MGM conceived and designed the study, CCR, MFP and MGM analyzed the data and wrote the manuscript. All authors read and approved the final version.

## References

- Abouelkhair MA, Bemis DA, Giannone RJ, Frank LA and Kania SA (2018) Characterization of a leukocidin identified in *Staphylococcus pseudintermedius*. *PLoS One* 13:e0204450.
- Allori MC, Jure MA, Romero C and Castillo ME (2006) Antimicrobial resistance and production of biofilms in clinical isolates of coagulase-negative *Staphylococcus* strains. *Biol Pharm Bull* 29:1592-1596.
- Almeida CC, Pizauro LJJ, Soltes GA, Slavic D, Ávila FA, Pizauro JM and Janet IM (2018) Some coagulase negative *Staphylococcus* spp. isolated from buffalo can be misidentified as *Staphylococcus aureus* by phenotypic and Sa442 PCR methods. *BMC Res Notes* 11:346.
- Amitai G and Sorek R (2016) CRISPR-Cas adaptation: Insights into the mechanism of action. *Nat Rev Microbiol* 14:67-76.
- Barros EM, Lemos M, Souto-Padron T and Giambiagi-deMarval M (2015) Phenotypic and genotypic characterization of biofilm formation in *Staphylococcus haemolyticus*. *Curr Microbiol* 70:829-834.
- Bastos MC, Mondino PJ, Azevedo ML, Santos KR and Giambiagi-deMarval M (1999) Molecular characterization and transfer among *Staphylococcus* strains of a plasmid conferring high-level resistance to mupirocin. *Eur J Clin Microbiol Infect Dis* 18:393-398.
- Becker K, Heilmann C and Peters G (2014) Coagulase-negative staphylococci. *Clin Microbiol Rev* 27:870-926.
- Black CE and Costerton JW (2010) Current concepts regarding the effect of wound microbial ecology and biofilms on wound healing. *Surg Clin North Am* 90:1147-1160.
- Buttner H, Mack D and Rohde H (2015) Structural basis of *Staphylococcus epidermidis* biofilm formation: Mechanisms and molecular interactions. *Front Cell Infect Microbiol* 5:14.
- Cabello FC, Godfrey HP, Tomova A, Ivanova L, Dolz H, Millanao A and Buschmann AH (2013) Antimicrobial use in aquaculture re-examined: Its relevance to antimicrobial resistance and to animal and human health. *Environ Microbiol* 15:1917-1942.
- Cafini F, Nguyen le TT, Higashide M, Roman F, Prieto J and Morikawa K (2016) Horizontal gene transmission of the *cf* gene to MRSA and *Enterococcus*: Role of *Staphylococcus epidermidis* as a reservoir and alternative pathway for the spread of linezolid resistance. *J Antimicrob Chemother* 71:587-592.
- Cao L, Gao CH, Zhu J, Zhao L, Wu Q, Li M and Sun B (2016) Identification and functional study of type III-A CRISPR-Cas systems in clinical isolates of *Staphylococcus aureus*. *Int J Med Microbiol* 306:686-696.
- Chang PH, Liu TP, Huang PY, Lin SY, Lin JF, Yeh CF, Chang SC, Wu TS and Lu JJ (2018) Clinical features, outcomes, and molecular characteristics of an outbreak of *Staphylococcus haemolyticus* infection, among a mass-burn casualty patient group, in a tertiary center in northern Taiwan. *J Microbiol Immunol Infect* 51:847-855.
- Czekaj T, Ciszewski M and Szewczyk EM (2015) *Staphylococcus haemolyticus* – an emerging threat in the twilight of the antibiotics age. *Microbiology* 161:2061–2068.
- Davey ME, Caiazza NC and O'Toole GA (2003) Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* 185:1027-1036.
- Devriese LA, Vancanneyt M, Baele M, Vanechoutte M, De Graef E, Snaauwaert C, Cleenwerck I, Dawyndt P, Swings J, Decostere A and Haesebrouck F (2005) *Staphylococcus pseudintermedius* sp. nov., a coagulase-positive species from animals. *Int J Syst Evol Microbiol* 55:1569-1573.
- Doroshenko N, Tseng BS, Howlin RP, Deacon J, Wharton JA, Thurner PJ, Gilmore BF, Parsek MR and Stoodley P (2014) Extracellular DNA impedes the transport of vancomycin in *Staphylococcus epidermidis* biofilms preexposed to subinhibitory concentrations of vancomycin. *Antimicrob Agents Chemother* 58:7273-7282.
- Dutta TK, Chakraborty S, Das M, Mandakini R, Vanrahmlim-phuii, Roychoudhury P, Ghorai S and Behera SK (2018) Multidrug-resistant *Staphylococcus pettenkoferi* isolated from cat in India. *Vet World* 11:1380-1384.
- Fitzgerald JR (2009) The *Staphylococcus intermedius* group of bacterial pathogens: Species re-classification, pathogenesis and the emergence of meticillin resistance. *Vet Dermatol* 20:490-495.
- Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA and Kjelleberg S (2016) Biofilms: An emergent form of bacterial life. *Nat Rev Microbiol* 14:563-575.
- Foster TJ, Geoghegan JA, Ganesh VK and Hook M (2014) Adhesion, invasion and evasion: The many functions of the surface proteins of *Staphylococcus aureus*. *Nat Rev Microbiol* 12:49-62.
- Ghebremedhin B, Layer F, König W and König B (2008) Genetic classification and distinguishing of *Staphylococcus* species based on different partial gap, 16s rRNA, *hsp60*, *rpoB*, *sodA*, and *tuf* gene sequences. *J Clin Microbiol* 46:1019-1025.
- Gomes F, Teixeira P and Oliveira R (2014) *Staphylococcus epidermidis* as the most frequent cause of nosocomial infections: Old and new fighting strategies. *Biofouling* 30:131-141.
- Guardabassi L, Schwarz S and Lloyd DH (2004) Pet animals as reservoirs of antimicrobial-resistant bacteria. *J Antimicrob Chemother* 54:321-332.

- Haaber J, Penades JR and Ingmer H (2017) Transfer of antibiotic resistance in *Staphylococcus aureus*. Trends Microbiol 25:893-905.
- Hannan S, Ready D, Jasni AS, Rogers M, Pratten J and Roberts AP (2010) Transfer of antibiotic resistance by transformation with eDNA within oral biofilms. FEMS Immunol Med Microbiol 59:345-349.
- Hardy KJ, Oppenheim BA, Gossain S, Gao F and Hawkey PM (2006) A study of the relationship between environmental contamination with methicillin-resistant *Staphylococcus aureus* (MRSA) and patients' acquisition of MRSA. Infect Control Hosp Epidemiol 27:127-132.
- Hatoum-Aslan A, Maniv I, Samai P and Marraffini LA (2014) Genetic characterization of antiplasmid immunity through a type III-A CRISPR-Cas system. J Bacteriol 196:310-317.
- Hauschild T, Stepanovic S and Zakrzewska-Czerwinska J (2010) *Staphylococcus stepanovicii* sp. nov., a novel novobiocin-resistant oxidase-positive staphylococcal species isolated from wild small mammals. Syst Appl Microbiol 33:183-187.
- Horvath P and Barrangou R (2010) CRISPR/Cas, the immune system of bacteria and archaea. Science 327:167-170.
- Hosseinkhani F, Tammes Buirs M, Jabalameh F, Emaneini M and van Leeuwen WB (2018) High diversity in SCCmec elements among multidrug-resistant *Staphylococcus haemolyticus* strains originating from paediatric patients; characterization of a new composite island. J Med Microbiol 67:915-921.
- Javid F, Taku A, Bhat MA, Badroo GA, Mudasar M and Sofi TA (2018) Molecular typing of *Staphylococcus aureus* based on coagulase gene. Vet World 11:423-430.
- King LJ, Anderson LR, Blackmore CG, Blackwell MJ, Lautner EA, Marcus LC, Meyer TE, Monath TP, Nave JE, Ohle J *et al.* (2008) Executive summary of the AVMA One Health Initiative Task Force report. J Am Vet Med Assoc 233:259-261.
- Kock R, Schaumburg F, Mellmann A, Koksai M, Jurke A, Becker K and Friedrich AW (2013) Livestock-associated methicillin-resistant *Staphylococcus aureus* (MRSA) as causes of human infection and colonization in Germany. PLoS One 8:e55040.
- Kohler V, Vaishampayan A and Grohmann E (2018) Broad-host-range Inc18 plasmids: Occurrence, spread and transfer mechanisms. Plasmid 99:11-21.
- Kraker ME, Stewardson AJ and Harbarth S (2016) Will 10 million people die a year due to antimicrobial resistance by 2050? PLoS Med 13:e1002184.
- Lax S, Smith DP, Hampton-Marcell J, Owens SM, Handley KM, Scott NM, Gibbons SM, Larsen P, Shogan BD, Weiss S *et al.* (2014) Longitudinal analysis of microbial interaction between humans and the indoor environment. Science 345:1048-1052.
- Laxminarayan R, Sridhar D, Blaser M, Wang M and Woolhouse M (2016) Achieving global targets for antimicrobial resistance. Science 353:874-875.
- Lee AS, de Lencastre H, Garau J, Kluytmans J, Malhotra-Kumar S, Peschel A and Harbarth S (2018) Methicillin-resistant *Staphylococcus aureus*. Nat Rev Dis Primers 4:18033.
- Lee CA, Thomas J and Grossman AD (2012) The *Bacillus subtilis* conjugative transposon ICEBs1 mobilizes plasmids lacking dedicated mobilization functions. J Bacteriol 194:3165-3172.
- Leid JG (2009) Bacterial biofilms resist key host defenses. Microbe 4:66-71.
- Li X, Xing J, Li B, Wang P and Liu J (2012) Use of *tuf* as a target for sequence-based identification of Gram-positive cocci of the genus *Enterococcus*, *Streptococcus*, coagulase-negative *Staphylococcus*, and *Lactococcus*. Ann Clin Microbiol Antimicrob 11:31.
- Li X, Arias CA, Aitken SL, Galloway Pena J, Panesso D, Chang M, Diaz L, Rios R, Numan Y, Ghaoui S *et al.* (2018) Clonal emergence of invasive multidrug-resistant *Staphylococcus epidermidis* deconvoluted via a combination of whole-genome sequencing and microbiome analyses. Clin Infect Dis 67:398-406.
- Liu P, Wu Z, Xue H and Zhao X (2017) Antibiotics trigger initiation of SCCmec transfer by inducing SOS responses. Nucleic Acids Res 45:3944-3952.
- Liu X, Ng C and Ferenci T (2000) Global adaptations resulting from high population densities in *Escherichia coli* cultures. J Bacteriol 182:4158-4164.
- Lozano C, Rezusta A, Ferrer I, Perez-Laguna V, Zarazaga M, Ruiz-Ripa L, Revillo MJ and Torres C (2017) *Staphylococcus pseudintermedius* human infection cases in Spain: Dog-to-human transmission. Vector Borne Zoonotic Dis 17:268-270.
- Maali Y, Badiou C, Martins-Simoes P, Hodille E, Bes M, Vandenesch F, Lina G, Diot A, Laurent F and Trouillet-Assant S (2018) Understanding the virulence of *Staphylococcus pseudintermedius*: a major role of pore-forming toxins. Front Cell Infect Microbiol 8:221.
- Madsen JS, Burmolle M, Hansen LH and Sorensen SJ (2012) The interconnection between biofilm formation and horizontal gene transfer. FEMS Immunol Med Microbiol 65:183-195.
- Marraffini LA and Sontheimer EJ (2008) CRISPR interference limits horizontal gene transfer in staphylococci by targeting DNA. Science 322:1843-1845.
- Marshall BM and Levy SB (2011) Food animals and antimicrobials: Impacts on human health. Clin Microbiol Reviews 24:718-733.
- Meric G, Miragaia M, de Been M, Yahara K, Pascoe B, Mageiros L, Mikhail J, Harris LG, Wilkinson TS, Rolo J *et al.* (2015) Ecological overlap and horizontal gene transfer in *Staphylococcus aureus* and *Staphylococcus epidermidis*. Genome Biol Evol 7:1313-1328.
- Mojica FJ, Diez-Villasenor C, Garcia-Martinez J and Soria E (2005) Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. J Mol Evol 60:174-182.
- Moon BY, Park JY, Hwang SY, Robinson DA, Thomas JC, Fitzgerald JR, Park YH and Seo KS (2015) Phage-mediated horizontal transfer of a *Staphylococcus aureus* virulence-associated genomic island. Sci Rep 5:9784.
- Morfin-Otero R, Martinez-Vazquez MA, Lopez D, Rodriguez-Noriega E and Garza-Gonzalez E (2012) Isolation of rare coagulase-negative isolates in immunocompromised patients: *Staphylococcus gallinarum*, *Staphylococcus pettenkoferi* and *Staphylococcus pasteurii*. Ann Clin Lab Sci 42:182-185.
- Morikawa K, Takemura AJ, Inose Y, Tsai M, Nguyen Thi le T, Ohta T and Msadek T (2012) Expression of a cryptic secondary sigma factor gene unveils natural competence for DNA transformation in *Staphylococcus aureus*. PLoS Pathog 8:e1003003.



- Moroni P, Pisoni G, Antonini M, Ruffo G, Carli S, Varisco G and Boettcher P (2005) Subclinical mastitis and antimicrobial susceptibility of *Staphylococcus caprae* and *Staphylococcus epidermidis* isolated from two Italian goat herds. *J Dairy Sci* 88:1694-1704.
- Murray AK, Lee J, Bendall R, Zhang L, Sunde M, Schau Slettemeas J, Gaze W, Page AJ and Vos M (2018) *Staphylococcus cornubiensis* sp. nov., a member of the *Staphylococcus intermedius* Group (SIG). *Int J Syst Evol Microbiol* 68:3404-3408.
- Novakova D, Pantucek R, Petras P, Koukalova D and Sedlacek I (2006a) Occurrence of *Staphylococcus nepalensis* strains in different sources including human clinical material. *FEMS Microbiol Lett* 263:163-168.
- Novakova D, Sedlacek I, Pantucek R, Stetina V, Svec P and Petras P (2006b) *Staphylococcus equorum* and *Staphylococcus succinus* isolated from human clinical specimens. *J Med Microbiol* 55:523-528.
- O'Brien FG, Ramsay JP, Monecke S, Coombs GW, Robinson OJ, Htet Z, Alshaikh FA and Grubb WB (2015a) *Staphylococcus aureus* plasmids without mobilization genes are mobilized by a novel conjugative plasmid from community isolates. *J Antimicrob Chemother* 70:649-652.
- O'Brien FG, Yui Eto K, Murphy RJ, Fairhurst HM, Coombs GW, Grubb WB and Ramsay JP (2015b) Origin-of-transfer sequences facilitate mobilisation of non-conjugative antimicrobial-resistance plasmids in *Staphylococcus aureus*. *Nucleic Acids Res* 43:7971-7983.
- O'Neill J (2014) Antimicrobial resistance: tackling a crisis for the health and wealth of nations. *Rev Antimicrob Resist* 2014:1-20.
- Okshevsky M and Meyer RL (2015) The role of extracellular DNA in the establishment, maintenance and perpetuation of bacterial biofilms. *Crit Rev Microbiol* 41:341-352.
- Oliveira WF, Silva PMS, Silva RCS, Silva GMM, Machado G, Coelho LCBB and Correia MTS (2018) *Staphylococcus aureus* and *Staphylococcus epidermidis* infections on implants. *J Hosp Infect* 98:111-117.
- Otto M (2013) Coagulase-negative staphylococci as reservoirs of genes facilitating MRSA infection: Staphylococcal commensal species such as *Staphylococcus epidermidis* are being recognized as important sources of genes promoting MRSA colonization and virulence. *Bioessays* 35:4-11.
- Otto M (2014) *Staphylococcus aureus* toxins. *Curr Opin Microbiol* 17:32-37.
- Parte AC (2018) LPSN - List of prokaryotic names with standing in nomenclature (bacterio.net), 20 years on. *Int J Syst Evol Microbiol* 68:1825-1829.
- Patel JB, Gorwitz RJ and Jernigan JA (2009) Mupirocin resistance. *Clin Infect Dis* 49:935-941.
- Poirel L, Jayol A and Nordmann P (2017) Polymyxins: Antibacterial activity, susceptibility testing, and resistance mechanisms encoded by plasmids or chromosomes. *Clin Microbiol Rev* 30:557-596.
- Powers ME and Wardenburg JB (2014) Igniting the Fire: *Staphylococcus aureus* virulence factors in the pathogenesis of sepsis. *PLoS Pathog* 10:e1003871.
- Ramsay JP, Kwong SM, Murphy RJ, Yui Eto K, Price KJ, Nguyen QT, O'Brien FG, Grubb WB, Coombs GW and Firth N (2016) An updated view of plasmid conjugation and mobilization in *Staphylococcus*. *Mob Genet Elements* 6:e1208317.
- Riegel P, Jesel-Morel L, Laventie B, Boisset S, Vandenesch F and Prevost G (2011) Coagulase-positive *Staphylococcus pseudintermedius* from animals causing human endocarditis. *Int J Med Microbiol* 301:237-239.
- Robinson TP, Bu DP, Carrique-Mas J, Fevre EM, Gilbert M, Grace D, Hay SI, Jiwakanon J, Kakkar M, Kariuki S *et al.* (2016) Antibiotic resistance is the quintessential One Health issue. *Trans R Soc Trop Med Hyg* 110:377-380.
- Rolo J, Worning P, Nielsen JB, Bowden R, Bouchami O, Damborg P, Guardabassi L, Perreten V, Tomasz A, Westh H *et al.* (2017) Evolutionary origin of the staphylococcal cassette chromosome mec (SCCmec). *Antimicrob Agents Chemother* 61:e02302-16.
- Rossi CC, Ferreira NC, Coelho ML, Schuenck RP, Bastos MC and Giambiagi-deMarval M (2016) Transfer of mupirocin resistance from *Staphylococcus haemolyticus* clinical strains to *Staphylococcus aureus* through conjugative and mobilizable plasmids. *FEMS Microbiol Lett* 363:fnw121.
- Rossi CC, Souza-Silva T, Araujo-Alves AV and Giambiagi-deMarval M (2017a) CRISPR-Cas systems features and the gene-reservoir role of coagulase-negative staphylococci. *Front Microbiol* 8:1545.
- Rossi CC, da Silva Dias I, Muniz IM, Lilenbaum W and Giambiagi-deMarval M (2017b) The oral microbiota of domestic cats harbors a wide variety of *Staphylococcus* species with zoonotic potential. *Vet Microbiol* 201:136-140.
- Rossi CC, de Oliveira LL, de Carvalho Rodrigues D, Urmenyi TP, Laport MS and Giambiagi-deMarval M (2017c) Expression of the stress-response regulators CtsR and HrcA in the uropathogen *Staphylococcus saprophyticus* during heat shock. *Antonie van Leeuwenhoek* 110:1105-1111.
- Rossi CC, Salgado BAB, Barros EM, de Campos Braga PA, Eberlin MN, Lilenbaum W and Giambiagi-deMarval M (2018) Identification of *Staphylococcus epidermidis* with transferrable mupirocin resistance from canine skin. *Vet J* 235:70-72.
- Rossi CC, Andrade-Oliveira AL and Giambiagi-deMarval M (2019) CRISPR tracking reveals global spreading of antimicrobial resistance genes by *Staphylococcus* of canine origin. *Vet Microbiol* 232:65-69.
- Ruscher C, Lubke-Becker A, Wleklinski CG, Soba A, Wieler LH and Walther B (2009) Prevalence of Methicillin-resistant *Staphylococcus pseudintermedius* isolated from clinical samples of companion animals and equidae. *Vet Microbiol* 136:197-201.
- Saber H, Jasni AS, Jamaluddin T and Ibrahim R (2017) A Review of staphylococcal cassette chromosome mec (SCCmec) types in coagulase-negative staphylococci (CoNS) species. *Malays J Med Sci* 24:7-18.
- Salehzadeh A, Zamani H, Langeroudi MK and Mirzaie A (2016) Molecular typing of nosocomial *Staphylococcus aureus* strains associated to biofilm based on the coagulase and protein A gene polymorphisms. *Iran J Basic Med Sci* 19:1325-1330.
- Santos DC, Lange CC, Avellar-Costa P, Santos KRN, Brito MAVP and Giambiagi-deMarval M (2016) *Staphylococcus chromogenes*, a coagulase-negative *Staphylococcus* species that can clot plasma. *J Clin Microbiol* 54:1372-1375.
- Sasaki T, Kikuchi K, Tanaka Y, Takahashi N, Kamata S and Hiramatsu K (2007) Reclassification of phenotypically identified *Staphylococcus intermedius* strains. *J Clin Microbiol* 45:2770-2778.

- Savage VJ, Chopra I and O'Neill AJ (2013) *Staphylococcus aureus* biofilms promote horizontal transfer of antibiotic resistance. *Antimicrob Agents Chemother* 57:1968-1970.
- Schwarz S, Fessler AT, Loncaric I, Wu C, Kadlec K, Wang Y and Shen J (2018) Antimicrobial resistance among staphylococci of animal origin. *Microbiol Spectr* 6:4.
- Sexton T, Clarke P, O'Neill E, Dillane T and Humphreys H (2006) Environmental reservoirs of methicillin-resistant *Staphylococcus aureus* in isolation rooms: correlation with patient isolates and implications for hospital hygiene. *J Hosp Infect* 62:187-194.
- Silbergeld EK, Graham J and Price LB (2008) Industrial food animal production, antimicrobial resistance, and human health. *Ann Rev Public Health* 29:151-169.
- Singhal N, Kumar M, Kanaujia PK and Viridi JS (2015) MALDI-TOF mass spectrometry: An emerging technology for microbial identification and diagnosis. *Front Microbiol* 6:791.
- Song SJ, Lauber C, Costello EK, Lozupone CA, Humphrey G, Berg-Lyons D, Caporaso JG, Knights D, Clemente JC, Nakielny S *et al.* (2013) Cohabiting family members share microbiota with one another and with their dogs. *eLife* 2:e00458.
- Spergser J, Wieser M, Taubel M, Rossello-Mora RA, Rosengarten R and Busse HJ (2003) *Staphylococcus nepalensis* sp. nov., isolated from goats of the Himalayan region. *Int J Syst Evol Microbiol* 53:2007-2011.
- Sugimoto S, Sato F, Miyakawa R, Chiba A, Onodera S, Hori S and Mizunoe Y (2018) Broad impact of extracellular DNA on biofilm formation by clinically isolated Methicillin-resistant and -sensitive strains of *Staphylococcus aureus*. *Sci Rep* 8:2254.
- Sundareshan S, Isloor S, Babu YH, Sunagar R, Sheela P, Tiwari JG, Wryyah CB, Mukkur TK and Hegde NR (2017) Isolation and molecular identification of rare coagulase-negative *Staphylococcus aureus* variants isolated from bovine milk samples. *Indian J Comp Microbiol Immunol Infect Dis* 38:66-73.
- Takeuchi F, Watanabe S, Baba T, Yuzawa H, Ito T, Morimoto Y, Kuroda M, Cui L, Takahashi M, Ankaï A *et al.* (2005) Whole-genome sequencing of *Staphylococcus haemolyticus* uncovers the extreme plasticity of its genome and the evolution of human-colonizing staphylococcal species. *J Bacteriol* 187:7292-7308.
- Taponen S, Supre K, Piessens V, Van Coillie E, De Vliegher S and Koort JM (2012) *Staphylococcus agnetis* sp. nov., a coagulase-variable species from bovine subclinical and mild clinical mastitis. *Int J Syst Evol Microbiol* 62:61-65.
- Thi LTN, Romero VM and Morikawa K (2016) Cell wall-affecting antibiotics modulate natural transformation in SigH-expressing *Staphylococcus aureus*. *J Antibiot* 69:464-466.
- Tong SY, Davis JS, Eichenberger E, Holland TL and Fowler VG (2015) *Staphylococcus aureus* infections: Epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* 28:603-661.
- Uribe-Alvarez C, Chiquete-Felix N, Contreras-Zentella M, Guerrero-Castillo S, Pena A and Uribe-Carvajal S (2016) *Staphylococcus epidermidis*: Metabolic adaptation and biofilm formation in response to different oxygen concentrations. *Pathog Dis* 74:ftv111.
- Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA and Laxminarayan R (2014) Global antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data. *Lancet Infect Dis* 14:742-750.
- Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP, Teillant A and Laxminarayan R (2015) Global trends in antimicrobial use in food animals. *Proc Natl Acad Sci U S A* 112:5649-5654.
- van der Oost J, Westra ER, Jackson RN and Wiedenhoff B (2014) Unravelling the structural and mechanistic basis of CRISPR-Cas systems. *Nat Rev Microbiol* 12:479-492.
- Van Hoovels L, Vankeerberghen A, Boel A, Van Vaerenbergh K and de Beenhouwer H (2006) First case of *Staphylococcus pseudintermedius* infection in a human. *J Clin Microbiol* 44:4609-4612.
- Vanderhaeghen W, Hermans K, Haesebrouck F and Butaye P (2010) Methicillin-resistant *Staphylococcus aureus* (MRSA) in food production animals. *Epidemiol Infect* 138:606-625.
- Varaldo PE, Kilpper-Bälz R, Biavasco F, Satta G and Schleifer KH (1988) *Staphylococcus delphini* sp. nov., a coagulase-positive species isolated from dolphins. *Int J Syst Evol Microbiol* 38:436-439.
- Vorkapic D, Pressler K and Schild S (2016) Multifaceted roles of extracellular DNA in bacterial physiology. *Curr Genet* 62:71-79.
- Wang R, Khan BA, Cheung GY, Bach TH, Jameson-Lee M, Kong KF, Queck SY and Otto M (2011) *Staphylococcus epidermidis* surfactant peptides promote biofilm maturation and dissemination of biofilm-associated infection in mice. *J Clin Invest* 121:238-248.
- Wendlandt S, Shen J, Kadlec K, Wang Y, Li B, Zhang WJ, Fessler AT, Wu C and Schwarz S (2015) Multidrug resistance genes in staphylococci from animals that confer resistance to critically and highly important antimicrobial agents in human medicine. *Trends Microbiol* 23:44-54.
- WHO - World Health Organization (2017) Global action plan on antimicrobial resistance. WHO Document Production Services, Geneva.
- Widerstrom M, Wistrom J, Edebro H, Marklund E, Backman M, Lindqvist P and Monsen T (2016) Colonization of patients, healthcare workers, and the environment with healthcare-associated *Staphylococcus epidermidis* genotypes in an intensive care unit: a prospective observational cohort study. *BMC Infect Dis* 16:743.
- Worthing K, Pang S, Trott DJ, Abraham S, Coombs GW, Jordan D, McIntyre L, Davies MR and Norris J (2018) Characterisation of *Staphylococcus felis* isolated from cats using whole genome sequencing. *Vet Microbiol* 222:98-104.
- Zhang YQ, Ren SX, Li HL, Wang YX, Fu G, Yang J, Qin ZQ, Miao YG, Wang WY, Chen RS *et al.* (2003) Genome-based analysis of virulence genes in a non-biofilm-forming *Staphylococcus epidermidis* strain (ATCC 12228). *Mol Microbiol* 49:1577-1593.
- Zurek L and Ghosh A (2014) Insects represent a link between food animal farms and the urban environment for antibiotic resistance traits. *Appl Environ Microbiol* 80:3562-3567.

Associate Editor: Célia Maria de Almeida Soares