



Sulfadiazine-induced crystalluria and non-oliguric renal failure in HIV-1 inaugural infection with presumed cerebral toxoplasmosis

Cristalúria induzida por sulfadiazina e insuficiência renal não oligúrica em infecção inaugural por HIV-1 com toxoplasmose cerebral presumida

Authors

Vasco Gaspar^{1,2} 

Nuno Moreira Fonseca^{2,3} 

Sara Lino^{2,4} 

¹Centro Hospital Universitário Lisboa Central, Departamento de Medicina Interna, Lisboa, Portugal.

²Universidade NOVA de Lisboa, Faculdade de Medicina NOVA, Lisboa, Portugal.

³Centro Hospital Universitário Lisboa Central, Hospital Curry Cabral, Departamento de Nefrologia, Lisboa, Portugal.

⁴Centro Hospital Universitário Lisboa Central, Hospital Curry Cabral, Departamento de Infeciologia, Lisboa, Portugal.

A 30-year-old woman with no previous medical history presented to the emergency department with a seizure and aggressive behavior. After extensive blood work and imaging studies, she was admitted with HIV inaugural infection (CDC stage C3) and cerebral toxoplasmosis. As the first line of treatment, the patient was started on sulfadiazine and pyrimethamine¹⁻³. After one week, she developed non-oliguric

acute kidney injury. Urinary sediment analysis revealed sulfonamide crystals with the morphologic appearance of shocks of wheat (Figures 1 and 2), confirmed by infrared spectroscopy⁴. Sulfadiazine was replaced with clindamycin, and a notable enhancement was observed after to the implementation of vigorous fluid hydration using an alkaline solution (sodium bicarbonate).



Figure 1. Sulfadiazine crystals have an amber color and radial striations (contrast phase microscopy, 400× magnification). Urinary analysis results – density: 1.008; pH: 5; proteins: 15 mg/dL; hemoglobin: 0.75 mg/dL; nitrites/glucose/ketones/bilirubin/urobilinogen: negative.

Submitted on: 10/08/2023.

Approved on: 03/22/2024.

Published on: 05/10/2024.

Correspondence to:

Vasco Gaspar.

Email: gaspar.vasco@gmail.com

DOI: <https://doi.org/10.1590/2175-8239-JBN-2023-0151en>




Figure 2. Sulfadiazine crystals are strongly birefringent under polarized light (polarized light, magnification 400×).

ACKNOWLEDGMENTS

We thank Dr. Vincent Frochot for the infrared spectroscopy analysis.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

AUTHORS' CONTRIBUTIONS

VG writing of the original draft. NMF, SL writing – review and editing.

REFERENCES

1. Elsheikha HM, Marra CM, Zhu XQ. Epidemiology, pathophysiology, diagnosis, and management of cerebral toxoplasmosis. *Clin Microbiol Rev.* 2020;34(1):e00115–19. doi: <http://dx.doi.org/10.1128/CMR.00115-19>. PubMed PMID: 33239310.
2. Perazella MA, Herlitz LC. The crystalline nephropathies. *Kidney Int Rep.* 2021;6(12):2942–57. doi: <http://dx.doi.org/10.1016/j.ekir.2021.09.003>. PubMed PMID: 34901567.
3. Chebion G, Bugni E, Gerin V, Daudon M, Castiglione V. Drug-induced nephrolithiasis and crystalluria: the particular case of the sulfasalazine derivatives. *C R Chim.* 2022;25(S1):295–306. doi: <http://dx.doi.org/10.5802/crchim.109>.
4. Fogazzi GB. *The urinary sediment: an integrated view.* 3rd ed. Milan: Elsevier; 2010. p. 159–61.