

SCIENTIFIC ARTICLE

Knowledge retention after focused cardiac ultrasound training: a prospective cohort pilot study

Glenio B. Mizubuti^a, Rene V. Allard^a, Anthony M.-H. Ho^a, Louie Wang^a, Theresa Beesley^b, Wilma M. Hopman^c, Rylan Egan^d, Devin Sydor^a, Dale Engen^a, Tarit Saha^a, Robert C. Tanzola^{Id, a,*}

^a Queen's University, Kingston General Hospital, Department of Anesthesiology and Perioperative Medicine, Kingston, Canada

^b McGill University, Department of Medicine, Montreal, Canada

^c Queen's University, Kingston Sciences Centre Research Institute and Public Health Sciences, Kingston, Canada

^d Queen's University, Healthcare Quality Graduate Programs, Kingston, Canada

Received 10 February 2018; accepted 31 October 2018

Available online 7 December 2018

KEYWORDS

POCUS;
FoCUS;
Knowledge retention;
Education;
Anesthesia

Abstract

Background and objectives: Focused Cardiac Ultrasound (FoCUS) has proven instrumental in guiding anesthesiologists' clinical decision-making process. Training residents to perform and interpret FoCUS is both feasible and effective. However, the degree of knowledge retention after FoCUS training remains a subject of debate. We sought to provide a description of our 4-week FoCUS curriculum, and to assess the knowledge retention among anesthesia residents at 6 months after FoCUS rotation.

Methods: A prospective analysis involving eleven senior anesthesia residents was carried out. At end of FoCUS Rotation (EOR) participants completed a questionnaire (evaluating the number of scans completed and residents' self-rated knowledge and comfort level with FoCUS), and a multiple-choice FoCUS exam comprised of written- and video-based questions. Six months later, participants completed a follow-up questionnaire and a similar exam. Self-rated knowledge and exam scores were compared at EOR and after 6 months. Spearman correlations were conducted to test the relationship between number of scans completed and exam scores, perceived knowledge and exam scores, and number of scans and perceived knowledge.

Results: Mean exam scores (out of 50) were 44.1 at EOR and 43 at the 6-month follow-up. Residents had significantly higher perceived knowledge (out of 10) at EOR (8.0) than at the 6-month follow-up (5.5), $p=0.003$. At the EOR, all trainees felt comfortable using FoCUS, and at 6 months 10/11 still felt comfortable. All the trainees had used FoCUS in their clinical practice after EOR, and the most cited reason for not using FoCUS more frequently was the lack of perceived clinical need. A strong and statistically significant ($\rho=0.804$, $p=0.005$) correlation between number of scans completed during the FoCUS rotation and 6-month follow-up perceived knowledge was observed.

* Corresponding author.

E-mail: Rob.Tanzola@kingstonhsc.ca (R.C. Tanzola).



Conclusion: Four weeks of intensive FoCUS training results in adequate knowledge acquisition and 6-month knowledge retention.

© 2018 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALAVRAS-CHAVE

POCUS;
USCLA;
Retenção de
conhecimento;
Educação;
Anestesia

Retenção do conhecimento após treinamento de ultrassonografia cardíaca focada: estudo-piloto prospectivo de coorte

Resumo

Justificativa e objetivos: A ultrassonografia cardíaca no local de atendimento (USCLA) provou ser importante para orientar o processo de tomada de decisão clínica dos anestesiologistas. Treinar os residentes para fazer e interpretar uma USCLA é viável e eficaz. No entanto, o grau de retenção do conhecimento após o treinamento permanece um assunto de debate. Procuramos fornecer uma descrição do currículo de quatro semanas do treinamento de USCLA e avaliar a retenção do conhecimento entre os residentes de anestesia seis meses após a rotação em USCLA.

Métodos: Uma análise prospectiva foi realizada com 11 residentes seniores de anestesia. Ao final da rotação em USCLA, os participantes preencheram um questionário (avaliando o número de exames ultrassonográficos concluídos, o conhecimento adquirido e o nível de conforto dos residentes com a USCLA) e fizeram um exame de múltipla escolha para USCLA, composto por perguntas escritas e baseadas em vídeo. Seis meses depois, os participantes preencheram um questionário de acompanhamento e um exame similar. A autoavaliação do conhecimento e os escores do exame foram comparados no final da rotação e após seis meses. Correlações de Spearman foram usadas para testar a relação entre o número de exames concluídos e os escores dos exames, o conhecimento percebido, os escores dos exames, o número de exames e o conhecimento percebido.

Resultados: Os escores médios dos exames (50) foram: 44,1 no final da rotação e 43 após seis meses. Os residentes tiveram conhecimento percebido significativamente maior (10) no final da rotação (8,0) que após seis meses (5,5), $p = 0,003$. No final da rotação, todos os residentes se sentiram confortáveis usando o aparelho de USCLA e, aos seis meses, 10/11 ainda se sentiam confortáveis. Todos os residentes haviam usado o USCLA em sua prática clínica após o final da rotação e a razão mais citada para não usar o USCLA com mais frequência foi a falta de necessidade clínica percebida. Uma correlação forte e estatisticamente significativa ($\rho = 0,804$, $p = 0,005$) foi observada entre o número de exames realizados durante a rotação em USCLA e o conhecimento percebido em seis meses de seguimento.

Conclusão: Quatro semanas de treinamento intensivo de USCLA resultaram em aquisição e retenção adequadas do conhecimento por seis meses.

© 2018 Sociedade Brasileira de Anestesiologia. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Point-of-Care Ultrasound (POCUS) has become a key diagnostic tool for several medical specialties including anesthesiology, emergency medicine, critical care, and internal medicine.¹⁻³ POCUS training is now an integral component of undergraduate medical curricula in several schools.^{4,5} Focused Cardiac Ultrasound (FoCUS), one of the POCUS modalities, is a valuable diagnostic tool in a variety of clinical settings.⁶ For instance, FoCUS-guided therapy in subacute shock is associated with improved survival.⁷ When performed by anesthesiologists, FoCUS can be instrumental in the clinical decision-making process, providing invaluable information that changes perioperative care during both elective and emergency situations.⁸⁻¹²

Training residents to perform and interpret FoCUS is both feasible and effective.¹³⁻¹⁵ Accordingly, a FoCUS rotation has recently been introduced into the anesthesia residency training program at Queen's University, providing residents with an opportunity to integrate this diagnostic modality into their clinical armamentarium. However, the degree of knowledge retention after this FoCUS training has yet to be determined. This is important because while many educational interventions have shown immediate gains in knowledge and skills, these gains are often subject to subsequent decay over a period of time.¹⁶ We hypothesized that 4 weeks of intensive FoCUS training would allow trainees to maintain knowledge and interpretation skills 6 months later. The objective of this pilot study, therefore, was to assess the knowledge retention (i.e., theoretical knowledge,

interpretation and clinical correlation skills) at 6 months among anesthesia residents following the FoCUS rotation. Additionally, we sought to provide a thorough description of our current curriculum (and the solutions we have found over time to major barriers to its implementation and maintenance), in the hopes that it may serve as a template to institutions willing to establish a FoCUS training program.

Structure of rotation

The anesthesia residency training program at Queen's University has a well established 4-week FoCUS rotation that allows its residents to pursue Level I (basic) competence in FoCUS.^{13,17} Learning objectives of the rotation include residents obtaining and displaying an understanding of Ultrasound (US) basic principles, showing consistently the ability to acquire and interpret images from 3 Windows (parasternal, apical and subcostal, including assessment of the inferior vena cava), and most importantly, showing the ability to integrate the acquired information into the clinical context. Given the current lack of formal training guidelines for FoCUS in anesthesiology, the structure and learning objectives of our curriculum were adapted from the American College of Chest Physicians statement on basic critical care echocardiography.¹⁸

The rotation (Fig. 1) consists of weekly didactic teaching sessions with staff anesthesiologists trained in echocardiography, a commercially available online course,¹⁹ scanning practice sessions on an echo simulator, followed by supervised and independent scanning of perioperative patients. The rotation is divided in three phases.

In Phase 1, trainees are required to complete "Part 1: Basic Principles and 2-D Echo" and "Part 2: Color Doppler, M-Mode and Harmonic Imaging" of a 6-part online course (CAE Healthcare (Montreal, Canada) ICCU "Focused Transthoracic Cardiac Echo" – 7.5 hour course).¹⁹ Successful completion of Parts 1 and 2 is achieved by obtaining a set pass score.

In Phase 2, residents complete the remaining modules of the online course – "Parasternal long-axis", "Parasternal short-axis", "Apical four- and five-chamber", and "Subcostal four-chamber, IVC, aorta",¹⁹ and supplement it with didactic mini-seminars of basic topics (US principles, knobology, basic views, left and right ventricles – structure and function, Doppler, bedside hemodynamic and volume assessment, and aortic stenosis – although briefly discussed throughout the rotation, formal training in other valvulopathies is not part of the curriculum) delivered throughout the rotation by attending anesthesiologists trained in echocardiography. Residents also begin scanning practice sessions on a simulator (CAE Vimedix US simulator), followed by supervised bedside scanning on patients. After 10 supervised exams, residents begin Phase 3 – independent scanning of perioperative patients with weekly preceptor debriefing.

During Phase 3, residents electively scan pre- and post-operative patients for educational purposes, and perform the initial perioperative scan of patients with a clinical indication. Images and/or video clips are recorded for subsequent review by a certified staff supervisor. Residents complete a report on all patients they have scanned. They also have at least 6 opportunities to shadow a sonographer in the cardiology echocardiography lab (Fig. 1) to further improve scanning technique. On 6 occasions a staff anesthesiologist supervisor reviews the scans performed in the preceding days and completes a feedback form – detailing strengths and areas for improvement. Additionally, trainees use these review days for proctored scanning, with direct real-time feedback from the supervisor.

In order to be deemed successful in the FoCUS rotation, trainees are required to (1) complete a minimum of fifty FoCUS scans reviewed by a staff supervisor, (2) complete the online course along with an acceptable "pass-score" (80%) in its final exam, (3) complete a written (multiple-choice) and practical (video clip-based) exam at the End Of the Rotation (EOR) with a minimum score of 80%, (4) complete an observed practical bedside FoCUS exam at the EOR with

	Monday	Tuesday	Wednesday	Thursday	Friday
Week 1		CAE-PLAX Introduction and views (AM) Simulator (PM)	CAE-PLAX	CAE-PLAX Review scans	CAE-PLAX Review scans
Week 2	CAE-PSAX LV and RV function Echo Lab (PM)	CAE-PSAX Echo Lab (PM)	CAE-PSAX	CAE - PSAX Review scans	CAE - PSAX
Week 3	CAE-apical 4/5c Pleura/pericardium Echo Lab (PM)	CAE - apical 4/5c Echo Lab (PM)	CAE - Apical 4/5C	CAE - apical 4/5C Review scans	CAE - apical 4/5C Review scans
Week 4	CAE-SC/IVC Volume status and hemodynamics Echo Lab (PM)	CAE - SC/IVC Echo Lab (PM)	CAE - SC/IVC	CAE - SC/IVC Review scans	CAE - SC/IVC Case Management Rounds
Final day	Final Exam Final Assessment				

Figure 1 Outline of FoCUS rotation. CAE, "Focused Transthoracic Cardiac Echo" online course by CAE Healthcare; PLAX, Parasternal long axis view; PSAX, Parasternal short axis view; Apical 4/5C, Apical 4-chamber and 5-chamber views; SC/IVC, subcostal and inferior vena cava views.

adequate image acquisition and interpretation as assessed by a staff supervisor, and (5) complete a Case Management Rounds presentation at the EOR, outlining the specific FoCUS findings and its impact on perioperative patient management.

Methods

Study design and participants

Following approval from the Queen's University Research Ethics Board, senior anesthesiology residents (Postgraduate Years – PGY: 3–5) were recruited to the study on a voluntary basis and gave written informed consent. The intervention was the Queen's University Anesthesiology FoCUS rotation. A prospective analysis was undertaken from July 2013 to June 2015.

Data collection

At the EOR, participants completed a brief questionnaire including PGY of training, number of scans completed, resident's self-rated knowledge (10-point scale) pre- and post-FoCUS rotation, and comfort level (yes/no) with FoCUS. In addition, they completed an exam consisting of 50 multiple choice questions (37 written- and 13 video-based) that assessed US fundamentals, anatomy, imaging windows, and interpretation of static images and video clips of normal and abnormal findings. Six months later, residents completed a follow-up questionnaire (aiming to collect similar data in addition to barriers to the application of FoCUS in clinical practice) and an exam consisting of similar content and questions. Self-rated knowledge and exam scores were compared between EOR and 6-month follow-up.

Statistical analysis

Data were analyzed using the Statistical Package for the Social Sciences (SPSS 24; IBM, Armonk, NY, USA, 2016). Descriptive statistics were provided for the EOR and 6-month follow-up questionnaire data. Non-parametric tests of association were used due to the small sample size. Wilcoxon signed rank tests were used to compare total, video- and written-based exam scores as well as perceived knowledge between EOR and 6-month follow-up. Spearman correlations were conducted to test the relationship between number of scans completed and exam scores, perceived knowledge and exam scores, and number of scans and perceived knowledge.

Results

Eleven anesthesiology residents completed the FoCUS rotation and the 6-month follow-up assessment. The mean exam score (out of 50) was 44.1 at the EOR and 43 for the 6-month follow-up (Table 1). The mean video exam scores (out of 13) for the two time points were 12.3 and 11.5 respectively; the equivalent values for the written exam scores (out of 37) were 31.8 and 31.5. Fig. 2 provides the median values; none of the three comparisons attained statistical significance using the Wilcoxon signed rank test.

At the EOR, all trainees felt comfortable using FoCUS in their clinical practice, whereas at 6 months 10 out of 11 trainees still felt comfortable. Residents had significantly higher perceived knowledge (out of 10) at EOR (8.0) than at the 6-month follow-up (5.5), $p = 0.003$, but their knowledge perception was not statistically significantly correlated with their exam scores either at the EOR ($\rho = 0.32$, $p = 0.34$), or at the 6-month follow-up ($\rho = 0.17$, $p = 0.62$).

The average number of scans completed during the FoCUS rotation and the following 6 months were 61 and 4, respectively, and the most cited reason for not using FoCUS more frequently was the lack of perceived clinical need (Table 1). There was no correlation between number of scans completed during the FoCUS rotation and the post exam score ($\rho = 0.032$, $p = 0.93$), but there was a strong and statistically significant ($\rho = 0.804$, $p = 0.005$) correlation between number of scans completed during the FoCUS rotation and 6-month follow-up perceived knowledge.

Discussion

One goal of medical education is to promote long-term knowledge and skill retention. Following a 13-day US course, novice anesthesia interns' scores in a knowledge test involving basic echocardiography and perioperative US were sustained and similar to senior residents 90 days later.²⁰ Similarly, residents' knowledge of ultrasound increased immediately after a 2.5-hour US workshop (which did not include cardiac imaging), but significantly decreased in a follow-up 12-month assessment.²¹ In the present study, residents experienced a negligible decrease in their exam scores 6 months after the completion of the FoCUS rotation (Fig. 2). Whereas infrequently used knowledge and skills generally decay over time, this was not the finding of this study at 6 months. In fact, despite the residents' perception that their knowledge was worse at 6 months (perhaps because of the less frequent clinical use of FoCUS after the EOR), this was not corroborated by their exam scores between the time points that they were tested. The explanation for such finding is likely multifactorial. Firstly, the fact that all trainees have incorporated FoCUS into their clinical practice may be considered a contributing factor as it is accepted that constant reinforcement of new knowledge leads to longer-term retention.²² In fact, dispersed learning and reinforcement in this manner is associated with improved long-term knowledge retention.²³ Secondly, proctored US training, a component of our FoCUS curriculum, has been shown to yield knowledge retention.²⁴ Finally, and perhaps most importantly, over-learning in the initial phases of knowledge acquisition is associated with strong long-term memory,²⁵ and this might have occurred in our cohort given the robustness (a dedicated 4 weeks of study and practice) of the initial training.

Anesthesiology societies have been repeatedly reminded of the importance of US education in contemporary anesthesia training.^{6,26} It is, therefore, foreseeable that FoCUS will become an integral component of anesthesiology training across Canada. Indeed, a recent survey revealed that FoCUS training is currently offered by 92% of Canadian anesthesiology training programs.²⁷ Overall, the goal would be to achieve Level I (basic) proficiency, for which guidelines have

Table 1 Demographic and descriptive information of anesthesiology residents in FoCUS ($n=11$).

Resident	Level of study (PGY level)	Pre ^a FoCUS self-rated knowledge (of 10)	Post FoCUS self-rated knowledge at EOR (of 10)	Post FoCUS self-rated knowledge at 6-month follow-up (of 10)	Number of scans completed during FoCUS rotation	Exam score at EOR (of 50)	Number of scans completed after EOR (6-month follow-up)	Exam score at 6-month follow-up (of 50)
1	4	2	8	6	64	44	10	44
2	3	2	9	7	80	41	2	41
3	3	0	9	7	57	47	3	42
4	4	4	8	6	85	47	2	46
5	3	1	7	6	65	41	n/a ^b	45
6	3	2	7.5	5	50	39	3	42
7	3	1	7.5	5	50	47	2	46
8	5	1	7	5	n/a ^b	44	5	42
9	4	2	9	5	52	44	4	40
10	4	1	9	5	53	47	5	46
11	4	1	8	4	50	44	6	39
Mean	4	1.5	8.0	5.5	60.6	44.1	4.2	43.0
SD		1.0	0.8	0.9	12.8	2.8	2.5	2.5

PGY, post-graduate year; FoCUS, focused cardiac ultrasound; EOR, end of rotation.

^a The pre FoCUS self-rated knowledge data were collected at the EOR.

^b n/a, Not available. The total number of scans was not reported by the resident on the respective time points. Hence, the mean scores and standard deviations (SD) for the columns ‘‘Number of scans completed during FoCUS rotation’’ and ‘‘Number of scans completed after EOR (6-month follow-up)’’ were calculated based on the 10 residents whose data were available.

already been proposed.^{28,29} However, unlike Emergency³⁰ and Critical Care Medicine,³¹ a formal FoCUS curriculum in Anesthesiology is still lacking, and the Royal College of Physicians and Surgeons of Canada has not yet taken the steps to establish FoCUS as part of the anesthesiology core training requirements.

In our cohort, residents who completed more scans during their FoCUS rotation had higher perceived knowledge at the 6-month follow-up. We speculate that a higher number of scans results in deeper learning, which in turn promotes self-efficacy (i.e., self-confidence). Furthermore, a premise of our FoCUS curriculum is that more scans equals more feedback. Feedback provided by expert staff on multiple occasions may promote experiential learning through enhancement of reflective observation (i.e., reviewing and reflecting on the scan and staff feedback) and abstract conceptualization (i.e., using feedback to learn from the experience).³² Finally, completing more scans may indicate highly intrinsically motivated individuals (although knowing they were going to be retested cannot be neglected as potentially influencing residents’ behavior during their FoCUS rotation). Intrinsic motivation has been favored in the educational literature as it is associated with self-initiated learning and deep learning.³³ Hence, feedback allows learners habituated in an intrinsically motivated learning approach to integrate new knowledge to already existing knowledge to solve clinical problems.³⁴

At present, different guidelines for FoCUS training have been proposed^{13,28,29,35}; however, they are limited by little evidence-based data. A subject of constant debate is the *amount* of training and clinical experience required to achieve competency.^{36,37} While studies have demonstrated that as little as a few hours of training allow participants

to reliably identify cardiac abnormalities,^{38,39} formal recommendations for proper credentialing are lacking. What seems clear, however, is that adequacy of training should be determined by competency-based assessment.⁴⁰ Establishing criteria for maintenance of certification is equally important as practitioners rapidly lose their skills without practice.⁴¹ Ideally, echocardiography societies in conjunction with specialty training committees would create a core set of curricular requirements and competencies pertaining to FoCUS training in anesthesiology postgraduate training programs.

Barriers to the implementation of a FoCUS rotation in anesthesiology training include time constraints, a lack of manpower and equipment, and limited expertise.⁴² As we faced all of these barriers at our institution, we endeavoured to implement from the outset a practical and sustainable training program with minimal resource utilization that is tailored to the reality of our institution. Key features to this program include the use of a commercially available online curriculum based on existing guidelines from the International Statement on Training Standards for Critical Care Ultrasonography,⁴³ and the self-directed nature of the rotation requiring residents to take initiative and ownership of their learning. The success of our curriculum, therefore, depends on highly motivated learners, which was never an issue given the perceived value of FoCUS training. Indeed, 91% of anesthesia residents in the United States agreed that FoCUS should be part of anesthesiology training.⁴² Furthermore, our curriculum has been constantly refined over the years in order to meet our local needs and requirements. Of note, given the small nature of our program (20 residents in total) and the great emphasis on self-directed learning, no more than 2 staff supervisors are required to adequately run




Figure 2 Wilcoxon signed rank test $p = 0.12$ (Video), $p = 0.7$ (Written), $p = 0.50$ (Total).

the FoCUS rotation. We therefore report on a self-directed FoCUS curriculum that has allowed many anesthesia trainees to achieve – and most importantly, to maintain – Level I competency over time.

This study has several limitations. First, residents identified their baseline self-perceived FoCUS knowledge following enrolment in the FoCUS curriculum rather than prior to enrolment. While pre-FoCUS self-perceived knowledge was lower than that at post-FoCUS rotation and at the 6-month follow-up, indicating that residents perceived that their participation in FoCUS training enhanced their knowledge, baseline data would have provided a comparison with the initial/baseline condition. Nevertheless, a previous study using similar assessments revealed that novices without FoCUS training perform poorly and rate their pre-FoCUS knowledge in the same range as residents in the current study (1.5 out of 10).¹³ Second, the small sample size may limit our ability to generalize our results to other populations. Additionally, the small nature of our program and the timeline of our study design precluded us from using a control group. Third, although our results showed similar exam scores at the EOR and 6 months later, the 6-month follow-up exam did not include a practical bedside FoCUS scan, and therefore, the trainees' technical ability to perform FoCUS was not assessed. Fourth, the FoCUS rotation in our institution was optional during our data collection. Therefore, a selection bias (including highly motivated individuals with a specific interest in FoCUS) may have positively influenced our results. Fifth, we did not have a follow-up beyond 6 months due to residents finishing their training. Finally, the multiple-choice examination we used at the EOR and at the 6-month assessment has not been formally validated to assess for a standard level of proficiency in echocardiography. However, we have previously shown that the exam used in this study is able to discriminate between trainees and experts, and that any improvement observed in exam scores after FoCUS training is not simply a result of re-testing.¹³

Summary

This pilot study suggests that 4 weeks of intensive FoCUS training consisting of a combination of didactic and practical

sessions results in adequate knowledge acquisition and 6-month knowledge retention. Further studies are required to determine if long-term retention is in fact related to clinical proficiency with FoCUS.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Jasudavicius A, Arellano R, Martin J, et al. A systematic review of transthoracic and transesophageal echocardiography in non-cardiac surgery: implications for point-of-care ultrasound education in the operating room. *Can J Anesth.* 2016;63:480–7.
- Barbariol F, Vetrugno L, Pompei L, et al. Point-of-care ultrasound of the diaphragm in a liver transplant patient with acute respiratory failure. *Crit Ultrasound J.* 2015;7:3.
- Leeson K, Leeson B. Pediatric ultrasound. Applications in the emergency department. *Emerg Med Clin North Am.* 2013;31:809–29.
- Ho AMH, Critchley LAH, Leung JYC, et al. Introducing final-year medical students to pocket-sized ultrasound imaging: teaching transthoracic echocardiography on a 2-week anesthesia rotation. *Teach Learn Med.* 2015;27:307–13.
- Nelson BP, Hojsak J, Dei Rossi E, et al. Seeing is believing: evaluating a point-of-care ultrasound curriculum for 1st-year medical students. *Teach Learn Med.* 2016;1334:1–8.
- Mizubuti GB, Allard RV, Tanzola RC, et al. Pro: focused cardiac ultrasound should be an integral component of anesthesia residency training. *J Cardiothorac Vasc Anesth.* 2015;29:1081–5.
- Kanji HD, McCallum J, Sironis D, et al. Limited echocardiography-guided therapy in subacute shock is associated with change in management and improved outcomes. *J Crit Care.* 2014;29:700–5.
- Augoustides JG, Hosalkar HH, Savino JS. Utility of transthoracic echocardiography in diagnosis and treatment of cardiogenic shock during noncardiac surgery. *J Clin Anesth.* 2005;17:488–9.
- Cowie B. Focused cardiovascular ultrasound performed by anesthesiologists in the perioperative period: feasible and alters patient management. *J Cardiothorac Vasc Anesth.* 2009;23:450–6.
- Ferguson EA, Paech MJ, Veltman MG. Hypertrophic cardiomyopathy and caesarean section: intraoperative use of transthoracic echocardiography. *Int J Obstet Anesth.* 2006;15:311–6.
- Gerlach RM, Saha TK, Allard RV, et al. Unrecognized tamponade diagnosed pre-induction by focused echocardiography. *Can J Anesth.* 2013;60:803–7.
- Oren-Grinberg A, Gulati G, Fuchs L, et al. Hand-held echocardiography in the management of cardiac arrest. *Anesth Analg.* 2012;115:1038–41.
- Tanzola RC, Walsh S, Hopman WM, et al. Brief report: Focused transthoracic echocardiography training in a cohort of Canadian anesthesia residents: a pilot study. *Can J Anesth.* 2013;60:32–7.
- Vignon P, Dugard A, Abraham J, et al. Focused training for goal-oriented hand-held echocardiography performed by non-cardiologist residents in the intensive care unit. *Intensive Care Med.* 2007;33:1795–9.
- Jones AE, Tayal VS, Kline JA. Focused training of emergency medicine residents in goal-directed echocardiography: a prospective study. *Acad Emerg Med.* 2003;10:1054–8.
- Semeraro F, Signore L, Cerchiari EL. Retention of CPR performance in anaesthetists. *Resuscitation.* 2006;68:101–8.

17. Beaulieu Y. Specific skill set and goals of focused echocardiography for critical care clinicians. *Crit Care Med.* 2007;35:S144–9.
18. Mayo PH, Beaulieu Y, Doelken P, et al. American College of Chest Physicians/La Société de Réanimation de Langue Française statement on competence in critical care ultrasonography. *Chest.* 2009;135:1050–60.
19. CAE Healthcare. CAE ICCU Web Portal. Available at: <https://www.caeiccu.com/> Accessed November 11, 2017.
20. Mitchell JD, Montealegre-Gallegos M, Mahmood F, et al. Multimodal perioperative ultrasound course for interns allows for enhanced acquisition and retention of skills and knowledge. *AA Case Rep.* 2015;5:119–23.
21. Town JA, Bergl PA, Narang A, et al. Internal medicine residents' retention of knowledge and skills in bedside ultrasound. *J Grad Med Educ.* 2016;8:553–7.
22. Bell DS, Harless CE, Higa JK, et al. Knowledge retention after an online tutorial: a randomized educational experiment among resident physicians. *J Gen Intern Med.* 2008;23:1164–71.
23. Raman M, McLaughlin K, Violato C, et al. Teaching in small portions dispersed over time enhances long-term knowledge retention. *Med Teach.* 2010;32:250–5.
24. Noble VE, Nelson BP, Sutingco AN, et al. Assessment of knowledge retention and the value of proctored ultrasound exams after the introduction of an emergency ultrasound curriculum. *BMC Med Educ.* 2007;7:40.
25. Custers EJFM. Long-term retention of basic science knowledge: a review study. *Adv Health Sci Educ Theory Pract.* 2010;15:109–28.
26. Mahmood F, Matyal R, Skubas N, et al. Perioperative ultrasound training in anesthesiology. *Anesth Analg.* 2016;122:1794–804.
27. Mizubuti G, Allard R, Ho AMH, et al. A survey of focused cardiac ultrasonography training in Canadian anesthesiology residency programs. *Can J Anesth.* 2017;64:441–2.
28. Burwash IG, Basmadjian A, Bewick D, et al. 2010 Canadian Cardiovascular Society/Canadian Society of Echocardiography guidelines for training and maintenance of competency in adult echocardiography. *Can J Cardiol.* 2011;27:862–4.
29. Royse CF, Carty DJ, Faris J, et al. Core review: Physician-performed ultrasound: the time has come for routine use in acute care medicine. *Anesth Analg.* 2012;115:1007–28.
30. Akhtar S, Theodoro D, Gaspari R, et al. Resident training in emergency ultrasound: consensus recommendations from the 2008 council of emergency medicine residency directors conference. *Acad Emerg Med.* 2009;16:32–6.
31. Mazraeshahi RM, Farmer JC, Porembka DT. A suggested curriculum in echocardiography for critical care physicians. *Crit Care Med.* 2007;35:S431–3.
32. Kolb DA. Experiential learning: experience as the source of learning and development. Prentice Hall Inc.; 1984. p. 20–38.
33. Entwistle N. Motivational factors in students' approaches to learning. In: Schmeck RR, editor. *Learning Strategies and Learning Styles.* Springer; 1988. p. 21–51.
34. Kusurkar RA, Croiset G, Mann KV, et al. Have motivation theories guided the development and reform of medical education curricula? A review of the literature. *Acad Med Acad Med.* 2012;87:735–43.
35. Sicari R, Galderisi M, Voigt JU, et al. The use of pocket-size imaging devices: a position statement of the European Association of Echocardiography. *Eur J Echocardiogr.* 2011;12:85–7.
36. Cowie B, Kluger R. Evaluation of systolic murmurs using transthoracic echocardiography by anaesthetic trainees. *Anesthesia.* 2011;66:785–90.
37. Lau G, Swanevelder J. Echocardiography in intensive care: where we are heading? *Anesthesia.* 2011;66:649–52.
38. Spurney CF, Sable CA, Berger JT, et al. Use of a hand-carried ultrasound device by critical care physicians for the diagnosis of pericardial effusions, decreased cardiac function, and left ventricular enlargement in pediatric patients. *J Am Soc Echocardiogr.* 2005;18:313–9.
39. Kobal SL, Trento L, Bahrami S, et al. Comparison of effectiveness of hand-carried ultrasound to bedside cardiovascular physical examination. *Am J Cardiol.* 2005;96:6–1002.
40. Melniker L. International evidence-based recommendations for focused cardiac ultrasound. *J Am Soc Echocardiogr.* 2014;27, e1-683.e33.
41. Kimura BJ, Sliman SM, Waalen J, et al. Retention of ultrasound skills and training in "Point-of-Care" cardiac ultrasound. *J Am Soc Echocardiogr.* 2016;29:992–7.
42. Conlin F, Roy Connelly N, Raghunathan K, et al. Focused transthoracic cardiac ultrasound: a survey of training practices. *J Cardiothorac Vasc Anesth.* 2016;30:102–6.
43. Cholley BP. International expert statement on training standards for critical care ultrasonography. *Intensive Care Med.* 2011;37:1077–83.