

Changes in functional capacity of patients two years after coronary artery bypass grafting surgery

Alterações na capacidade funcional de pacientes após dois anos da cirurgia de revascularização do miocárdio

Rosane Maria NERY¹, Marcio Roberto MARTINI², Cristiane da Rocha VIDOR³, Mahmud Ismail MAHMUD⁴, Maurice ZANINI⁵, Aderson LOUREIRO⁶, Juarez Neuhaus BARBISAN⁷

RBCCV 44205-1177

Abstract

Objective: To check changes in the functional capacity of patients undergoing coronary artery bypass grafting (CABG) through the six-minute walk test (6MWT) in a 2-year follow-up.

Methods: In this prospective cohort study in 215 patients who underwent CABG, 13 did not meet the criteria for inclusion. Of these, 202 patients were evaluated preoperatively, 13 died during hospitalization and 6 in the 2-year follow-up. Four patients were considered lost-to-follow-up. This study followed 179 patients for two years classified into active and sedentary, according to physical activity in leisure time and to the 6MWT preoperatively and 2 years later.

Results: One hundred and twenty patients were evaluated in the day before CABGS, being 67% male with an average age of 63 (± 9.75) years. Before surgery and 2 years later, 52 were active and the 6MWT distances performed had been 359m (± 164.47) and 439m (± 171.34), respectively; $P = 0.016$. Forty five patients were classified as sedentary before and 2 years after surgery. The 6MWT distances walked before and after surgery had been 255m (± 172.15) and 376m (± 210.92), respectively; $P < 0.001$. Eighty two patients transited between these two groups, 71 passed from sedentary

to active and had walked before and after surgery 289m (± 157.15) and 380m (± 125.44), respectively; $P = 0.001$. The 11 patients who were active and passed to the sedentary group walked 221m (± 191.91) and 384m (± 63.73), respectively; $P = 0.007$.

Conclusion: The functional capacity of the patients who underwent CABGS had a significant improvement in a medium-term follow-up.

Descriptors: Myocardial revascularization. Functional residual capacity. Total lung capacity. Walking.

Resumo

Objetivo: Verificar alterações na capacidade funcional dos pacientes que se submetem à cirurgia de revascularização do miocárdio (CRM) por meio do teste de caminhada de seis minutos (TC6) no seguimento de dois anos.

Métodos: Estudo de coorte prospectivo, no qual foram arrolados 215 pacientes submetidos a CRM, 13 não preencheram os critérios de inclusão. Foram 202 pacientes avaliados no pré-operatório, 13 morreram no período da internação e seis no seguimento de dois anos. Quatro pacientes foram considerados perdidos de seguimento. Foram

1. Master's Degree in Health Science, Cardiology; Physical Educator, HCPA.
2. Undergraduate Health Science Student, Cardiology; Physical Educator, HCPA.
3. Graduated in Physical Education and Training; Physical Educator, UNISINOS.
4. Master's Degree in Epidemiology; Physical Therapist, HCPA.
5. Graduated in Physical Education and Training; Physical Educator, UFRGS.
6. Master's Degree in Sciences of Human Movement (Kinesiology); Physical Educator, UNISINOS.
7. M. D., Cardiology; Cardiologist in charge of the Tilt Test Sector - Instituto de Cardiologia do Rio Grande do Sul/ FUC.

This study was carried out at Instituto de Cardiologia do Rio Grande do Sul/ Fundação Universitária de Cardiologia, Porto Alegre, RS, Brazil.

Correspondence address:

Rosane Maria Nery - Caixa Postal 5032, Bom Fim - Porto Alegre, RS, Brasil. CEP: 90041-970

E-mail: barbisan.pesquisa@cardiologia.org.br

Article received on: March 25th, 2010

Article accepted on: May 18th, 2010

acompanhados 179 pacientes no período de 2 anos, classificados em ativos e sedentários, conforme a prática de atividade física no tempo livre e submetidos ao TC6 no pré-operatório e 2 anos depois.

Resultados: Dos 202 pacientes avaliados no pré-operatório da CRM, 67% eram do sexo masculino, com idade média de 63 ($\pm 9,75$) anos. Pré e após 2 anos da CRM, 52 pacientes se mantiveram ativos e as distâncias caminhadas foram 359m ($\pm 164,47$) e 439m ($\pm 171,34$), respectivamente, $P= 0,016$. A distância caminhada no pré e pós-operatório, dos 45 pacientes que permaneceram sedentários, foi, respectivamente, 255m ($\pm 172,15$) e 376m ($\pm 210,92$) $P < 0,001$. Oitenta e dois pacientes

transitaram entre estes dois grupos, 71 passaram de sedentários para ativos e caminharam 289m ($\pm 157,15$) no pré e 380m ($\pm 125,44$) no pós-operatório, $P= 0,001$; os 11 pacientes que eram ativos e passaram a sedentários caminharam no pré 221m ($\pm 191,91$) e, no pós-operatório, 384m ($\pm 63,73$) $P= 0,007$.

Conclusão: A capacidade funcional dos pacientes submetidos à CRM melhorou de forma importante no seguimento de médio prazo.

Descritores: Revascularização miocárdica. Capacidade residual funcional. Capacidade pulmonar total. Caminhada.

INTRODUCTION

The coronary artery bypass grafting (CABG) has been proposed to minimize symptoms, improve cardiac function, survival and reduce the recurrence of major cardiac adverse events in selected subgroups of patients [1]. Studies have suggested that physical activity is beneficial in both the prevention and treatment of patients with ischemic heart disease, and it should be recommended for those undergoing CABG [2]. Therefore, the maintenance of physical activity in operated patients should be recommended. A controlled clinical trial evaluated the cardiopulmonary rehabilitation in CABG patients, which consisted of muscular exercises and ventilation, as well as preoperative education for at least 5 days and postoperatively until discharge. A sharp reduction in pulmonary complications, a significant difference in the incidence of cardiac arrhythmias, and an improved functional capacity measured on the seventh day after surgery through the six-minute walk test (6MWT) were observed [3].

In a controlled study described by Stein et al. [4], a group attended exercise-based cardiac rehabilitation, carried out bronchial hygiene, and used EPAP mask within seven days of hospitalization after CABG and another group received usual care only. At the time of hospital discharge, the patients in the rehabilitation group have walked a significantly greater distance in the 6MWT.

Physical capacity can be assessed by different methods. The cardiopulmonary exercise test is the gold standard. However, it has some limitations including a relatively high cost and reduced applicability in patients with physical deficits, and it does not reproduce the activities of daily

living [5]. The 6MWT was well established to evaluate the functional capacity, especially in elderly patients with comorbidities [6]. The 6MWT is a valid instrument to assess the progression of functional capacity for exercise in different clinical interventions. It presents easy to use, and it is inexpensive. Even as a submaximal exercise test it can identify patients with poor prognosis and help in making the most appropriate therapeutic measures [7,8].

Therefore, our objective was to assess changes in the functional capacity of patients classified into active and sedentary who undergo CABG using the 6MWT after two years.

METHODS

A prospective, multicenter, cohort study to evaluate 215 consecutive patients undergoing elective CABG was carried out. Of these patients, 13 did not meet the inclusion criteria.

Inclusion and Exclusion criteria

The study included patients referred for elective CABG, in stable clinical condition at the time of preoperative evaluation performed by the responsible staff in each institution. The patients were considered fitted to take the 6MWT. The main reasons for exclusion were patients operated on an emergency basis with acute coronary syndrome and concomitant congenital heart disease or terminal illnesses, and patients with unstable angina and musculoskeletal impairment, which would make it impossible for them to take the 6MWT.

Ethical Aspects

This study was approved by the ethical committee and

scientific committee of Instituto de Cardiologia do Rio Grande do Sul (IC/RS), Hospital de Clínicas de Porto Alegre (HCPA) e Santa Casa de Misericórdia de Porto Alegre (SCMPOA), and a free and informed consent was obtained from each patient.

Instruments

A structured questionnaire was applied and the information on demographic, anthropometric and clinical data was examined.

Patients were divided into two groups, according to leisure-time physical activity (LTPA): Group I - active patients who performed physical activities in their free time three or more times per week and for 30 minutes or more over the past two weeks before surgery and Group II – the sedentary patients.

In order to give more consistency to this classification, it was applied the Baecke Questionnaire of Habitual Physical Activity (BQHPA), which investigates the physical activity of the last 12 months [9]. The 6MWT was performed in the corridors of the hospital, in a level ground, with the distance previously marked. Patients were monitored for blood pressure (BP) and heart rate (HR) at rest and immediately after the test; the same was done as a pattern of the American Thoracic Society (ATS) [10].

Statistical Analysis

Statistical analysis was performed using the software *Statistical Package for Social Sciences* (SPSS, version 15.0). Categorical variables were presented as absolute frequencies and percentages. Continuous variables were expressed as mean ± standard deviation when normally distributed. Demographic characteristics and comparison between groups of active and sedentary patients were analyzed by Student's *t*-test for continuous variables and the chi-square test for categorical variables. A *P* < 0.05 was considered statistically significant.

RESULTS

Of the 202 patients evaluated preoperatively, 13 died during hospitalization and six after two years. We were unable to contact four patients; they were considered lost-to-follow-up; 179 patients were tracked over a 2-year follow-up.

In Table 1 are shown the demographic, anthropometric, and clinical variables. The study group showed a majority of male patients, 120 (67%). The mean age of the patients was 63 (± 9.75) years. Fifty-seven patients started physical activity two years after CABG.

Table 2 shows the distance walked during the preoperative period and in the 2-year follow-up of patients who remained active and those who remained sedentary. Both groups showed a significant increase in distance walked during the study period.

Table 3 shows those patients who went from being active preoperatively to be sedentary after two years. They showed no difference in walking distance; however, patients who went from being sedentary to active, over the

Table 1. Characteristics of patients undergoing CABG in the preoperative and postoperative period classified into active and sedentary.

Characteristics	Actives n = 66	Sedentaries n = 136	P Value
Age, mean age (±SD), years	60(± 10)	62 (± 10)	0.14
Male, n%	51 (77.3)	83 (61)	0.02
BMI, mean (±SD)	27 (± 4)	27 (± 5)	0.97
WHR, mean (±SD)	0.96 (± 0.07)	0.96 (± 0.11)	0.99
Ejection fraction, mean (±SD)	61(± 14)	60 (± 13)	0.70
Smoking, n%	42 (63.6)	87 (64)	0.95
Diabetes mellitus, n%	21 (31.8)	51 (37.5)	0.43
SAH, n%	58 (87.9)	119 (87.5)	0.94
Previous AMI, n%	20 (30.3)	36 (26.5)	0.57
PVD, n%	8 (12.1)	20 (14.7)	0.62
COPD, n%	4 (6.1)	9 (6.6)	1.00
Dyslipidemia, n%	14 (21.2)	38 (27.9)	0.31

BMI= Body mass index; WHR= Waist-hip ratio; SAH= Systemic arterial hypertension; PVD= Peripheral vascular disease; COPD= Chronic obstructive pulmonary disease

Table 2. Distance (in meters) walked by the groups in the preoperative and postoperative period that kept their characteristics of active and sedentary.

Patients	Preoperative walked distance	Postoperative walked distance	P Value
Actives (n=52)*	358.52 (±164.47)	439.43 (±171.34)	0.016
Sedentaries (n=45)*	254.56 (±172.15)	375.53 (±210.92)	<0.001

* Data presented as mean (±SD)

Table 3. Distance (in meters) walked by both groups active and sedentary in the preoperative and postoperative period that had changed their habits.

Patients	Preoperative walked distance	Postoperative walked distance	P value
Sedentaries to	289.03	379.62	0.001
Actives (n=71)*	(±157.15)	(±125.44)	
Actives to	221.36	384.41	0.21
Sedentaries (n=11)*	(±191.91)	(±63.73)	

* Data presented as mean (±SD)

same period, show a significant difference in walking distanced.

DISCUSSION

Our main finding shows that in a population of patients undergoing CABG, those who remained active in the period of two years and those who went from being sedentary to active improve their functional capacity.

Nery et al. [11] showed the importance of physical activity in the preoperative period in the outcome of CABG. The physically active patients had shorter hospital stays and fewer trans- and postoperative complications within a year. The cardiac surgery promoted a change in patients' lifestyle, increasing the number of physically active individuals in a 1-year follow-up.

Traditionally, exercise tolerance has been used as an indicator of overall physical capacity. It has also been demonstrated that exercise intolerance predicts a poor physical ability to perform tasks of daily living [12]. Besides, regular physical exercise produces a variety of beneficial effects on health, such as increased longevity, decreased mortality and incidence of metabolic disorders, among other comorbidity [13].

Kavanagh et al. [14] evaluated the influence of a change in both physical fitness and physical activity after a year of a walking-based cardiac rehabilitation program; the influence on cardiac mortality, and mortality from other causes. Patients were followed-up over nine years. The distance walked increased significantly during the training period, constituting an important predictor of cardiac mortality, in which each increase in every mile walked corresponded up to 20% reduction in mortality.

The effects of different interventions conducted only during the postoperative period of CABG were approached by some studies. Among them, Hirschhorn et al. [15] evaluated 93 patients who were randomized into three groups. Among the results, only the groups who were walking with or without breathing exercises, increased performance on the 6MWT distance at the time of discharge.

Herdy et al. [3], in a controlled clinical trial have evaluated the cardiopulmonary rehabilitation which consisted of muscular exercise and ventilation, as well as preoperative and postoperative education for at least 5 days until discharge. As a result, the authors could observe an expressive reduction in pulmonary complications, a significant difference in the incidence of cardiac arrhythmias, and an improved functional capacity measured on the seventh day after surgery using the 6MWT.

CONCLUSION

Physically active postoperatively patients improved their

walking distance and functional capacity. Patients who stopped their physical activities showed no change. As a suggestion to the patients after CABG, in cardiac rehabilitation phase II, we consider important to recommend sessions of walking as part of a healthy lifestyle.

REFERENCES

1. Cutlip D, Levin T, Aroesty J. Bypass surgery versus percutaneous intervention in the management of stable angina pectoris: Recommendations. Disponível em: <http://www.uptodate.com> Acesso em: 15/04/2009.
2. Nery RM. Valor prognóstico da atividade física no pós-operatório da cirurgia de revascularização do miocárdio [Dissertação de Mestrado]. Porto Alegre: Instituto de Cardiologia; Programa de Pós-Graduação em Ciências da Saúde, Área de Concentração: Ciências Cardiovascular;2007.
3. Herdy AH, Marcchi PL, Vila A, Tavares C, Collaço J, Niebauer J, et al. Pre- and postoperative cardiopulmonary rehabilitation in hospitalized patients undergoing coronary artery bypass surgery: a randomized controlled trial. *Am J Phys Med Rehabil*. 2008;87(9):714-9.
4. Stein R, Maia CP, Silveira AD, Chiappa GR, Myers J, Ribeiro JP. Inspiratory muscle strength as a determinant of functional capacity early after coronary artery bypass graft surgery. *Arch Phys Med Rehabil*. 2009;90(10):1685-91.
5. Osada N, Chaitman BR, Miller LW, Yip D, Cishek MB, Wolford TL, et al. Cardiopulmonary exercise testing identifies low risk patients with heart failure and severely impaired exercise capacity considered for heart transplantation. *J Am Coll Cardiol*. 1998;31(3):577-82.
6. De Feo S, Tramarin R, Lorusso R, Faggiano P. Six-minute walking test after cardiac surgery: instructions for an appropriate use. *Eur J Cardiovasc Prev Rehabil*. 2009;16(2):144-9.
7. Enright PL, McBurnie MA, Bittner V, Tracy RP, McNamee R, Arnold A, et al. The 6-min walk test: a quick measure of functional status in elderly adults. *Chest*. 2003;123(2):387-98.
8. Rodrigues SL, Mendes HF, Viegas CAA. Teste de caminhada de seis minutos: estudo do efeito do aprendizado em portadores de doença pulmonar obstrutiva crônica. *J Bras Pneumol*. 2004;30(2):121-5.

-
9. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr.* 1982;36(5):936-42.
 10. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Resp Crit Care Med.* 2002;166(1):111-7.
 11. Nery RM, Barbisan JN, Mahmud MI. Influência da prática da atividade física no resultado da cirurgia de revascularização miocárdica. *Rev Bras Cir Cardiovasc* 2007;22(3):297-302.
 12. Calsamiglia G, Camera F, Mazza A, Villa P, Gigli Berzolari F, Tramarin R, et al. A new test (VITTORIO Test) for functional fitness assessment in rehabilitation after cardiac surgery. *Monaldi Arch Chest Dis.* 2005;64(1):8-18.
 13. Lippi G, Maffulli N. Biological influence of physical exercise on hemostasis. *Semin Thromb Hemost.* 2009;35(3):269-76.
 14. Kavanagh T, Hamm LF, Beyene J, Mertens DJ, Kennedy J, Campbell R, et al. Usefulness of improvement in walking distance versus peak oxygen uptake in predicting prognosis after myocardial infarction and/or coronary artery bypass grafting in men. *Am J Cardiol.* 2008;101(10):1423-7.
 15. Hirschhorn AD, Richards D, Mungovan SF, Morris NR, Adams L. Supervised moderate intensity exercise improves distance walked at hospital discharge following coronary artery bypass graft surgery. *Heart, Lung Circ.* 2008;17(2):129-38.