



# Revista Brasileira de Farmacognosia

BRAZILIAN JOURNAL OF PHARMACOGNOSY

www.journals.elsevier.com/revista-brasileira-de-farmacognosia



## Original article

# Halistanol sulfate A and rodriguesines A and B are antimicrobial and antibiofilm agents against the cariogenic bacterium *Streptococcus mutans*

Bruna de A. Lima<sup>a,b,\*</sup>, Simone P. de Lira<sup>c</sup>, Miriam H. Kossuga<sup>d</sup>, Reginaldo B. Gonçalves<sup>e</sup>, Roberto G.S. Berlinck<sup>d</sup>, Regianne U. Kamiya<sup>f</sup>

<sup>a</sup>Faculdade de Odontologia de Piracicaba, Universidade Estadual de Campinas, Piracicaba, SP, Brazil

<sup>b</sup>Instituto de Biologia, Universidade Estadual de Campinas, Campinas, SP, Brazil

<sup>c</sup>Escola Superior de Agronomia Luiz de Queiroz, Universidade de São Paulo, Piracicaba, SP, Brazil

<sup>d</sup>Instituto de Química de São Carlos, Universidade de São Paulo, São Carlos, SP, Brazil

<sup>e</sup>Faculté de Médecine Dentaire, Université Laval, Québec, Canada

<sup>f</sup>Instituto de Ciências Biológicas e da Saúde, Universidade Federal de Alagoas, Maceió, AL, Brazil

## ARTICLE INFO

### Article history:

Received 17 June 2014

Accepted 17 November 2014

### Keywords:

Biofilms

Molecular oral microbiology

Dental caries

*Streptococcus*

Marine natural products

## ABSTRACT

In the present investigation we report the antibacterial activity of halistanol sulfate A isolated from the sponge *Petromica ciocalyptoides*, as well as of rodriguesines A and B isolated from the ascidian *Didemnum* sp., against the caries etiologic agent *Streptococcus mutans*. The transcription levels of *S. mutans* virulence genes *gtfB*, *gtfC* and *gpbB*, as well as of house-keeping genes *groEL* and *16S*, were evaluated by sqRT-PCR analysis of *S. mutans* planktonic cells. There were no alterations in the expression levels of *groEL* and *16S* after antimicrobial treatment with halistanol sulfate A and with rodriguesines A and B, but the expression of the genes *gtfB*, *gtfC* and *gpbB* was down-regulated. Halistanol sulfate A displayed the most potent antimicrobial effect against *S. mutans*, with inhibition of biofilm formation and reduction of biofilm-associated gene expression in planktonic cells. Halistanol sulfate A also inhibited the initial oral bacteria colonizers, such as *Streptococcus sanguinis*, but at much higher concentrations. The results obtained indicate that halistanol sulfate A may be considered a potential scaffold for drug development in *Streptococcus mutans* antibiofilm therapy, the main etiologic agent of human dental caries.

© 2014 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. All rights reserved.

## Introduction

Caries is an infectious disease caused by the accumulation of biofilm on the tooth surface. *Streptococcus mutans* is considered the main etiologic agent of human dental caries (Whiley and

Beighton, 1998). *S. mutans* can metabolize diet carbohydrates, producing high levels of water-insoluble glucan matrices from sucrose, which are responsible for the initial stage of oral biofilm formation. *S. mutans* expresses several virulence genes that regulate its growth and accumulation on tooth surfaces,

\* Corresponding author.

E-mail: b069508@dac.unicamp.br (B.A. Lima).

promoting acid-mediated tooth demineralization and acid tolerance. *S. mutans* is adapted to the biofilm lifestyle, which is essential for the pathogenesis of dental caries (Stipp et al., 2008).

One of the strategies for preventing the formation and development of dental caries is to control the growth and adherence of mutans streptococci on the dental surface (Smith, 2002) by inhibiting cell-cell signaling (Lonn-Stensrud et al., 2007) and/or acid production. Several *S. mutans* virulence genes and corresponding functions have been identified. For example, glucosyltransferases GtfB and GtfC catalyze the extracellular synthesis of a water-insoluble glucan matrix from sucrose, and are essential for the accumulation of bacteria in the dental biofilm. GbpB (*Glucan Binding Proteins*), an essential protein of *S. mutans*, might also mediate cell surface interaction with glucan, increasing the adherence ability of the bacterial pathogen in the dental biofilm (Mattos-Graner et al., 2001). Inhibitors of these virulence factors represent important candidates for antibiofilm therapy.

Fluoride, chlorhexidine and their combinations have been used to control dental caries (Bader et al., 2001). Additionally, natural product-based treatments, such as propolis and plant extracts, have been successfully used against dental caries (Koo et al., 2000, 2006; Galvão et al., 2012). Examples of anticariogenic natural products include anthraquinones, apigenin, tt-farnesol, chitosan, 7-epiclusianone, 2-aminoimidazole and 2-aminobenzimidazole, several of which have shown antibiofilm activity against *Streptococcus mutans* (Koo et al., 2003; Coenye et al., 2007; Murata et al., 2008; Pasquantonio et al., 2008; Liu et al., 2011). However, very few natural products isolated from marine organisms have been evaluated as inhibitors of bacterial oral pathogens.

Marine natural products isolated from invertebrates, such as sponges, ascidians, octocorals and bryozoans, as well as from associated microorganisms, such as bacteria, fungi, cyanobacteria and dinoflagellates, are considered very promising sources of new drug candidates, including antibiotics (Selegim et al., 2007; Liu et al., 2011; Mai et al., 2011; Kossuga et al., 2012; Mayer et al., 2013; Butler et al., 2014; Blunt et al., 2014).


The aim of this investigation was to evaluate the antimicrobial activity of halistanol sulfate A and of rodriguesines A and B against three species of oral bacterial and three species of pathogens. The same compounds were further evaluated for their effect on the transcription levels of known biofilm-associated genes, including *gtfB*, *gtfC* and *gbpB*, of two *Streptococcus mutans* strains. In addition, the transcription levels of *groEL* and 16S genes were also evaluated on *Streptococcus mutans*.

## Material and methods

### Isolation of halistanol sulfate A and of rodriguesines A and B

Halistanol sulfate A (**1**) was obtained from the sponge *Petromica ciocalyptoides* as previously described (Kossuga et al., 2007). Rodriguesines A (**2**) and B (**3**) were obtained from an ascidian of the genus *Didemnum* as an inseparable mixture of homologues,

as previously described (Kossuga et al., 2009) and tested as such. Both halistanol sulfate A and rodriguesines A and B have been identified by a complete analysis of spectroscopic data, as previously reported (Kossuga et al., 2007, Kossuga et al., 2009).



### Microorganisms

Pure compounds obtained from marine organisms were tested against *Streptococcus mutans* UA 159, *Streptococcus mutans* CI (clinical isolate), *Streptococcus sanguinis* ATCC 15300, *Streptococcus sobrinus* ATCC 27607, *Staphylococcus aureus* ATCC 6538, *Enterococcus faecalis* ATCC 14506 and *Escherichia coli* NCTC 8196. *Streptococcus mutans* CI was isolated from an active caries individual, as previously described (Klein et al., 2004).

### Antimicrobial assays

The minimal inhibitory concentration (MIC) test or broth microdilution assay was carried out using microplates (96 wells), following the procedure described by Oliveira et al. (2006), with modifications. Aliquots of pure compounds were diluted in 1% (v/v) dimethylsulfoxide (DMSO - 1250  $\mu\text{g}/\text{ml}$ ). Serial dilutions of antimicrobial agents were prepared in 80  $\mu\text{l}$  of BHI (Brain Heart Infusion) medium, and 20  $\mu\text{l}$  of bacterial suspension (approximately  $1 \times 10^8$  CFU/ml) were added to the wells containing different final concentrations of the antimicrobial agents ranging from 0.05 to 500  $\mu\text{g}/\text{ml}$ , at a final volume of 100  $\mu\text{l}/\text{well}$ . Microplates were incubated at 37°C and 10%  $\text{CO}_2$  for 24 h. The bacterial growth was measured by absorbance at  $\text{OD}_{550-560}$  nm in an automated microplate reader (Molecular Devices, Versa Max Program). The MIC values were defined as the lowest antimicrobial concentration that inhibited 90 to 100% of bacterial growth compared to a positive control. Microbial cultures grown in medium (1% DMSO v/v) without antimicrobial samples and with doxycycline (0.05 to 5  $\mu\text{g}/\text{ml}$ ) were used as the negative and positive controls, respectively. All samples were assayed in triplicate.

To determine the minimal bactericidal concentration (MBC) of the tested samples, an aliquot of each sample in

concentrations  $\geq$  MIC was inoculated on BHA medium. The plates were incubated at 37°C and 10% CO<sub>2</sub> for 24 to 48 h. The MBC values were defined as the lowest concentration of a pure compound that promoted no visible bacterial growth in the tested medium. To determine the nature of the antibacterial effect of the samples, the bacterial MBC/MIC ratio was calculated. When the MBC/MIC ratio was between 1 and 2, the extract or pure compound was considered as bactericidal against the tested pathogen. When the MBC/MIC ratio was higher than 2, the active extract or pure compound was considered as bacteriostatic (Galvão et al., 2012).

### Inhibitory activity against biofilm formation

The selected microorganisms were *Streptococcus mutans* UA 159 and *Streptococcus mutans* CI. The samples of halistanol sulfate A (**1**) and of rodriguesines A (**2**) and B (**3**) that displayed broad-spectrum antimicrobial activity against cariogenic species were selected based on data obtained after the MIC and MBC testing. Serial dilutions of the samples were added to wells of sterile polystyrene U-bottom microtiter plates. Final concentrations between 0.05 to 62.5 µg/ml of the antimicrobial agents in 0.5% (w/v) sucrose-containing THB medium (Toddy Hewitt Broth) were separately added to inocula (approximately 10<sup>8</sup> CFU/ml) of *Streptococcus mutans* UA 159 or *Streptococcus mutans* CI. The microplates were incubated in 10% CO<sub>2</sub> at 37°C for 18 h. Biofilm growth was examined and quantified using the crystal violet staining method by measuring the absorbance at OD<sub>575</sub> nm (O'Toole and Kolter, 1998). Briefly, after 18 h of incubation, the spent medium was aspirated, non-adhering cells were removed, the wells were washed three times with sterile distilled water, and the plates were dried for 45 min. The wells were stained with crystal violet at 1% (w/v) for 30 min, washed three times with distilled water and destained with 98% ethanol (v/v) before carrying out the biofilm quantification in an automated microplate reader (Molecular Devices, Versa Max Program). Doxycycline in concentrations between 0.05 and 1.0 µg/ml was used as a positive control. DMSO 1% (v/v) was used as a negative control.

The MIC of biofilm inhibition was determined as the lowest antimicrobial concentration capable of inhibiting biofilm formation without interfering with planktonic growth. Tests were performed in quadruplicate. Planktonic cultures in the wells containing biofilms at the MIC of biofilm inhibition, as well as 1% (v/v) DMSO vehicle, were serially diluted and plated on BHA (Brain Heart Agar). The number of viable cells (in CFU/ml) at the MIC of biofilm inhibition, relative to control, was measured to confirm the minimal interference of the antimicrobial activity on planktonic growth.

### RNA extraction, synthesis of cDNA and semi-quantitative RT-PCR

RNA extraction of planktonic cell cultures was performed. Growth curves of two *Streptococcus mutans* strains were determined to establish the middle log phase (D.O.  $\approx$  0.5). Briefly, frozen *Streptococcus mutans* cultures were reactivated in 5 ml TSB (Trypticase Soy Broth) and incubated at 37°C in 10% CO<sub>2</sub> for 10 to 12 h. Standard inocula of the strains

(approximately 10<sup>8</sup> CFU/ml, at OD<sub>550</sub> nm) were transferred to tubes containing 5 ml of fresh pre-warmed BHI medium and incubated in the same conditions described above for the determination of the growth curves. The optical density (O.D.) was measured spectrophotometrically at OD<sub>550-560</sub> nm every 1-2 h. Bacterial cell cultures in the middle log phase (O.D.  $\approx$  0.5) were submitted to pure compounds, doxycycline or 1% (v/v) DMSO control. Antimicrobial samples were evaluated for biofilm inhibition for 1 min (named T1) and 30 min (named T30) (Koo et al., 2006) with treated cells, which were collected by centrifugation (10,000 x g for 2 min, at 2 °C) and then frozen at -70 °C for RNA extraction. The experiment was conducted in duplicate, using two distinct cultures for each strain. RNA extraction of the bacterial cells was achieved using a Qiagen RNeasy kit according to the manufacturer's protocol. Cells were lysed by incubating them with lysozyme (20 µg/ml) in a water bath at 37°C for 30 min and vortexing the sample in 5 min intervals (Duque et al., 2011). The residual DNA was removed with deoxyribonuclease I according to the manufacturer's instructions. cDNA synthesis was performed with 24 ng of total RNA using a random primer Mix (20 µM) Ea1, Ea7, Es1, Es3 and Es8 (Chia et al., 2001) and the enzyme Super Transcript RT III according to the manufacturer's protocol, with some previously described modifications (Kamiya et al., 2008).

The PCR experiment was performed in triplicate for each cDNA. One microliter of cDNA (obtained from 24 ng of RNA) was used as a template in the PCR (25 µl) analysis, with 50 mM of MgCl<sub>2</sub>, 0.3 mM of each specific primer, 200 µM of dNTPs, 1.25 U of Taq polymerase and 10x PCR buffer, to generate amplicons ranging from 150-200 base pairs (Table 1). The amplicons were visualized on agarose gels (2% w/v) with ethidium bromide (0.3 µg/ml). Positive controls and a 100 bp DNA ladder were included in each gel. Digital images were captured under ultraviolet light using the Gel logic 100 Imaging System (Eastman Kodak Co.). The curves of the amplicons per number of thermal cycles (1-60 cycles), specific for each primer pair, were then generated to identify the middle of the exponential phase amplification or to establish the optimal number of cycles for each target gene. The thermal cycling parameters were as follows: 94°C for 45 s, annealing temperature (Table 1) for 60 s, and 72°C for 60 s.

All cDNA samples were tested with their respective primers for a specific target gene as well as with primers for the housekeeping gene 16S, a constitutively transcribed control gene whose expression is invariant under the used experimental conditions (Stipp et al., 2008). The intensities of the amplicons captured in the gels, using the same imaging system, were expressed as the relative arbitrary units of expression (UR). Controls for the RT-PCR included: reaction mixtures lacking template cDNA to effectively rule out (detect) the presence of contaminating DNA, and/or the formation of primer dimers. The positive controls included genomic DNA of the respective pathogenic bacterial strain.

### Statistical analysis

Student's t test was used to compare the significant differences between the planktonic growth or biofilm formation of cells submitted to the antimicrobial activity and the control group. The levels of genetic transcription variation of the treated cells

**Table 1**

Primer sequence, Gene Bank reference, amplicon size, and thermal conditions applied in the semi-quantitative reverse transcription polymerase chain reactions of known biofilm-associated genes and house-keeping gene 16S.

Primers	Gene Bank	Primers Sequences (5' → 3')	Product Size	Number of Cycles	Annealing
<i>gfpB</i> _Forward	1029610	TTAATACGATTCAAGGACAA	121pb	35	52
<i>gfpB</i> _Reverse		CTTGAAAGTGTGTTGAATTTG			
<i>gtfB</i> _Forward	1028336	AGATGGAACCTTACGCTTATT	150 pb	35	52
<i>gtfB</i> _Reverse		TCATCAAAGTATTGAACCTG			
<i>gtfC</i> _Forward	1028343	TGATTAACATGGATAACAGG	102 pb	35	52
<i>gtfC</i> _Reverse		CCAAACTGTTAGTGATCAGA			
<i>groEL</i> _Forward	1029159	ACCTTGACAGGATCAATAAT	132 pb	35	52
<i>groEL</i> _Reverse		TAAAAATGCTGGTTATGAAG			
16S_Foward*	NP722374	TATCTTCCTCAATTAACCA	181pb	27	54
16S_Reverse*		TAGTTAAGATTGCCAGTGAT			

\* Stipp et al., 2008.

(after 30 min of exposure to the antimicrobial agent) and their respective controls (after 1 min of exposure) were applied an ANOVA test. The gene expression by planktonic cells submitted to 1% (v/v) DMSO vehicle for 30 min was also used as a control. The *p* values < 5% were considered statistically significant for all tests.

## Results

Halistanol sulfate A (**1**), isolated from the sponge *Petromica ciocalyptoides* (Kossuga et al., 2007) and the modified diketopiperazines rodriguesines A (**2**) and B (**3**) isolated from the ascidian *Didemnum* sp. (Kossuga et al., 2009) (Fig. 1) inhibited the growth of *Streptococcus mutans* strains in low concentrations. Other bacteria growth were inhibited with higher concentrations of these antimicrobials agents (> 62.5 µg/ml).


Table 2 shows the MIC and MBC values, as well as the MBC/MIC ratio, displayed by the halistanol sulfate A (**1**) and by rodriguesines A (**2**) and B (**3**). Rodriguesines A (**2**) and B (**3**) were inseparable under several HPLC conditions, as previously reported by Kossuga et al., (2009), and were tested in combination. Both samples presented a broad inhibitory spectrum against the tested bacterial strains. Rodriguesines A (**2**) and B (**3**) inhibited all of the tested bacteria, with MIC values between 15-125 µg/ml and MBC values between 31.2-250 µg/ml. These modified diketopiperazines had a bactericidal effect on all the bacterial strains except *Staphylococcus aureus* ATCC 6538. Halistanol sulfate A showed the lowest MIC and MBC values when tested on *Streptococcus mutans*. However, its inhibitory effect against other species was much less potent (MIC > 125 µg/ml) (Table 2). Therefore, it appears that halistanol sulfate A selectively inhibits the *Streptococcus mutans* pathogen at low concentrations, without modifying the ecological balance between more and less pathogenic microorganisms. The specificity of an oral antimicrobial agent at low concentrations

against *Streptococcus mutans* is important for reducing the selective pressure on other bacterial species, particularly initial colonizers, such as *Streptococcus sanguinis*. Therefore, both halistanol sulfate A and rodriguesines A and B were selected for additional studies of their antibiofilm activities.

The pathogenic *Streptococcus mutans* CI and *Streptococcus mutans* UA159 showed response to CV-stained biofilms greater than 2.0 at  $\lambda_{\max}$  OD<sub>575</sub> nm in control conditions and were categorized as good biofilm formers (Loo et al., 2000). Fig. 1 shows the effect of rodriguesines A and B and halistanol sulfate A on planktonic cells and on biofilm formation, as well as the MIC values of biofilm formation by *Streptococcus mutans* strains. No significant differences in the inhibition of planktonic cells were observed in these MIC compared to the DMSO control (Student's *t* test, *p* > 0.05). As for the MIC of biofilm formation, we observed reductions of up to 38% in the optical density of planktonic cells (at  $\lambda_{\max}$  OD<sub>550</sub> nm) relative to the DMSO vehicle (Fig. 1). There were no significant differences in the viable cells counts of the planktonic cultures (Student's *t* test, *p* > 0.05, data not shown). These results indicate that these compounds present modest selectivity in their inhibition of biofilm formation, despite their ability to inhibit the growth of planktonic cells.

Halistanol sulfate A (**1**) showed potent antibacterial activity against *Streptococcus mutans* CI and *Streptococcus mutans* UA159 strains at 3.0 µg/ml, inhibiting approximately 85.2 and 99.5% the biofilm formation by these species, respectively. The strain *Streptococcus mutans* UA159 was more susceptible to the antimicrobial activity of all tested substances in comparison with *Streptococcus mutans* CI. Biofilm formation by *Streptococcus mutans* UA159 was inhibited by 77.5 to 100%, while *Streptococcus mutans* CI biofilm formation was inhibited by 34.8 to 85.2% after antimicrobial treatment during 18 h.

Antimicrobial samples were evaluated on planktonic cells for 1 min (T1) for control samples and after 30 min (T30) of treatment. Diketopiperazines (**2**) and (**3**) and halistanol



**Figure 1** – Antimicrobial effect of rodriguesines A/B, halistanol sulfate A and doxycycline on *Streptococcus mutans* Clinical Isolated CI and *Streptococcus mutans* UA159 planktonic cells and biofilm formation after 18 h. The percentage of planktonic cell growth (white bars) or biofilm formation (gray bars) in the treated groups was calculated based on the DMSO control groups (100%). All the bars represent the means ± standard deviations from three independent experiments. An asterisk indicates that the cell growth between the DMSO and treated groups was significantly different (Student test,  $p < 0.05$ ).

**Table 2**

Antimicrobial activities (values of MIC, MBC and the MBC/MIC ratio in µg/ml, bactericidal or bacteriostatic effects) of halistanol sulfate A and rodriguesines A/B against oral streptococci and medical pathogenic bacteria.

Antimicrobial samples	Antimicrobial activities	Oral streptococci				Medical pathogenic bacteria		
		<i>Streptococcus mutans</i> clinical isolate	<i>Streptococcus mutans</i> UA159	<i>Streptococcus sobrinus</i> ATCC 27607	<i>Streptococcus sanguinis</i> ATCC 15300	<i>Escherichia coli</i> NCTC 8196	<i>Enterococcus faecalis</i> ATCC 14506	<i>Staphylococcus aureus</i> ATCC 6538
Rodriguesines A and B	MIC	31.25	62.5	125	125	125	125	62.5
	MBC	62.5	62.5	250	125	250	125	250
	MBC:MIC	2	1	2	1	2	1	4
	Effect	bactericidal	bactericidal	bactericidal	bactericidal	bactericidal	bactericidal	bacteriostatic*
Halistanol sulfate A	MIC	15	15	250	250	-	250	-
	MBC	31.25	31.25	500	500	-	500	-
	MBC:MIC	2	2	2	2	-	2	-
	Effect	bactericidal	bactericidal	bactericidal	bactericidal	-	bactericidal	-

The antimicrobials were considered bactericidal when the MBC:MIC ratio was between 1:1 to 2:1 and bacteriostatic if this ratio was higher than 2:1 (Galvão et al., 2012). DMSO (1%) was used as a negative control. (\*)The bacteriostatic effect of the tested samples could be bactericidal if bacterial inocula were used at approximately  $1 \times 10^5$  CFU/ml, as described by CLSI (2009).

sulfate A (**1**) affected the expression of both *S. mutans* UA 159 and *S. mutans* CI genes involved in cell adhesion and biofilm formation, except the expression of *groEL* and 16S genes. The ANOVA test showed a statistically significant difference ( $p < 0.05$ ) in the levels of *gtfB*, *gtfC* and *gpbB* gene transcription after the treatment of planktonic cells in the mid-log phase with the antimicrobial compounds. A 15 to 85% reduction of the gene transcription levels relatively to the respective controls (T1) was observed. The gene expression was normalized relatively to 16S gene transcription, which showed invariable expression in all tested conditions (data not shown), as well as relative to *groEL*, which codes for the chaperone GroEL, a class I heat shock protein of *Streptococcus mutans*.


Comparing DMSO 1% at T1 and at T30, we observed an increase of 12 to 55% (average  $26.3 \pm 15.3$ ) in the transcription levels of biofilm-associated genes of *S. mutans* CI only, at T30 (data not shown). Fig. 2 shows the percentage of *gtfB*, *gtfC*, *gpbB* and *groEL* transcription levels after treatment with the compounds **1** - **3** after 30 min. Treatment with compounds **1** - **3** inhibited gene expression between 1.5 and 82% ( $51.56 \pm 18.52$ ) (Fig. 2). We observed a decrease in the gene expression levels of all tested genes, except *gtfC* transcription in *S. mutans* CI, which was not altered after treatment with halistanol sulfate A (ANOVA,  $p > 0.05$ ). In addition, there were no alterations in the expression levels of the *groEL* and 16S genes after the antimicrobial treatments.

The gene *gpbB* was more severely affected after treatment for 30 min with the antimicrobial compounds, with an expression reduction of 53 to 82% for both pathogenic bacterial strains. The expression of *gtfB* and *gtfC* decreased by approximately 1.5 to 57%, depending on the tested strain and the antimicrobial compounds (Fig. 2). The *gpbB* gene of *Streptococcus mutans* UA159 strain was inhibited (average  $72.5\% \pm 13.4$ ) to a higher level than the *gtf* genes (average  $35.8\% \pm 8.0$ ). A higher inhibition of *gtfB* expression by *Streptococcus mutans* CI was observed relatively to *Streptococcus mutans* UA159.

## Discussion

A series of marine natural products inhibit biofilm formation of *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Bordetella bronchiseptica*, *Staphylococcus aureus* and *Streptococcus mutans* (Huigens et al., 2007, Rogers et al., 2008, Liu et al., 2011). In the present investigation we verified that halistanol sulfate A (**1**), isolated from the sponge *Petromica ciocalyptoides* (Kossuga et al., 2007), and the mixture of modified diketopiperazines rodriguesines A (**2**) and B (**3**) isolated from an ascidian of the genus *Didemnum* (Kossuga et al., 2009), displayed low MIC values against *Streptococcus mutans*. The rodriguesines A and B had broad antimicrobial spectra, inhibiting both oral streptococci as well as pathogenic bacteria with low MIC values.

Diketopiperazines are dipeptides that have also been isolated from Gram-positive bacteria, fungi and higher organisms (De Carvalho and Abraham, 2012). Diketopiperazines are reported to modulate the LuxR-mediated quorum-sensing systems of Gram-negative and Gram-positive bacteria and are considered to influence cell-cell bacterial signaling, offering alternative ways to control biofilms by interfering with



**Figure 2** – The percentage of *gtfB*, *gtfC* and *groEL* relative expression by planktonic cells of Clinical Isolated (CI) *Streptococcus mutans* and *Streptococcus mutans* UA159 after treatment for 30 min with rodriguesines A/B, halistanol sulfate A and doxycycline in the MIC of biofilm inhibition. Data were expressed relative to treatment with 1% DMSO for 30 min (control vehicle). An asterisk represents that there is no significant difference relative to control (ANOVA  $p > 0.05$ ). The mRNA expression levels were calibrated using 16S rRNA. Bars represent the means  $\pm$  standard deviations of triplicate experiments.

microbial communication (Withers et al., 2001; Abraham, 2005; Ryan and Dow, 2008). *Streptococcus mutans* LuxS affects acid and oxidative-stress tolerance, bacteriocin production and biofilm formation (Merritt et al., 2003; 2005, Wen and Burne, 2004; Yoshida et al., 2005; Wen et al., 2011).

In the present investigation, the modified diketopiperazines rodriguesines A (**2**) and B (**3**) inhibited *Streptococcus mutans* biofilm formation and downregulated *gtfB*, *gtfC* and *gpbB* expression but did not change *groEL* and 16S expression. Diketopiperazines may act on the Lux-RS quorum-sensing system (Withers et al., 2001; Abraham, 2005; Ryan and Dow, 2008), although it appears that the *S. mutans* *gtfB*, *gtfC* and *gpbB* genes are under the control of a distinct quorum-sensing system (two-component system - TCS). Inhibitors of the quorum-sensing TCS system and/or PknB may be important

for the control of *S. mutans* because these systems regulate the transcription of several virulence target genes, some of which have been studied in detail (Li et al., 2001; Van Der Ploeg Jr, 2005). However, the lack of a detailed knowledge of the involved cell-cell communication mechanisms has hindered the development of effective therapeutic strategies to fight bacterial infections. The increased interest in studying the bacterial lifestyle in biofilm formation should lead to the discovery of new therapeutic strategies for caries treatment (Bayles, 2007).

Halistanol sulfate A inhibited biofilm formation by two *Streptococcus mutans* strains at low MIC, but it did not inhibit the initial colonizers represented by *S. sanguinis*. This activity profile is highly suitable because antibacterial agents that inhibit pathogenic bacteria and biofilms without affecting the healthy normal flora can be used in preventive treatments based on specifically targeted molecules. Agents with this mechanism tend to not disturb the ecological balance between pathogens and commensal residents in the oral cavity (Liu et al., 2011).

Halistanol sulfates A-E were first described as antifungal and antithrombin agents (Kanazawa et al., 1992). Previously it has been observed that halistanol sulfate A affects *Staphylococcus aureus* cells, possibly via a disruption membrane mechanism (Marinho et al., 2012). Sterol sulfates related to halistanol trisulfate A may act as detergents, inhibiting the growth of Gram-positive bacteria without affecting Gram-negative bacteria, possibly via a membranolytic mechanism and may also inhibit bacterial protein-tyrosine kinases (Slate, et al., 1994).

*Streptococcus mutans* CI showed higher resistance to the antibiofilm action of small-molecules than *S. mutans* UA159. The resistance mechanisms or intermediary resistance to antimicrobials agents are known to be variable among bacterial genotypes. The existence of a high diversity of *S. mutans* genotypes in human oral cavities (Kamiya et al., 2005) suggests that such antibacterial small-molecules should be tested against different strains with possible distinct resistance phenotype/genotypes to antimicrobial agents. The mechanisms controlling the expression of the *gtfB*, *gtfC* and *gbpB* virulence genes are strain-specific, illustrating the genetic diversity of *S. mutans* (Stipp et al., 2008). Such genotypic variation may have further implications in the selection of therapeutic targets to control this cariogenic bacterium.

After antimicrobial treatment with halistanol, rodriguesines or doxycycline, there was a significant reduction in the *Streptococcus mutans* biofilm formation, although there were no alterations in the *groEL* expression levels. The gene *groEL* codes for a chaperone, that has essential roles in cellular metabolism by assisting in the folding of newly synthesized or denatured proteins, as well as in the assembly, transport, and degradation of cellular proteins (Bukau and Horwich, 1998). In *S. mutans*, GroEL is classified as a class I heat shock protein as well is the chaperone DnaK. Lemos et al. (2007), showed that the down-regulation of *dnaK* and *groEL* reduced *S. mutans* biofilm formation and acid tolerance, suggesting that the regulatory response by *S. mutans* to environmental stresses for the development and maturation of a biofilm has a significant influence on the biofilm structure. Our data suggest that *groEL*

expression is not essential for the development of *S. mutans* biofilms because *groEL* is not a target gene of halistanol sulfate A and rodriguesines. These antimicrobial agents inhibited biofilm formation by the down regulation of *gtfB* / *gtfC* and *gbpB*, which codes for a water-insoluble glucan matrix and increases the interaction between *S. mutans* and glucans of the matrix.

The evaluation of natural products against different *Streptococcus mutans* genotypes is important for establishing the antibacterial effectiveness of these substances and for elucidating their effects on the selective pressure of possible *S. mutans* genotypes that are resistant to antimicrobial agents. The results presented herein illustrate the potential of marine secondary metabolites to act as antibacterial agents against *S. mutans* with distinct modes-of-action. Further studies are necessary to clarify the mechanisms-of-action of rodriguesines and halistanol sulfate A, their effects on the regulation of other *S. mutans* virulence genes and/or *S. mutans* quorum sensing signaling pathways, and their respective species-specific antibacterial activities in multi-species biofilm models, both *in vitro* and *in vivo*, associated with transcriptome and/or proteome analyses.

---

## Conclusion

The modified diketopiperazines rodriguesines A and B as well as halistanol sulfate A displayed inhibitory effects on biofilm formation by *Streptococcus mutans* and down regulate the expression of the *gtfB*, *gtfC* and *gbpB* genes. The antimicrobial properties of halistanol sulfate A suggest that it may be useful in the development of new drug candidates for anti-*Streptococcus mutans* therapy because it selectively inhibits biofilm formation by *S. mutans* strains at low concentrations, not affecting initial buccal colonizing species such as *S. sanguinis*.

---

## Authors' contributions

BAL performed antimicrobial assays, inhibitory assays on biofilm formation, gene expression assays, statistical analysis and contributed extensively for the manuscript preparation. SPL performed the isolation and identification of halistanol sulfate from the sponge *Petromica ciocalyptoides*. MHK performed the isolation and identification of rodriguesines A and B from the ascidian *Didemnum* sp. RBG obtained financial support, provided laboratory infrastructure and facilities for the antibacterial experiments at the Faculdade de Odontologia de Piracicaba, Universidade Estadual de Campinas and conceived the experiments and the methodology to evaluate the results. RGSB obtained financial support, laboratory infrastructure and facilities for the natural products isolation and identification experiments at the Instituto de Química de São Carlos, Universidade de São Paulo and designed the isolation and identification procedures and also contributed extensively for the manuscript preparation. RUK performed the inhibitory tests against microorganisms and inhibition of biofilm formation, standardized the protocols of RNA extraction and

semi-quantitative RT-PCR and performed statistical analysis. Contributed extensively to the preparation of the manuscript. All the authors have read the final manuscript and approved its submission.

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgments

This study was financially supported by Fundação de Amparo à Pesquisa de São Paulo (FAPESP, grants 2005/60175-2 and 2010/50190-2) and Fundação de Desenvolvimento da Unicamp (FUNCAMP, grant Faepex 68807). B.A.L. was supported by a FAPESP MSc scholarship (proc. 2007/01078-2), R.U.K. was supported by FAPESP (proc. 2005/58479-8 and 2006/60037-1), while S.P.L. and M.H.K. were supported by CAPES and CNPq scholarships.

## REFERENCES

- Abraham, W-R, 2005. Controlling gram-negative pathogenic bacteria by interfering with their biofilm formation. *Drug Des. Rev.* 2, 13-33.
- Bader, J.D., Shugars, D.A., Bonito, A.J., 2001. Systematic reviews of selected dental caries diagnostic and management methods. *J. Dent. Educ.* 65, 960-968.
- Bayles, K.W., 2007. The biological role of death and lysis in biofilm development. *Nat. Rev. Microbiol.* 5, 721-726.
- Blunt, J.W., Copp, B.R., Keyzers, R.A., Munro, M.H.G., Prinsep, M.R., 2014. Marine natural products. *Nat. Prod. Rep.* 31, 160-258.
- Bukau, B., Horwich, A.L., 1998. The Hsp70 and Hsp60 chaperone machines. *Cell.* 92, 351-366.
- Butler, M.S., Robertson, A.A.B., Cooper, M.A., 2014. Natural product and natural product derived drugs in clinical trials. *Nat. Prod. Rep.* 31, 1612-1661.
- Chia, J.S., Lee, Y.Y., Huang, P.T., Chen, J.Y., 2001. Identification of stress-responsive genes in *Streptococcus mutans* by differential display reverse transcription-PCR. *Infect. Immun.* 69, 2493-2501.
- CLSI, 2009. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically*, Approved Standard, 29, no. 2, Clinical And Laboratory Standards Institute document M07-A8, Wayne, Pa, USA, 8<sup>th</sup> ed.
- Coenye, T., Honraet, K., Rigole, P., Nadal, J.P., Nelis, H.J., 2007. In vitro inhibition of *Streptococcus mutans* biofilm formation on hydroxyapatite by subinhibitory concentrations of anthraquinones. *Antimicrob. Agents. Chemother.* 51, 1541-1544.
- De Carvalho, M.P., Abraham, W.R., 2012. Antimicrobial and biofilm inhibiting diketopiperazines. *Curr. Med. Chem.* 19, 3564-3577.
- Duque, C., Stipp, R.N., Wang, B., Smith, D.J., Höfling, J.F., Kuramitsu, H.K., Duncan, M.J., Mattos-Graner, R.O., 2011. Downregulation of GbpB, a component of the VicRK regulon, affects biofilm formation and cell surface characteristics of *Streptococcus mutans*. *Infect. Immun.* 79, 786-796.
- Galvão, L.C., Furletti, V.F., Bersan, S.M., Da Cunha, M.G., Ruiz, A.L., De Carvalho, J.E., Sartoratto, A., Rehder, V.L.G., Figueira, G.M., Duarte, M.C.T., Ikegaki, M., de Alencar, S. M., Rosalen, P.L., 2012. Antimicrobial activity of essential oils against *Streptococcus mutans* and their antiproliferative effects. *Evid. Based Complement. Alternat. Med.* 2012, <http://dx.doi.org/10.1155/2012/751435>
- Huigens III, R.W., Richards, J.J., Parise, G., Ballard, T.E., Zeng, W., Deora, R., Melander, C., 2007. Inhibition of *Pseudomonas aeruginosa* biofilm formation with Bromoageliferin analogues. *J. Am. Chem. Soc.* 129, 6966-6967.
- Kamiya, R.U., Höfling, J.F., Gonçalves, R.B., 2008. Frequency and expression of mutacin biosynthesis genes in isolates of *Streptococcus mutans* with different mutacin-producing phenotypes. *J. Med. Microbiol.* 57, 626-635.
- Kamiya, R.U., Napimoga, M.H., Rosa, R.T., Höfling, J.F., Gonçalves, R.B., 2005. Mutacins production in *Streptococcus mutans* genotypes isolated from caries-affected and caries-free individuals. *Oral. Microbiol. Immunol.* 20, 20-24.
- Kanazawa, S., Fusetani, N., Matsunaga, S., 1992. Halistanol sulfates A-E, new steroid sulfates, from a marine sponge, *Epipolysis* sp. *Tetrahedron.* 48, 5467-5472.
- Klein, M.I., Flório, F.M., Pereira, A.C., Höfling, J.F., Gonçalves, R.B., 2004. Longitudinal study of transmission, diversity, and stability of *Streptococcus mutans* and *Streptococcus sobrinus* genotypes in Brazilian nursery children. *J. Clin. Microbiol.* 42, 4620-4626.
- Koo, H., Gomes, B.P., Rosalen, P.L., Ambrosano, G.M., Park, Y.K., Cury, J.A., 2000. In vitro antimicrobial activity of propolis and *Arnica montana* against oral pathogens. *Arch. Oral Biol.* 45, 141-148.
- Koo, H., Hayacibara, M.F., Schobel, B.D., Cury, J.A., Rosalen, P.L., Park, Y.K., Vacca-Smith, A.M., Bowen, W.H., 2003. Inhibition of *Streptococcus mutans* biofilm accumulation and polysaccharide production by apigenin and tt-farnesol. *J. Antimicrob. Chemother.* 52, 782-789.
- Koo, H., Seils, J., Abranches, J., Burne, R.A., Bowen, W.H., Quivey, R.G. Jr., 2006. Influence of apigenin on *gtf* gene expression in *Streptococcus mutans* UA159. *Antimicrob. Agents Chemother.* 50, 542-546.
- Kossuga, M.H., De Lira, S.P., McHugh, S., Torres, Y., Lima, B.A., Gonçalves, R.B., Veloso, K., Ferreira, A.G., Rocha, R.M., Berlinck, R.G.S., 2009. Antibacterial modified diketopiperazines from two ascidians of the genus *Didemnum*. *J. Braz. Chem. Soc.* 20, 704-711.
- Kossuga, M.H., De Lira, S.P., Nascimento, A.M., Gambardella, M.T.P., Berlinck, R.G.S., Torres, Y.R., Nascimento, G.G.F., Pimenta, E.F., Silva, M., Thiemann O.H., Oliva, G., Tempone, A.G., Melhem, M.S.C., de Souza, A.O., Galetti, F.C.S., Silva, C.L., Cavalcanti, B., Pessoa, C.O., Moraes, M.O., Hajdu, E., Peixinho, S., Rocha, R.M., 2007. Isolamento e atividades biológicas de produtos naturais das esponjas *Monanchora arbuscula*, *Aplysina* sp., *Petromica ciocalyptoides* e *Topsentia ophiraphidites*, da ascídia *Didemnum ligulum* e do octocoral *Carijoa riisei*. *Quim. Nova.* 30, 1194-1202.
- Kossuga, M.H., Romminger, S., Xavier, C., Milanetto, M.C., Do Valle, M.Z., Pimenta, E.F., Morais, R.P., Carvalho, E., Mizuno, C.M., Coradello, L.F.C., Barroso, V. M., Vacondio, B., Javaroti, D.C.D., Seleglim, M.H.R., Cavalcanti, B.C., Pessoa, C., Moraes, M.O., Lima, B.A., Gonçalves, R.B., Bonugli-Santos, R.C., Sette, L.D., Berlinck, R.G.S., 2012. Evaluating methods for the isolation of marine-derived fungal strains and production of bioactive secondary metabolites. *Rev. Bras. Farmacogn.* 22, 257-267.
- Lemos, J.A., Luzardo, Y., Burne, R.A., 2007. Physiologic effects of forced down-regulation of *dnaK* and *groEL* expression in *Streptococcus mutans*. *J. Bacteriol.* 189, 1582-1588.



- Li, Y.H., Lau, P.C., Lee, J.H., Ellen, R.P., Cvitkovitch, D.G., 2001. Natural genetic transformation of *Streptococcus mutans* growing in biofilms. *J. Bacteriol.* 183, 897-908.
- Liu, C., Worthington, R.J., Melander, C., Wu, H., 2011. A new small molecule specifically inhibits the cariogenic bacterium *Streptococcus mutans* in multispecies biofilms. *Antimicrob. Agents Chemother.* 55, 2679-2687.
- Lonn-Stensrud, J., Petersen, F.C., Benneche, T., Scheie, A.A., 2007. Synthetic bromated furanone inhibits autoinducer-2-mediated communication and biofilm formation in oral streptococci. *Oral Microbiol. Immunol.* 22, 340-346.
- Loo, C.Y., Corliss, D.A., Ganeshkumar, N., 2000. *Streptococcus gordonii* biofilm formation: identification of genes that code for biofilm phenotypes. *J. Bacteriol.* 182, 1374-1382.
- Mai, J., Tian, X-L., Gallant, J.W., Merkle, N., Biswas, Z., Syvitski, R., Douglas, S.E., Ling, J., Li, Y.H., 2011. A novel target-specific, salt-resistant antimicrobial peptide against the cariogenic pathogen *Streptococcus mutans*. *Antimicrob. Agents Chemother.* 55, 5205-5213.
- Marinho, P.R., Simas, N.K., Kuster, R.M., Duarte, R.S., Fracalanza, S.E., Ferreira, D.F., Romanos, M.T., Muricy, G., Giambiagi-Demarval, M., Laport, M.S., 2012. Antibacterial activity and cytotoxicity analysis of halistanol trisulphate from marine sponge *Petromica citrina*. *J. Antimicrob. Chemother.* 67, 2396-2400.
- Mattos-Graner, R.O., Jin, S., King, W.F., Chen, T., Smith, D.J., Duncan, M.J., 2001. Cloning of the *Streptococcus mutans* gene encoding glucan binding protein B and analysis of genetic diversity and protein production in clinical isolates. *Infect. Immun.* 69, 6931-6941.
- Mayer, A.M., Rodríguez, A.D., Tagliatalata-Scafati, O., Fusetani, N., 2013. Marine pharmacology in 2009-2011: marine compounds with antibacterial, antidiabetic, antifungal, anti-inflammatory, antiprotozoal, antituberculosis, and antiviral activities; affecting the immune and nervous systems, and other miscellaneous mechanisms of action. *Mar. Drugs.* 11, 2510-2573.
- Merritt, J., Kreth, J., Shi, W., Qi, F., 2005. LuxS controls bacteriocin production in *Streptococcus mutans* through a novel regulatory component. *Mol. Microbiol.* 57, 960-969.
- Merritt, J., Qi, F., Goodman, S.D., Anderson, M.H., Shi, W., 2003. Mutation of LuxS affects biofilm formation in *Streptococcus mutans*. *Infect. Immun.* 71, 1972-1979.
- Murata, R.M., Almeida, L.S.B., Yatsuda, R., Dos Santos, M.H., Nagem, T.J., Rosalen, P.L., Koo, H., 2008. Inhibitory effects of 7-epiclusianone on glucan synthesis, acidogenicity and biofilm formation by *Streptococcus mutans*. *FEMS Microbiol. Lett.* 282, 174-181.
- O'Toole, G.A., Kolter, R., 1998. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol. Microbiol.* 28, 449-461.
- Oliveira, J.H., Selegim, M.H., Timm, C., Grube, A., Kock, M., Nascimento, G.F.G., Martins, A.C.T., Silva, E.G.O., de Souza, A.O., Minarini, P.R.R., Galetti, F.C.S., Silva, C.L., Hajdu E., Berlinck, R.G.S., 2006. Antimicrobial and antimycobacterial activity of cyclostelletamine alkaloids from sponge *Pachychalina* sp. *Mar. Drugs.* 4, 1-8.
- Pasquantonio, G., Greco, C., Prenna, M., Ripa, C., Vitali, L.A., Petrelli, D., Di Luca, M.C., Ripa, S., 2008. Antibacterial activity and anti-biofilm effect of chitosan against strains of *Streptococcus mutans* isolated in dental plaque. *Int. J. Immunopathol. Pharmacol.* 21, 993-997.
- Rogers, S.A., Melander, C., 2008. Construction and screening of a 2-aminoimidazole library identifies a small molecule capable of inhibiting and dispersing bacterial biofilms across order, class, and phylum. *Angew. Chem. Int. Ed. Engl.* 47, 5229-5231.
- Ryan, R.P., Dow, J.M., 2008. Diffusible signals and interspecies communication in bacteria. *Microbiol.* 154, 1845-1858.
- Selegim, M.H.R., De Lira, S.P., Kossuga, M.H., Batista, T., Berlinck, R.G.S., Hajdu, E., Muricy, G., da Rocha, R.M., do Nascimento, G.G.F., Silva, M., Pimenta, E.F., Thiemann, O.H., Oliva, G., Cavalcanti, B.C., Pessoa, C., de Moraes, M.O., Galetti, F.C.S., Silva, C.L., de Souza, A.O., Peixinho, S., 2007. Antibiotic, cytotoxic and enzyme inhibitory activity of crude extracts from Brazilian marine invertebrates. *Rev. Bras. Farmacogn.* 17, 287-318.
- Slate, D.L., Lee, R.H., Rodriguez, J., Crews, P., 1994. The marine natural product, halistanol trisulfate, inhibits pp60v-src protein tyrosine kinase activity. *Biochem. Biophys. Res. Commun.* 203, 260-264.
- Smith, D.J., 2002. Dental caries vaccines: Prospects and concerns. *Crit. Rev. Oral Biol. Med.* 13, 335-349.
- Stipp, R.N., Goncalves, R.B., Höfling, J.F., Smith, D.J., Mattos-Graner, R.O., 2008. Transcriptional analysis of *gtfB*, *gtfC*, and *gcbP* and their putative response regulators in several isolates of *Streptococcus mutans*. *Oral Microbiol. Immunol.* 23, 466-473.
- Van Der Ploeg Jr, 2005. Regulation of bacteriocin production in *Streptococcus mutans* by the quorum-sensing system required for development of genetic competence. *J. Bacteriol.* 187, 3980-3989.
- Wen, Z.T., Burne, R.A., 2004. LuxS-mediated signaling in *Streptococcus mutans* is involved in regulation of acid and oxidative stress tolerance and biofilm formation. *J. Bacteriol.* 186, 2682-2691.
- Wen, Z.T., Nguyen, A.H., Bitoun, J.P., Abranches, J., Baker, H.V., Burne, R.A., 2011. Transcriptome analysis of LuxS-deficient *Streptococcus mutans* grown in biofilms. *Mol. Oral Microbiol.* 26, 2-18.
- Whiley, R.A., Beighton, D., 1998. Current classification of the oral streptococci. *Oral Microbiol. Immunol.* 13, 195-216.
- Withers, H., Swift, S., Williams, P., 2001. Quorum sensing as an integral component of gene regulatory networks in gram-negative bacteria. *Curr. Opin. Microbiol.* 4, 186-193.
- Yoshida, A., Ansai, T., Takehara, T., Kuramitsu, H.K., 2005. LuxS-based signaling affects *Streptococcus mutans* biofilm formation. *Appl. Environ. Microbiol.* 71, 2372-2380.