

Low levels of crustacean parasite infestation in fish species from the Matapi River in the state of Amapá, Brazil

Baixos níveis de infestação de parasitos crustáceos em espécies de peixes do Rio Matapi no estado do Amapá (Brasil)

Ligia Rigôr Neves¹ ; Marcos Tavares-Dias^{1,2*} 

¹ Programa de Pós-graduação em Biodiversidade e Biotecnologia da Amazônia Legal – Bionorte, Universidade Federal do Amapá – UNIFAP, Macapá, AP, Brasil

² Embrapa Amapá, Macapá, AP, Brasil

Received November 17, 2018

Accepted January 30, 2019

Abstract

This first study investigated the crustacean parasite fauna in 66 species of fish from the Matapi River basin, state of Amapá (Brazil). Fish were collected every two months between March 2012 and August 2013, encompassing dry and rainy seasons. Among the 66 species examined (corresponding to 722 fish specimens) only 11 species were parasitized. The infestation prevalence was 2.2%, and a total of 48 specimens of parasites were distributed between three different parasite groups. These included *Argulus elongatus*, *Dolops reperta* and *Argulus multicolor* (Branchiura), *Ergasilus xinguensis* and *Gamidactylus* sp. (Copepoda), and Isopoda (*Braga patagonica*), but branchiuran species were predominant. This was the first report of these parasite species for *Leporinus fasciatus*, *Astyanax bimaculatus*, *Curimata incompta*, *Pygocentrus nattereri*, *Crenicichla cincta*, *Crenicichla johanna*, *Geophagus camopiensis*, *Pterophyllum scalare*, *Plagioscion squamosissimus*, *Hypostomus plecostomus* and *Propimelodus eigenmanni*. Lastly, this study expands the range of occurrence of these six parasite species to the Matapi River basin in eastern Amazon.

Keywords: Amazon, ectoparasites, freshwater fish, infestation.

Resumo

Este primeiro estudo investigou a fauna parasitária de crustáceos em 66 espécies de peixes da bacia do Rio Matapi, estado do Amapá (Brasil). Os peixes foram coletados bimestralmente no período de março de 2012 a agosto de 2013, nas estações de estiagem e chuvosa. Entre as 66 espécies (correspondendo a 722 espécimes) somente 11 espécies estavam parasitadas. A prevalência de infestação foi 2,2% e um total de 48 espécimes foram distribuídos em três grupos de parasitos. Esses incluem Branchiura (*Argulus elongatus*, *Dolops reperta* e *Argulus multicolor*), Copepoda (*Ergasilus xinguensis* e *Gamidactylus* sp.) e Isopoda (*Braga patagonica*), mas a dominância foi de espécies de branchiuras. Este foi o primeiro relato dessas espécies de parasitos para *Leporinus fasciatus*, *Astyanax bimaculatus*, *Curimata incompta*, *Pygocentrus nattereri*, *Crenicichla cincta*, *Crenicichla johanna*, *Geophagus camopiensis*, *Pterophyllum scalare*, *Plagioscion squamosissimus*, *Hypostomus plecostomus* e *Propimelodus eigenmanni*. Por fim, este estudo expande a ocorrência dessas seis espécies de parasitos para a bacia do Rio Matapi na Amazônia oriental.

Palavras-chave: Amazônia, ectoparasitos, peixes de água doce, infestação.

The Matapi River basin is located in the coastal-estuarine sector of the state of Amapá, within the municipality of Santana (Brazil). The Flexal and Pirativa rivers and the Maruanum and Lago creeks are the main tributaries of the basin. The predominant vegetation is composed of aquatic macrophytes. There is still little human influence in this basin, despite various urban and agricultural activities. This basin is flooded daily by tides of the Amazon River, which considerably influence the hydrodynamics

of the floodplain forest environments and other flooded areas (CUNHA et al., 2011; SILVA et al., 2016).

The highest water velocities, reaching approximately 1 m/s, occur during relatively short periods, amounting to close to two-fifths of the complete tidal cycle (12.9 h). Outside of these periods, the water velocity is approximately 0.5 m/s (CUNHA et al., 2011). Consequently, these hydrodynamic conditions also influence the lives of the 104 known species of fish in this basin, which include Characiformes (70.2%), Cichliformes (17.2%), Siluriformes (8.8%), Clupeiformes (1.7%), Tetraodontiformes (1.6%), Gymnotiformes (0.2%) and Beloniformes (0.05%) (SILVA et al., 2016). Despite

*Corresponding author: Marcos Tavares-Dias. Embrapa Amapá, Rodovia Juscelino Kubitschek, Km 5, 2600, CEP 68903-419, Macapá, AP, Brasil. e-mail: marcos.tavares@embrapa.br



the rich ichthyofauna of the Matapi River, little is known about the diversity of crustacean parasites infesting these fishes.

Among crustaceans, there are many ectoparasites of fish. These are found in various habitats and require a host during at least one phase of their life cycle (MAMANI et al., 2004; TAVARES-DIAS et al., 2015; OLIVEIRA et al., 2017). Ergasilidae, Argulidae, Lernaeidae, Lernaeopodidae and Cymothoidae are the families most frequently found, and these infest the gills, oral cavity, nostrils and tegument of fish in Brazil. Parasitic crustaceans are just some of the ectoparasites taxa found on freshwater fish species in Brazil (TAVARES-DIAS et al., 2015). Many of these parasites have received significant attention because of the damage that they cause to fishery resources, which includes reduced fish growth, reproduction and host activities such as natation (MAMANI et al., 2004; TAVARES-DIAS et al., 2015). Some of these parasite crustaceans require specific fish as hosts and parasitize specific sites, particularly in fish with certain lifestyles, while other parasites do not have any preferences (TAVARES-DIAS et al., 2015; OLIVEIRA et al., 2017). Thus, some species of these ectoparasites have a broad pattern of distribution in different

places, while others are restricted to certain geographical areas (TAVARES-DIAS et al., 2015) and host species.

The objective of the present study was to investigate the species of crustaceans infesting the gills of 66 species of fish in the Matapi River basin, in state of Amapá, northern Brazil.

Between March 2012 and August 2013, 66 species of fish were sampled every two months from the Matapi River (Table 1), in the municipality of Santana, state of Amapá, Brazil (Figure 1) for analyses on crustacean parasites in their gills, operculum and tegument. Fish were collected using gill nets of various mesh sizes (20, 25, 30, 35, 40 and 70 mm between knots), matapi traps, casting nets, hand lines and longlines, and the mean duration of fishing was 8 hours. Sampling was conducted in two seasonal periods: the dry period between July and November 2012 and in August 2013 (temperature of 28.3 ± 1.6 °C, pH 5.5 ± 0.7 , electrical conductivity 3.1 ± 1.9 $\mu\text{S}/\text{cm}$, suspended solids 24.0 ± 22.5 mg/L, transparency 83.9 ± 55.3 cm and rainfall 100.5 ± 128.1 mm); and the rainy period between March and May 2012 and between January and March 2013 (temperature of 27.6 ± 1.0 °C, pH 5.5 ± 0.6 , electrical

Table 1. Body parameters of the fish species collected in Matapi River, state of Amapá (Brazil).

Order/Family/Species	N	Weight (g)	Length (cm)
CHARACIFORMES			
Acestrorhynchidae			
<i>Acestrorhynchus falcatus</i> Bloch, 1794	2	88.0 \pm 8.4	23.5 \pm 0.7
<i>Acestrorhynchus falcistrostris</i> Cuvier, 1819	2	62.0 \pm 0	17.8 \pm 0.4
Anostomidae			
<i>Leporinus fasciatus</i> Bloch, 1794	2	165.6 \pm 110.9	22.7 \pm 9.6
<i>Leporinus friderici</i> Bloch, 1794	50	44.6 \pm 31.2	14.0 \pm 3.8
<i>Schizodon fasciatum</i> Spix & Agassiz, 1829	1	85.0 \pm 34.8	18.7 \pm 2.0
Characidae			
<i>Astyanax bimaculatus</i> Linnaeus, 1758	75	11.7 \pm 4.8	8.2 \pm 0.9
<i>Bryconamericus stramineus</i> Eigenmann, 1908	5	20.4 \pm 8.0	12.2 \pm 1.1
<i>Hemibrycon surinamensis</i> Géry, 1962	4	13.5 \pm 5.7	7.6 \pm 0.4
<i>Moenkhausia lepidura</i> Kner, 1858	8	9.6 \pm 5.5	9.5 \pm 1.5
<i>Tetragonopterus chalcus</i> Spix & Agassiz, 1829	40	19.6 \pm 17.4	9.3 \pm 2.1
Triporthidae			
<i>Triporthus albus</i> Cope, 1872	1	12.0 \pm 0	12.5 \pm 0
<i>Triporthus angulatus</i> Spix & Agassiz, 1829	4	51.5 \pm 15	15.1 \pm 1.6
<i>Triporthus elongatus</i> Günther, 1864	2	135.0 \pm 9.8	23.7 \pm 0.3
<i>Triporthus rotundatus</i> Jardine, 1841	1	32.0 \pm 0	10.0 \pm 0
Curimatidae			
<i>Curimata acutirostris</i> Vari & Reis, 1995	1	64.0 \pm 0	16.0 \pm 0
<i>Curimata cyprinoides</i> Linnaeus, 1766	8	45.7 \pm 30.1	13.5 \pm 3.0
<i>Curimata incompta</i> Vari, 1984	132	33.7 \pm 17.9	12.5 \pm 2.7
<i>Curimatella alburna</i> Muller & Troschel, 1844	1	96.0 \pm 0	18.5 \pm 0
<i>Curimata inornata</i>	5	30.0 \pm 6.0	10.1 \pm 0.8
Cynodontidae			
<i>Rhaphiodon vulpinus</i> Spix & Agassiz, 1829	1	28.0 \pm 0	15.0 \pm 0
Erythrinidae			
<i>Hoplias malabaricus</i> Bloch, 1794	2	68.0 \pm 28.2	16.7 \pm 0.3
Hemiodontidae			
<i>Hemiodus unimaculatus</i> Bloch, 1794	48	30.8 \pm 13.1	12.4 \pm 3.2

N: Sample number.

Table 1. Continued...

Order/Family/Species	N	Weight (g)	Length (cm)
Serrasalimidae			
<i>Metynnys lippincottianus</i> Cope, 1870	89	11.1 ± 6.8	7.4 ± 1.2
<i>Myleus rubripinnis</i> Muller & Troschel, 1844	4	41.5 ± 33.1	10.3 ± 2.5
<i>Pygocentrus nattereri</i> Kner, 1858	6	36.6 ± 28.7	10.5 ± 2.5
<i>Serrasalmus calmoni</i> Steindachner, 1908	4	24.2 ± 31.9	9.7 ± 2.8
<i>Serrasalmus rhombeus</i> Linnaeus, 1766	9	17.7 ± 20.5	6.5 ± 2.5
<i>Serrasalmus spilopleura</i> Kner, 1858	6	14.6 ± 9.8	9.3 ± 2.5
CLUPEIFORMES			
Engraulidae			
<i>Anchoviella guianensis</i> Eigenmann, 1912	3	6.6 ± 1.1	8.0 ± 2.5
<i>Lycengraulis batesii</i> Günther, 1868	1	16.0 ± 0	14.0 ± 0
<i>Pterengraulis atherinoides</i> Linnaeus, 1766	4	45.0 ± 23.0	16.6 ± 2.1
Pristigasteridae			
<i>Ilisha amazonica</i> Miranda Ribeiro, 1920	3	26.6 ± 3.0	14.8 ± 1.0
<i>Pellona flavipinnis</i> Valenciennes, 1847	2	86.0 ± 16.9	19.2 ± 1.7
<i>Pellona harroweri</i> Fowler, 1917	1	286.0 ± 0	34.5 ± 0
<i>Pristigaster cayana</i> Cuvier, 1829	3	1.5 ± 0.5	5.0 ± 1.0
GYMNOTIFORMES			
Gymnotidae			
<i>Gymnotus carapo</i> Linnaeus, 1758	1	10.0 ± 0	15.5 ± 0
CICHLIFORMES			
Cichlidae			
<i>Crenicichla cincta</i> Regan, 1905	1	0.188 ± 0	24.0 ± 0
<i>Crenicichla johanna</i> Heckel, 1840	2	136.0 ± 73.5	21.7 ± 3.8
<i>Crenicichla strigata</i> Günther, 1862	2	25.0 ± 21.2	13.2 ± 4.5
<i>Geophagus camopiensis</i> Pellegrin, 1903	4	47.2 ± 22.2	13.5 ± 2.7
<i>Mesonauta festivus</i> Heckel, 1840	6	19.6 ± 10.8	9.5 ± 1.5
<i>Pterophyllum scalare</i> Schultze, 1823	12	9.8 ± 5.4	7.6 ± 1.5
Sciaenidae			
<i>Pachypops fourcroyi</i> La Cepède, 1802	4	18.6 ± 13.2	11.6 ± 2.2
<i>Plagioscion auratus</i> Castelnau, 1855	14	101.6 ± 86.6	19.9 ± 5.5
<i>Plagioscion squamosissimus</i> Heckel, 1840	51	56.3 ± 54.4	15.7 ± 5.7
<i>Plagioscion surinamensis</i> Bleeker, 1973	5	13.6 ± 8.1	8.9 ± 3.1
Eleotridae			
<i>Eleotris pisonis</i> Gmelin, 1789	1	2.0 ± 0	6.0 ± 0
SILURIFORMES			
Ageneiosidae			
<i>Ageneiosus ucayalensis</i> Castelnau, 1855	10	55.8 ± 19.6	19.8 ± 4.5
Auchenipteridae			
<i>Centromochlus heckelii</i> De Filippi, 1853	9	2.1 ± 0	5.9 ± 1.1
<i>Parauchenipterus galeatus</i> Linnaeus, 1766	1	38.0 ± 0	15.5 ± 0
Heptapteridae			
<i>Pimelodella eigenmanni</i> Boulenger, 1891	8	18.0 ± 8.3	16.1 ± 2.1
<i>Pimelodella altipinnis</i>	7	29.7 ± 4.3	13.1 ± 0.8
Loricariidae			
<i>Ancistrus hoplogenyis</i> Gunther, 1864	3	22.3 ± 2.0	11.6 ± 0.2
<i>Ancistrus</i> sp.	1	28.0 ± 0	9.0 ± 0
<i>Hypostomus plecostomus</i> Linnaeus, 1758	7	155.7 ± 104.8	45.2 ± 61.7
<i>Hypostomus ventromaculatus</i> Boeseman, 1968	1	86.0 ± 0	18.5 ± 0
<i>Hypostomus watwata</i> Hancock, 1828	1	82.0 ± 0	19.0 ± 0
<i>Loricaria cataphracta</i> Linnaeus, 1758	8	1186.5 ± 3310.6	15.3 ± 1.6
<i>Panaque</i> Eigenmann & Eigenmann, 1889	1	20.0 ± 0	8.0 ± 0
<i>Peckoltia brevis</i> La Monte, 1935	1	16.0 ± 0	9.0 ± 0

N: Sample number.

Table 1. Continued...

Order/Family/Species	N	Weight (g)	Length (cm)
<i>Peckoltia lineola</i> Armbruster, 2008	1	38.0 ± 0	11.5 ± 0
Pimelodidae			
<i>Pimelodus blochii</i> Valenciennes, 1840	1	10.0 ± 0	9.5 ± 0
<i>Pimelodus ornatus</i> Kner, 1858	4	76.0 ± 73.9	16.3 ± 5.0
<i>Platynemataichthys notatus</i> Jardine, 1841	1	72.0 ± 0	20.0 ± 0
<i>Propimelodus eigenmanni</i> Van der Stigchel, 1946	8	20.8 ± 9.4	15.9 ± 2.0
TETRAODONTIFORMES			
Tetraodontidae			
<i>Colomesus asellus</i> Muller & Troschel, 1849	17	17.7 ± 9.1	8.6 ± 1.2
Total	722	-	-

N: Sample number.




Figure 1. Collection sites of the fish species in Matapi River, state of Amapá, Brazil.

conductivity $2.5 \pm 2.0 \mu\text{S}/\text{cm}$, suspended solids $26.8 \pm 30.5 \text{ mg}/\text{L}$, transparency $86.2 \pm 61.5 \text{ cm}$ and rainfall $338.4 \pm 111.6 \text{ mm}$.

The present study was conducted in accordance with the recommendations of the Brazilian College for Animal Experimentation (Colégio Brasileiro de Experimentação Animal, COBEA) and with authorization from the Ethics Committee for Use of Animals of Embrapa Amapá (Protocol N° 014 - CEUA/CPAFAP).

The tegument and operculum of all the fish were examined at the collection site and the gills were then fixed in formalin (5%). The gills were removed and analyzed using a stereomicroscope. The crustaceans found were fixed in 70% ethyl alcohol with 10% glycerin (EIRAS et al., 2006) and were then identified (LUQUE et al., 2013; TABORDA et al., 2016). The ecological terms used were those recommended by Bush et al. (1997).

A total of 722 fish of 66 species were collected. Their length and weight are described in Table 1. The host families Serrasalmidae and Characidae predominated, and nine species accounted for 70% of all the individuals examined: *Metynnis lippincottianus*, *Curimata incompta*, *Astyanax bimaculatus*, *Hemiodus unimaculatus*, *Tetragonopterus chalcus*, *Moenkhausia lepidura*, *Leporinus frederici* and *Plagioscion squamosissimus*. However, *M. lippincottianus* and *C. incompta* were the most abundant species (Table 1), and they were present in all sampling sites. The number of individuals collected was greater during the dry season, corresponding to 63% of the total sample.

The prevalence of crustacean parasites was 2.2% among the 722 fish of 66 species that were examined. Among the 48 parasite specimens collected, there was high abundance of three taxa (Branchiura, Copepoda and Isopoda) with diversity of six species (Table 2). Alsarakibi et al. (2014) reported that the density of argulids was lower in lotic environments such as rivers, in comparison with lentic environments such as fish farms. Vasconcelos & Tavares-Dias (2016) studied the crustacean parasite fauna in six species of fish in a reservoir in the state of Amapá and reported prevalence of 30.4%. They collected 878 parasites, which included one species of Branchiura, one of Copepoda and one of Isopoda. Oliveira et al. (2017) reported that in 13 host species

of the Jari River (state of Amapá), the prevalence of parasites was 63.8%; they collected 399 parasite specimens. However, the diversity of crustacean parasites and their levels of infestation can be influenced by various factors relating to the biology of parasites and hosts, and by environmental factors, among others (CARVALHO et al., 2003; MAMANI et al., 2004; FONTANA et al., 2012; ALSARAKIBI et al., 2014; MIKHEEV et al., 2015; VASCONCELOS & TAVARES-DIAS, 2016). Fish hosts may be are used by these parasites for transportation (CARVALHO et al., 2003), which could be facilitated by migrating fish species. However, this low prevalence of parasites in the hosts of the Matapi River may be due to the influence of daily tides from the Amazon River, considering that certain crustacean parasites respond to water movement to reach their hosts (MIKHEEV et al., 2015).

No crustacean parasites were found on the body surface of the fish from the Matapi River that were examined. However, among fish living in lotic environments, it is more difficult to estimate the prevalence of parasite crustaceans in the tegument when hosts are caught using fishing nets because these fish make strong movements and attempt to resist. Moreover, with time, the hosts' stress increases and their metabolism tends to decrease, which could stimulate these ectoparasites in the tegument of the fish to explore new habitats for their survival (BRANDÃO et al., 2013). Another factor that needs to be considered is the strong influence of diurnal tides from the Amazon River on the velocity of the Matapi River (CUNHA et al., 2011; SILVA et al., 2016), given that these parasites need to swim to find adequate hosts, while others depend on the flow of water and swimming speed (FONTANA et al., 2012; MIKHEEV et al., 2015).

Among the hosts in the Matapi River, Argulidae species predominated. Among these, *Argulus elongatus* Heller, 1857, and *Dolops reperta* Bouvier, 1899, were the most prevalent. However, *Ergasilus xinguensis* Taborda, Paschoal & Luque, 2016, was the most abundant species, even though it only infested the cichlids *Crenicichla johanna* Heckel, 1840, *Crenicichla cincta* Regan, 1905 and *Geophagus camopiensis* Pellegrin, 1903, and sciaenid *Plagioscion squamosissimus* Heckel, 1840 (Table 2). Branchiurans

Table 2. Species of parasite crustaceans in fish gills from the Matapi River, state of Amapá (Brazil).

Host species	<i>Dolops reperta</i>			<i>Argulus elongatus</i>			<i>Argulus multicolor</i>			<i>Ergasilus xinguensis</i>			<i>Braga patagonica</i>			<i>Gamidactylus sp.</i>		
	P (%)	MI	MA	P (%)	MI	MA	P (%)	MI	MA	P (%)	MI	MA	P (%)	MI	MA	P (%)	MI	MA
<i>Leporinus fasciatus</i>	0	0	0	20.0	1.0	0.2	0	0	0	0	0	0	20.0	1.0	0.2	0	0	0
<i>Astyanax bimaculatus</i>	1.3	1.0	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Curimata incompta</i>	0.7	1.0	0.007	0	0	0	0	0	0	0	0	0	0	0	0	0.7	1.0	0.007
<i>Pygocentrus nattereri</i>	0	0	0	12.0	1.0	0.1	0	0	0	0	0	0	0	0	0	0	0	0
<i>Crenicichla cincta</i>	0	0	0	0	0	0	0	0	0	100	18.0	18.0	0	0	0	0	0	0
<i>Crenicichla johanna</i>	0	0	0	0	0	0	0	0	0	100	7.5	7.5	0	0	0	0	0	0
<i>Geophagus camopiensis</i>	0	0	0	0	0	0	0	0	0	25.0	2.0	0.5	0	0	0	0	0	0
<i>Pterophyllum scalare</i>	8.3	1.0	0.08	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Plagioscion squamosissimus</i>	0	0	0	0	0	0	0	0	0	3.9	1.0	0.03	0	0	0	0	0	0
<i>Hypostomus plecostomus</i>	0	0	0	0	0	0	28.0	1.0	0.2	0	0	0	0	0	0	0	0	0
<i>Propimelodus eigenmanni</i>	0	0	0	0	0	0	0	0	0	0	0	0	12.0	2.0	0.2	0	0	0

P: Prevalence, MA: Mean abundance, MI: Mean intensity.

are known for frequently switching hosts and for having lower host specificity than that of other parasite groups. Among the factors that influence the infestation rates of these ectoparasites are the genetic similarities of the hosts and ecological factors that may be involved (MAMANI et al., 2004; OLIVEIRA et al., 2017).

To conclude, in 66 host species, the rates of infestation with crustacean parasites were low, and most of the fish examined had low abundance of parasites, except *C. johanna* and *G. camopiensis*. Lastly, this study reports these fish species as new hosts for the crustacean parasites species that were found and expands the range of occurrence of the six parasite species to the Matapi River basin in eastern Amazon (Brazil).

Acknowledgements

The authors thank to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) by the doctoral grant was granted to the first author and to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the productivity research grant awarded to M. Tavares-Dias (# 303013/2015-0).

References

- Alsarakibi M, Wadeh H, Li G. Influence of environmental factor on *Argulus japonicus* occurrence of Guangdong province, China. *Parasitol Res* 2014; 113(11): 4073-4083. <http://dx.doi.org/10.1007/s00436-014-4076-5>. PMID:25127735.
- Bush AO, Lafferty KD, Lotz JM, Shostak W. Parasitology meets ecology on its own terms: Margolis et al. revisited. *J Parasitol* 1997; 83(4): 575-583. <http://dx.doi.org/10.2307/3284227>. PMID:9267395.
- Brandão H, Toledo GM, Wunderlich AC, Ramos IP, Carvalho ED, Silva RJ. Occurrence of *Braga cigarra* (Cymothoidae) parasitizing *Galeocharax knerii* (Characidae) from affluents of Jurumirim reservoir, Brazil. *Rev Bras Parasitol Vet* 2013; 22(2): 292-296. <http://dx.doi.org/10.1590/S1984-29612013005000002>. PMID:23459850.
- Carvalho LN, Del-Claro K, Takemoto RM. Host-parasite interaction between branchiurans (Crustacea: Argulidae) and piranhas (Osteichthyes: Serrasalminidae) in the Pantanal wetland of Brazil. *Environ Biol Fishes* 2003; 67(3): 288-296. <http://dx.doi.org/10.1023/A:1025899925545>.
- Cunha AC, Pinheiro LAR, Cunha HFA, Schulz HE, Brasil ACP Jr, Souza EB. Simulação da hidrodinâmica e dispersão de poluentes com monitoramento virtual no Rio Matapi-AP. *REA* 2011; 13(2): 18-32. <http://dx.doi.org/10.7867/1983-1501.2011v13n2p18-32>.
- Eiras JC, Takemoto RM, Pavanelli GC. *Métodos de estudo e técnicas laboratoriais em parasitologia de peixes*. Maringá: Eduem; 2006.
- Fontana M, Takemoto RM, Malta JCO, Mateus LAF. Parasitism by argulids (Crustacea: Branchiura) in piranhas (Osteichthyes: Serrasalminidae) captured in the Caiçara bays, upper Paraguay River, Pantanal, Mato Grosso State, Brazil. *Neotrop Ichthyol* 2012; 10(3): 653-659. <http://dx.doi.org/10.1590/S1679-62252012005000019>.
- Luque JL, Vieira FM, Takemoto RM, Pavanelli GC, Eiras JC. Checklist of Crustacea parasitizing fishes from Brazil. *Check List* 2013; 9(6): 1449-1470. <http://dx.doi.org/10.15560/9.6.1449>.
- Mamani M, Hamel C, Van Damme PA. Ectoparasites (Crustacea: Branchiura) of *Pseudoplatystoma fasciatum* (surubí) and *P. tigrinum* (chuncuina) in Bolivian white water floodplains. *Ecol Bolív* 2004; 39(2): 9-20.
- Mikheev VN, Pasternak AF, Valtonen ET. Behavioural adaptations of argulid parasites (Crustacea: Branchiura) to major challenges in their life cycle. *Parasit Vectors* 2015; 8(1): 394. <http://dx.doi.org/10.1186/s13071-015-1005-0>. PMID:26205259.
- Oliveira MSB, Corrêa LL, Oliveira Ferreira D, Neves LR, Tavares-Dias M. Records of new localities and hosts for crustacean parasites in fish from the eastern Amazon in northern Brazil. *J Parasit Dis* 2017; 41(2): 565-570. <http://dx.doi.org/10.1007/s12639-016-0852-8>. PMID:28615880.
- Silva LMA, Lima, JF, Tavares-Dias M. *Ichtiofauna como indicadora da qualidade ambiental do Rio Matapi, Afluente do Rio Amazonas no estado do Amapá (Brasil)*. Macapá: Embrapa Amapá; 2016. (Boletim de Pesquisa e Desenvolvimento; vol. 92).
- Taborda NL, Paschoal F, Luque JL. A new species of *Ergasilus* (Copepoda: Ergasilidae) from *Geophagus altifrons* and *G. argyrostictus* (Perciformes: Cichlidae) in the Brazilian Amazon. *Acta Parasitol* 2016; 61(3): 549-555. <http://dx.doi.org/10.1515/ap-2016-0073>. PMID:27447219.
- Tavares-Dias M, Dias-Júnior MB, Florentino AC, Silva LM, da Cunha AC. Distribution pattern of crustacean ectoparasites of freshwater fish from Brazil. *Rev Bras Parasitol Vet* 2015; 24(2): 136-147. <http://dx.doi.org/10.1590/S1984-29612015036>. PMID:26154954.
- Vasconcelos HCG, Tavares-Dias M. Host-parasite interaction between crustaceans of six fish species from the Brazilian Amazon. *Acta Sci Biol Sci* 2016; 38(1): 113-123. <http://dx.doi.org/10.4025/actascibiolsci.v38i1.29601>.