

## Successful Improvement of Frequency and Symptoms of Premature Complexes after Oral Magnesium Administration

Cristina Nádja Muniz Lima De Falco<sup>1,2</sup>, Cesar Grupi<sup>1</sup>, Eduardo Sosa<sup>1</sup>, Maurício Scanavacca<sup>1</sup>, Denise Hachul<sup>1</sup>, Sissy Lara<sup>1</sup>, Luciana Sacilotto<sup>1</sup>, Cristiano F. Pisani<sup>1</sup>, José A. F. Ramires<sup>1</sup>, Francisco Darrieux<sup>1</sup>

Instituto do Coração da Faculdade de Medicina da Universidade de São Paulo<sup>1</sup>, São Paulo; Clínica Médica Core Vita<sup>2</sup>, Mogi Mirim, SP, Brazil

### Abstract

**Background:** Premature ventricular and supraventricular complexes (PVC and PsVC) are frequent and often symptomatic. The magnesium (Mg) ion plays a role in the physiology of cell membranes and cardiac rhythm.

**Objective:** We evaluated whether the administration of Mg Pidolate (MgP) in patients with PVC and PsVC is superior to placebo (P) in improving symptoms and arrhythmia frequency.

**Methods:** Randomized double-blind study with 60 consecutive symptomatic patients with more than 240 PVC or PsVC on 24-hour Holter monitoring who were selected to receive placebo or MgP. To evaluate symptom improvement, a categorical and a specific questionnaire for symptoms related to PVC and PsVC was made. Improvement in premature complex density (PCD) per hour was considered significant if percentage reduction was  $\geq 70\%$  after treatment. The dose of MgP was 3.0 g/day for 30 days, equivalent to 260mg of Mg element. None of the patients had structural heart disease or renal failure.

**Results:** Of the 60 patients, 33 were female (55%). Ages ranged from 16 to 70 years old. In the MgP group, 76.6% of patients had a PCD reduction  $>70\%$ , 10% of them  $>50\%$  and only 13.4%  $<50\%$ . In the P group, 40% showed slight improvement,  $<30\%$ , in the premature complexes frequency ( $p < 0.001$ ). Symptom improvement was achieved in 93.3% of patients in the MgP group, compared with only 16.7% in the P group ( $p < 0.001$ ).

**Conclusion:** Oral Mg supplementation decreases PCD, resulting in symptom improvement. (Arq Bras Cardiol 2012;98(6):480-487)

**Keywords:** Arrhythmias, cardiac; ventricular premature complexes; magnesium; ion channels.

### Introduction

Premature ventricular and supraventricular complexes (PVC and PsVC) are frequent and often symptomatic. Their prevalence can occur in up to 50% of the general population, especially<sup>1</sup>. The incidence of this arrhythmia increases with age<sup>2-5</sup>. The studies show that most of these patients had  $<1$  PVC per hour, usually monomorphic and single forms<sup>2,3</sup>.

Symptoms related to PVC and PsVC can be very troublesome or even disabling. Patients may refer to these symptoms as a "skipped beat," "punch in the chest," palpitations, dyspnea, cough, dizziness, atypical chest pain, and near syncope<sup>1,6</sup>, affecting quality of life. These symptoms are usually noticed when the premature complex density is high. Premature complexes are directly related to heart cell excitability, which is influenced by electrolyte balance in intracellular fluid. The interaction between magnesium (Mg) and calcium (Ca) has particular

relevance in the regulation of nerve and muscle cell permeability<sup>7-9</sup>, and in the ATPase – Na<sup>+</sup>/K<sup>+</sup> pump<sup>8,9</sup>. By acting on the physiology of cell membranes, Mg has a special role in cardiac rhythm maintenance<sup>8,9</sup>.

Mg is the second most abundant intracellular cation<sup>8,9</sup> and plays an important role in the activity of many coenzymes and ATP-dependent reactions, including membrane-dependent energy transport<sup>9,10</sup>. Less than 1% of magnesium is found in blood<sup>9</sup> and only approximately 0.3% in serum<sup>11,12</sup>. Therefore, the intracellular deficiencies can be underdiagnosed<sup>8</sup>.

Lifestyles characterized by stress, low micronutrient intake, physical training, sleep deprivation, and the use of certain medications (diuretics, aminoglycosides, and cyclosporine) may lead to Mg deficiency. This ion is mainly found in seeds, nuts, vegetables, and wheat bran. In the general population, magnesium deficiency probably occurs due to low magnesium dietary intake<sup>13,14</sup>, that needs to maintain adequate intracellular values and, in the elderly, by decreasing appetite<sup>14</sup>.

This study aimed to assess whether oral administration of magnesium pidolate in patients with PVC or PsVC is superior to placebo in improving symptoms and the frequency of premature complex (PC), as well as whether symptom improvement is related to a significant reduction in arrhythmia frequency.

**Mailing Address:** Cristina Nádja Muniz Lima De Falco •

Rua Cesar Henrique Costa, 76, Santa Cruz. Postal Code 13806-401, Mogi Mirim, SP - Brazil

E-mail: cnmfalco@cardiol.br, cristinanadjadefalco@usp.br

Manuscript received January 17, 2012; manuscript revised January 17, 2012; accepted March 05, 2012.

## Methods

### Study Design and Participants

Patients were recruited from the Arrhythmia Unit of the Heart Institute (InCor), University of São Paulo Medical School and Core Vita Clinics. Patients were eligible if they were symptomatic and had more than 240 PVC or PsVC per day on 24h Holter monitoring (or more than 10/hour). Exclusion criteria were impaired renal function, structural heart disease (except mitral valve prolapse without regurgitation), or the use of concomitant drugs. The study was approved by the ethics committees of the participating centers in the study with the number CAPPesq-0613/10.

### Procedures

After providing written informed consent, all trial participants were randomly assigned to receive placebo (P) or magnesium pidolate (MgP), each administered in a blinded manner. All the patients did echocardiogram and none of them had structural heart disease or electrolyte disturbances; the determination of electrolytes (magnesium, sodium, calcium and potassium) and the renal function were normal.

The dose of the MgP was 3.0 g/day for 30 days, which contains 260mg of Mg element. The magnesium, sodium, calcium, and potassium serum dosage was taken at baseline, at 15 days, and at 30 days after randomization.

### Holter

The 24-hour Holter (3 channels) was performed at baseline and 30 days after the medication use. The count of the premature complexes was performed following the institution protocol and the premature complex density (PCD) was performed dividing the total number of atrial and ventricular extra systoles Holter counted in the number of hours of recording.

### Outcomes

Follow-up visits occurred at randomization, after 15 days, and after 30 days.

To evaluate symptom improvement, a specific questionnaire related to PC was made with the following

### Quiz

Symptom	Points
Failures or "leaps" in the chest	
Couch with palpitation	
Dizziness	
Dyspnea	
Sudoresis and/or chest pain	
Total	

### Points

- A - Absent (never) - 0 point
- B - Infrequent (once to 5 times/day) - 1 point
- C - Frequent (6 to 20 times/day) - 2 points
- D - Very Frequent ( $\geq 21$  times/day) - 3 points

### "Quality of Life" Score

#### Symptom Classification

- I - Asymptomatic - 0 point
- II - Mid Symptomatic - 01 a 05 points
- III - Moderate Symptomatic - 06 a 11 points
- IV - Severe Symptomatic - more than 12 points

questions: 1- Failure or "leaps" in the chest; 2-Couch with palpitations; 3-Dizziness; 4-Dyspnea; 5-Sudoresis and/or chest pain. According to frequency of symptoms was made a "score" (Fig. 1). For this "score", was only considered an improvement, if the patient had a reduction of at least two categories, for example, was in the score IV before treatment and migrated to the score I or II after treatment. Furthermore, it was made a categorical classification of patients with questioning whether there was improvement of symptoms, with only answers "yes" or "no".

### Statistical Analysis

To meet the objectives of the study, we calculated the percentage changes in PCD /hour, and was considered success criteria, after treatment, the percentage reduction  $\geq 70\%$  per 24-hour Holter. Data are presented using summary statistics (mean, standard deviation, median, minimum, maximum). Outcome variables were compared between groups using a Mann-Whitney test<sup>15</sup>. The improvement of either PCD or symptoms was described using absolute and relative values. The existence of an association between groups and the improvement of each criterion was performed using Fisher's exact test<sup>15</sup>. A p value of 0.05 was considered statistically significant. We described the values of magnesium and potassium before medication, and at 15 and 31 days after medication use, according to groups, and also compared the values between groups and moments using analysis of variance with repeated measures and 2 factors<sup>16</sup>.

A statistical power of 80% was chosen to detect a 60% symptom reduction with MgP and 30% with P, with a confidence interval of 95%.

## Results

A total of 60 patients were enrolled in the program between October 2010 and August 2011. Both treatment groups had similar baseline characteristics. The mean age was 46.47 (MgP) to 48.53 (P), and 55% were women (Table 1). Age description and PCD variation according to each group are described in Table 1. The average extra-systole in 24 hours in both groups was higher than 4.955 PC/day. Twelve patients had mitral valve prolapse without insufficiency (5 in group P and 7 in group MgP).

Figure 1 - This figure shows the score system to assess improvement of symptoms before and after the drugs in both groups (placebo and magnesium pidolate).

### Frequency of Premature Complexes

In the MgP group, 76.6% of patients had a PCD reduction >70%, 10% of them >50% and only 13.4% <50%. In the P group, 40% showed slight improvement, <30%, in the PC frequency and none of them had improvement >70% (Table 2/ Figure 2). Patients in the MgP group had an average reduction of 77.13% (SD = 24.57%) of PCD, whereas the placebo group had an average increase of 47.99% (SD = 158.93%) in PCD ( $p < 0.001$ ) (Table 2/Figure 3). This difference was also consistent for both PVC and PsVC ( $p < 0.001$ ) (Table 2). A few patients (13.4%) had only a slight improvement in PC density in the MgP group (<50% in PC frequency), but had improvement in symptoms.

### Symptoms

In the P group, only 16.7% reported symptom improvement (score), whereas in the MgP group there was an improvement of 93.3% ( $p < 0.001$ ) (Table 2). As a categorical variable, there was improvement of symptoms in 93.3% of patients in the MgP, compared with only 13.3% in the placebo group ( $p < 0.001$ ) (Table 2). Figures 4 and 5 show the superiority of symptom improvement in patients receiving MgP compared with patients receiving P.

### Laboratory Findings

There were no significant changes in serum magnesium, potassium, sodium and calcium during the study in both groups. Serum magnesium did not differ significantly between groups ( $p = 0.743$ ) or between periods of time (before and after oral supplementation;  $p = 0.154$ ); moreover, serum potassium did not differ statistically between groups or during follow-up ( $p = 0.415$ ,  $p = 0.804$ , respectively). Table 2 and Figure 6 illustrate the changes in magnesium.

### Study Discontinuation and Adverse Events

Only one patient in the MgP group had to discontinue the protocol after 10 days due to diarrhea, which was promptly resolved within 24 hours.

## Discussion

Oral magnesium supplementation not only decreases

the density of premature ventricular and supraventricular complexes, but also improves symptoms compared to placebo. In the placebo group, only 16.7% showed improvement in symptoms compared to 93.3% of patients using MgP. Although most patients have improved PC density and symptoms, some patients have improved symptoms without a significant drop in PC density. The worsening of PCD in some patients in the placebo group can perhaps be explained by the great variability that involves spontaneous idiopathic arrhythmias. However, it is important to note that both groups were subjected to this variability, and even then, there was a statistically significant reduction of PCD in the magnesium group compared with the placebo group.

The mechanisms by which magnesium administration reduces the incidence of PC are not entirely known. Magnesium is regarded as a significant regulator of cardiac cell function. Depletion of magnesium, as shown in some studies, may be proarrhythmic<sup>14</sup>. Zehender et al<sup>17</sup> demonstrated that increased intake of potassium and magnesium in patients with frequent ventricular arrhythmias can result in a moderate but significant antiarrhythmic effect, although the frequency of tachyarrhythmia and symptoms has not been changed<sup>17</sup>. However, this sample differs from our results, probably because the population is different: most patients had other cardiac comorbidities, and some were using other medications.

On the possibility of oral magnesium overload, it is important to note that patients in the MgP group had a normal Mg serum dosage during the 30-day follow-up, without major adverse effects or the need for suspension (except for a single patient with diarrhea). It is likely that intracellular levels of Mg are low, despite normal serum dosage, since serum dosage corresponds to only 0.3% of total magnesium. It would be interesting to find out minimally invasive, accurate methods to detect intracellular Mg levels<sup>18</sup> to better understand many diseases, including cardiac arrhythmias.

There are some studies with fluorescent markers specific to magnesium (and Mg-Fluo-4/AM KMG-20/AM)<sup>19</sup>, that aim at the cation dosage in various cell types, such as platelets, red blood cells, lymphocytes<sup>12</sup>, cardiomyocytes (in rats)<sup>19</sup> and epithelial sublingual cells<sup>18</sup>. Silver et al<sup>18</sup> showed a significant correlation between the magnesium values in heart muscle biopsies from bypass surgery, with the values measured in

Table 1 - Baseline Characteristics of Study Patients

Characteristic	MgP	Placebo	p value
Gender Female, n (%)	17 (56,7%)	16 (53,3%)	0,795
Age (SD) #	46,47 ± 17,42	48,53 ± 13,18	0,606
Valve Mitral Prolapses (SD)	07 (23,3%)	05 (16,7%)	0,519
PCD*	256,41 ± 294,56	206,46 ± 249,89	0,496
PVC density	129,31 ± 219,13	90,90 ± 157,72	0,639
PsVC density	126,83 ± 225,82	115,55 ± 242,03	0,066
Magnesium	2,05 ± 0,13	2,08 ± 0,14	0,439
Potassium	4,20 ± 0,36	4,23 ± 0,36	0,690

\*Density of PC, PVC, PsVC: premature complexe per hour; # - Standard Deviation.

Table 2 - Clinical Outcomes at 30 Days

Variable	MgP	Placebo	p
Improvement of PCD, n (%)			< 0.001
< 0%	0 (0.0%)	18 (60%)	
0 to 30%	2 (6.7%)	12 (40%)	
30 to 50%	2 (6.7%)	0 (0.0%)	
50 to 70%	3 (10.0%)	0 (0.0%)	
70% or more	23 (76.6%)	0 (0.0%)	
Improvement of symptoms, n (%)			< 0.001
No	2 (6.7%)	26 (86.7%)	
Yes	28 (93.3%)	4 (13.3%)	
Improvement of score, n (%)			< 0.001
Less than 2 points	2 (6.7%)	25 (83.3%)	
Two points or more	28 (93.3%)	5 (16.7%)	
Reduction of PCD, % (SD)	77.13 ± 24.57	-47.99 ± 158.93	< 0.001
Reduction of PVC*	31.83 ± 280.62	-40.78 ± 187.34	< 0.001
Reduction of PsVC	22.64 ± 223.05	-212.78 ± 732.13	< 0.001
Magnesium (SD)	2.09 ± 0.10	2.09 ± 0.12	< 0.743
Potassium (SD)	4.17 ± 0.27	4.26 ± 0.29	< 0.415

\*Density of PC, PVC, PsVC: premature complexe per hour.




Figure 2 - Distribution of patients according to the percentage of improvement in the density of ventricular premature beats after 30 days in the placebo group (green) and magnesium pidolate (blue).




Figure 3 - Box-plot of the density of ventricular premature beats per time, before and after the treatment, according to each group: magnesium pidolate (blue) and placebo (green).




Figure 4 - Percentage of patients who presented improvement of the symptoms: Yes (blue) or No (green), according to the group of treatment.




Figure 5 - Percentage of patients who presented improvement in the score of symptoms. Smaller than 02 points (green) and greater than 02 points (blue).




Figure 6 - Serum levels of magnesium before and on the 15th and 30th day of treatment with placebo (black line) and magnesium pidolate (red line).

sublingual epithelial cell smear. Moreover, magnetic resonance imaging (MRI) is a noninvasive test for tissue Mg analysis. Recently, the Transient Receptor Potential Melastation channel and its role in hypomagnesemia were identified. Chronic Mg deficiency may be explained by changes in these specific membrane transporters<sup>12,20,21</sup>.

Magnesium is closely related to the maintenance of cellular ionic balance in combination with sodium, potassium, and calcium; it cooperates with the Na<sup>+</sup>/K<sup>+</sup> ATPase pump<sup>8,9,14</sup>. Magnesium deficiency causes an increase in intracellular Na<sup>+</sup>, which allows cellular K<sup>+</sup> loss<sup>9</sup>. Mg also affects calcium homeostasis, and many calcium channels are magnesium dependent. In addition, Mg is needed for the release and action of parathyroid hormone<sup>9</sup>.

Some conditions are associated with magnesium deficiency, such as metabolic syndrome<sup>22,23</sup>, hypertension, congestive heart failure, diabetes, preeclampsia, and arrhythmias, and some clinical studies show that supplementation was beneficial in controlling these diseases<sup>14</sup>. A single study showed that 365 mg of magnesium per day for 8 weeks lowers blood pressure<sup>24</sup>. According to Tong and Rude, magnesium reduces irregular heartbeat, and because intracellular Mg depletion may be present despite a normal serum Mg, magnesium deficiency must always be considered as a potential factor in cardiac dysrhythmias<sup>25</sup>. In the intensive care unit, magnesium is used when arrhythmias do not respond to conventional medications, and also in digitalis intoxication<sup>26</sup>. In heart failure, magnesium deficiency is caused by diuretic therapy, which increases the incidence of arrhythmia, among them premature complexes<sup>8,14</sup>.

The Mg balance among different compartments of the body occurs slowly, so that the concentration of Mg in one tissue does not correspond to that in another<sup>12</sup>. Less than 1% of total body Mg is present in blood<sup>9</sup> and only 0.3% in serum<sup>11,12</sup>, so serum levels do not reflect the total body stores<sup>27,28</sup>. This may be one explanation for the improvement of arrhythmia with the replacement of Mg compared to the placebo group, even without changes in serum levels of this ion.

Although symptom improvement could be associated with a reduction in PC density, even in some cases in which this decline is not significant, symptoms also improved. This is partly explained by the placebo effect, as well as magnesium action in nerve cells. A nutrition education rich in magnesium may be a practical alternative for the treatment of this arrhythmia. However, this information must be evaluated in another study. It is also important to control serum Mg during treatment and to assess renal function.

### Limitations

Long-term follow-up was not performed; therefore, it is not possible to establish whether these patients are free from symptom recurrence after oral replacement of magnesium. Intracellular magnesium was not measured, but Mg serum

dosages have shown that this replacement was safe and effective. The goal of reduction in the density of premature ventricular and supraventricular complexes and symptom improvement was achieved. The specific symptom score was not validated, because it does not exist in the literature. However, a simple categorical score was also made, with a good correlation between reduced PCD and symptom improvement. Moreover, the aim of this study was not to prevent life threatening arrhythmic events. It shows that the data should not be used as justification for treating patients with this objective, especially those with heart disease.

### Conclusion

Simple oral Mg replacement reduced the density of premature ventricular and supraventricular complexes and specially improved symptoms in our study population (no cardiac heart disease). Clinical and molecular studies are needed to evaluate intracellular Mg and develop better targets for the daily needs of this ion, show probable deficiencies, and explain how to prevent and better treat patients with symptomatic premature ventricular and supraventricular complexes and no apparent heart disease.

### Acknowledgments

The authors wish to thank the external collaborators Mrs. Alessandra Manera, Cacilda Motta, and Kérolem Ignatius for their technical support and Prof. Dr. Paulo Sampaio Gutierrez for valuable guidance. And also by the Laboratory Baldacci free supply of medication "Magnesium Pidolate and Placebo"

### Potential Conflict of Interest

No potential conflict of interest relevant to this article was reported.

### Sources of Funding

There were no external funding sources for this study.

### Study Association

This article is part of the thesis of doctoral submitted by Cristina Nádja Muniz Lima de Falco, from Faculdade de Medicina Instituto do Coração Universidade de São Paulo.

### Erratum

The version of "Successful Improvement of Frequency and Symptoms of Premature Complexes after Oral Magnesium Administration" published as ahead of print by Arquivos Brasileiros de Cardiologia underwent the following modification as required by the Editor on 10/11/2012: In the Methods section of the Abstract, PsVC/h was replaced by PsVC.

## References

1. Barret PA, Peter CT, Swan HJ, Singh BN, Mandel WJ. The frequency and prognostic significance of electrocardiographic abnormalities in clinically normal individuals. *Prog Cardiovasc Dis.* 1981;23(4):299-319.
2. DePaula RS, Antelmi I, Vincenzi MA, André CD, Artes R, Grupi CJ, et al. Cardiac arrhythmias and atrioventricular block in a cohort of asymptomatic individuals without heart disease. *Cardiology.* 2007;108(2):111-6.
3. Rasmussen V, Jensen G, Schnohr P, Hansen JF. Premature ventricular beats in healthy adult subjects 20 to 79 years of age. *Eur Heart J.* 1985;6(4):335-41.
4. Wajngarten M, Grupi C, Bellotti GM, Da Luz PL, Azul LG, Pileggi F. Frequency and significance of cardiac rhythm disturbances in healthy elderly individuals. *J Electrocardiol.* 1990;23(2):171-6.
5. Hiss RG, Averill KH, Lamb LE. Electrocardiographic findings in 67,375 asymptomatic subjects. III. Ventricular rhythms. *Am J Cardiol.* 1960;6:96-107.
6. Darrieux FC, Scanavacca MI, Hachul DT, Melo SL, D'Ávila AB, Gruppi CJ, et al. Ablação com radiofrequência de extra-sístoles da via de saída do ventrículo direito. *Arq Bras Cardiol.* 2007;88(3):265-72.
7. Huskisson E, Maggini S, Ruf M. The role of vitamins and minerals in energy metabolism and well-being. *J Int Med Res.* 2007;35(3):277-89.
8. Swaminathan R. Magnesium metabolism and its disorders. *Clin Biochem Rev.* 2003;24(2):47-66.
9. Wu J, Carter A. Magnesium: the forgotten electrolyte. *Aust Prescr.* 2007;30:102-5.
10. Ebel H, Gunther TJ. Magnesium metabolism: a review. *Clin Chem Clin Biochem.* 1980;18(5):257-70.
11. Elin RJ. Laboratory tests for the assessment of magnesium status in humans. *Magn Trace Elem.* 1991-1992;10(2-4):172-81.
12. Arnaud MJ. Update on the assessment of magnesium status. *Br J Nutr.* 2008;99(Suppl.3):S24-36.
13. Marier JR. Magnesium content of the food supply in the modern-day world. *Magnesium.* 1986;5(1):1-8.
14. Gums JG. Magnesium in cardiovascular and other disorders. *Am J Health Syst Pharm.* 2004;61(15):1569-76.
15. Kirkwood BR, Sterne JA. *Essential medical statistics.* 2<sup>nd</sup> ed. Massachusetts, USA: Wiley Blackwell; 2003.
16. Neter J, Kuntner MH, Nachtsheim CJ, Wasserman W. *Applied linear statistical models.* 4th ed. Illinois: McGraw Hill/Irwing; 1996.
17. Zehender M, Meinertz T, Faber T, Caspary A, Jeron A, Bremm K, et al. Antiarrhythmic effects of increasing the daily intake of magnesium and potassium in patients with frequent ventricular arrhythmias. Magnesium in Cardiac Arrhythmias (MAGICA) Investigators. *J Am Coll Cardiol.* 1997;29(5):1028-34.
18. Silver BB. Development of cellular magnesium nano-analysis in treatment of clinical magnesium deficiency. *J Am Coll Nutr.* 2004;23(6):732S-7S.
19. Kawahara K, Sato R, Iwabuchi S, Matsuyama D. Rhythmic fluctuations in the concentration of intracellular Mg<sup>2+</sup> in association with spontaneous rhythmic contraction in cultured cardiac myocytes. *Chronobiol Int.* 2008;25(6):868-81.
20. Chubanov V, Waldegger S, Mederos y Schnitzler M, Vitzthum H, Sassen MC, Seyberth HW, et al. Disruption of TRPM6/TRPM7 complex formation by a mutation in the TRPM6 gene causes hypomagnesemia with secondary hypocalcemia. *Proc Natl Acad Sci U S A.* 2004;101(9):2894-9.
21. Voets T, Nilius B, Hoefs S, van der Kemp AW, Droogmans G, Bindels RJ, et al. TRPM6 forms the Mg<sup>2+</sup> influx channel involved in intestinal and renal Mg<sup>2+</sup> absorption. *J Biol Chem.* 2004;279(1):19-25.
22. Belin RJ, He K. Magnesium physiology and pathogenic mechanisms that contribute to the development of the metabolic syndrome. *Magn Res.* 2007;20(2):107-29.
23. He K, Liu K, Daviglus ML, Morris SJ, Loria CM, Van Horn L, et al. Magnesium intake and incidence of metabolic syndrome among young adults. *Circulation.* 2006;113(13):1675-82.
24. Wirell MP, Wester PO, Stegmayr BC. Nutritional dose of magnesium in hypertensive patients on beta blockers lowers systolic blood pressure: a double-blind, cross-over study. *J Intern Med.* 1994;236(2):189-95.
25. Tong GM, Rude RK. Magnesium deficiency in critical illness. *J Intensive Care Med.* 2005;20(1):3-17.
26. Iseri LT, Allen BJ, Brodsky MA. Terapia de magnesio en arritmias cardiacas en medicina de cuidados críticos. *Revista de Cuidados Intensivo en Medicina.* 2005;20(1):3-17.
27. Franz KB. A functional biological marker is needed for diagnosing magnesium deficiency. *J Am Coll Nutr.* 2004;23(6):738S-41S.
28. Moe SM. Disorders of calcium, phosphorus, and magnesium. *Am J Kidney Dis.* 2005;45(1):213-8.