

Two New Oleanane Saponins from *Chiococca alba* (L.) Hitch.

Ricardo M. Borges,*^a Luzineide W. Tinoco,^a José D. de Souza Filho,^b
Nancy dos Santos Barbi^c and Antonio Jorge R. da Silva^a

^aNúcleo de Pesquisas de Produtos Naturais and ^cFaculdade de Farmácia, Centro de Ciências da Saúde,
Universidade Federal do Rio de Janeiro, 21941-590 Rio de Janeiro-RJ, Brazil

^bDepartamento de Química, Instituto de Ciências Exatas, Universidade Federal de Minas Gerais,
31270-901 Belo Horizonte-MG, Brazil

Duas saponinas triterpênicas foram isoladas do extrato etanólico das raízes de *Chiococca alba* (L.) Hitch. (Rubiaceae). Suas estruturas foram elucidadas por RMN de ¹H e ¹³C 1D e 2D e por espectrometria de massas de alta resolução com ionização por eletronebulização como 3-*O*-β-D-glucopiranosil-3-β-hidroxiolean-12,15-dien-28-oato de *O*-α-D-apiofuranosil (1→3)-[α-D-apiofuranosil (1→4)]-α-L-rhamnopiranosil (1→2)-α-L-arabinopiranosila (**1**) e 3-*O*-β-D-glucopiranosil-3-β-hidroxiolean-12,15-dien-28-oato de 28-*O*-α-D-apiofuranosil (1→3)-α-L-rhamnopiranosil (1→2)-α-L-arabinopiranosila (**2**).

Two triterpene saponins were isolated from the ethanolic extract of the roots of *Chiococca alba* (L.) Hitch (Rubiaceae). Their structures were determined by ¹H and ¹³C 1D and 2D NMR and high resolution electrospray mass spectrometry as 3-*O*-β-D-glucopyranosyl-3β-hydroxyolean-12,15-dien-28-oic acid 28-*O*-α-D-apiofuranosyl (1→3)-[α-D-apiofuranosyl (1→4)]-α-L-rhamnopyranosyl (1→2)-α-L-arabinopyranosyl ester (**1**) and 3-*O*-β-D-glucopyranosyl-3β-hydroxyolean-12,15-dien-28-oic acid 28-*O*-α-D-apiofuranosyl (1→3)-α-L-rhamnopyranosyl (1→2)-α-L-arabinopyranosyl ester (**2**).

Keywords: *Chiococca alba*, cainca, Rubiaceae, triterpene saponins

Introduction

Chiococca alba (L.) Hitch. (Rubiaceae), commonly known as “cainca” in Brazil, is a tropical and sub-tropical shrub spread all over the American continent. Its root bark is used in Brazilian traditional medicine for the treatment of several illnesses.¹ Toxicological studies demonstrated absence of mutagenic activity in the ethanolic extract of the roots of *C. alba* and also low acute and sub acute toxicity by the oral route. Nevertheless, toxicity by parenteral routes was indeed pronounced.²

Previous works on *C. alba* reported the occurrence of lignans, coumarins, ketoalcohols,³ triterpenes,⁴ iridoids,⁵ flavonoids,⁶ two quinoline alkaloids⁷ and a nor-*seco*-pimarane.⁸

The present paper reports for the first time the isolation and structure elucidation of two triterpenoidal saponins from the ethanolic extract of the roots of *C. alba*.

Experimental

General experimental procedures

Optical rotations were measured on a Perkin Elmer model 243B polarimeter. Infrared spectra were recorded on a Nicolet Magna spectrometer. ¹H NMR and ¹³C NMR spectra were recorded at 200 and 50 MHz (Varian Gemini 200), at 400 and 100 MHz (Bruker DRX 400) and at 600 and 150 MHz (Bruker DRX 600) in methanol-*d*₄ and chloroform-*d*₁ with TMS as internal standard.

The chemical shifts are reported in ppm. 1D and 2D experiments were made with the standard software provided. HRES/ESI/TOF MS analyses were made with a Bruker Daltonics microTOF instrument. Column chromatography: ODS silica gel (Aldrich Chemical Company, Inc); TLC: silica gel (0.25 mm pre-coated plates 60 F₂₅₄, Merck). For preparative HPLC a 7.6 × 300 mm ODS column and UV detection at 210 nm were used. GC-MS analyses were recorded on a Shimadzu QP5000

*e-mail: rmborges@nppn.ufjf.br

GC/MS instrument (5% phenylmethyl silicone column, 30 m × 0.25 mm ID, 0.25 μm film thickness; programmed column temperature from 110 to 290 °C, 5 °C min⁻¹).

Plant material

Samples of *C. alba* were collected in Nova Friburgo, Rio de Janeiro, Brazil. The botanical identification was provided by Dr. Sebastião Neto, and a voucher specimen (RB395399) is deposited in the Herbarium of Rio de Janeiro Botanical Garden.

Extraction and isolation


Air-dried and powdered roots of *C. alba* (400 g) were extracted with ethanol. The extract was evaporated; the residue obtained (12 g) was suspended in water and successively partitioned with methylene chloride and butanol. The butanol fractions were combined, evaporated and the residue (4 g) was suspended in methanol and subjected to controlled precipitation with diethyl ether. The precipitate (2 g) was fractionated by column chromatography (octadecylsilane, 60 × 20 cm) using H₂O with increasing proportions of methanol (0% to 100%) to afford 10 fractions. TLC tests carried out with Liebermann-Bouchard and sulfuric orcinol reagents together with the observation the abundant foam formation, allowed the identification of the saponin enriched fractions. Further purification was carried out with reversed phase (octadecylsilane) preparative HPLC using methanol: 0.02% aqueous trifluoroacetic acid (60:40; v/v) to afford chiococcasaponin I (78 mg) and chiococcasaponin II (48 mg).

Chiococcasaponin I (1)


Amorphous white solid, UV transparent, $[\alpha]_D^{25} = -58$ (MeOH; *c* 1). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$: 3428, 2944, 1732, 1040, 986. ¹H NMR (methanol-*d*₄, 400 MHz) and ¹³C NMR (methanol-*d*₄, 100 MHz) (Table 1). HRES/ESI/TOF MS (negative mode) calculated for C₅₇H₈₈O₂₅, [M-H]⁻: *m/z* 1171.5536; found: 1171.5546. HRES/ESI/TOF MS (positive mode), calculated for C₅₇H₈₈O₂₅Na, [M+Na]⁺: *m/z* 1195.5512; found: 1195.5513.

Chiococcasaponin II (2)

Amorphous white solid, UV transparent, $[\alpha]_D^{25} = -46$ (MeOH; *c* 1). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$: 3426, 2946, 1736, 1137, 1077, 1053, 1034, 980. ¹H NMR (methanol-*d*₄, 400 MHz) and ¹³C NMR (methanol-*d*₄, 100MHz) (Table 1). HRES/ESI/TOF MS (negative mode) calculated for C₅₂H₈₀O₂₁, [M-H]⁻: *m/z* 1039.5114; found: 1039.5122. HRES/ESI/TOF MS (positive mode), calculated for C₅₂H₈₀O₂₁Na, [M+Na]⁺: *m/z* 1063.5089 found: 1063.5084.



1 Chiococcasaponin I



2 Chiococcasaponin II

Acid hydrolysis

A solution of each saponin (5 mg per 2 mL of 0.5 mol L⁻¹ aqueous TFA) was poured in a Kontes heavy wall pressure resistant vessel and heated for 2 min in a microwave oven at full power. The reaction mixture was cooled, diluted with water and extracted with diethyl ether. The organic layer was concentrated under vacuum and the residue was methylated with diazomethane. 3β-hydroxyolean-12, 15-dien-28-oate methyl ester was identified by MS and NMR as the aglycone for both saponins.

MS (EI, 70eV) *m/z* 468 (2%); 260 (42%); 245 (14%); 231; 207; 201 (100%); 187 (14%); 173; 157; 145; 131 (46%); 119; 105; 95; 81; 69; 55. ¹³C NMR (chloroform-*d*₇, 100 MHz): 16.1 (C-25); 17.0 (C-23); 18.3 (C-26); 19.6 (C-6); 23.6 (C-30); 24.4 (C-11); 25.1 (C-27); 27.1 (C-2); 28.5 (C-24); 31.3 (C-20); 33.8 (C-22); 33.9 (C-29); 33.9 (C-17); 34.1 (C-7); 35.7 (C-21); 38.2 (C-10); 39.8 (C-1); 40.3 (C-4); 41.3 (C-8); 43.5 (C-18); 44.6 (C-19); 45.6 (C-14); 48.7 (C-9); 57.2 (C-5); 91.2 (C-3); 124.7 (C-12); 128.6 (C-15); 137.8 (C-16); 141.5 (C-13); 176.1 (C-28).

The aqueous phase was lyophilized, the monosaccharides were reduced with sodium borohydride in water and the

Table 1. Carbon and proton NMR data for Chiococcasaponins I and II (100 and 400 MHz, CD₃OD, internal TMS)

Carbon	Chiococcasaponin I		Chiococcasaponin II	
	¹ H	¹³ C	¹ H	¹³ C
1 CH ₂	1.60 / 1.01	39.7	1.60 / 1.01	39.8
2 CH ₂	1.87 / 1.7	27.1	1.85 / 1.68	27.1
3 CHOH	3.17 (<i>dd, J</i> 11.1 / 3.5 Hz)	91.1	3.17 (<i>dd, J</i> 11.1 / 3.5 Hz)	91.2
4 C	--	40.2	--	40.3
5 CH	0.82	57.0	0.82	57.2
6 CH ₂	1.45	19.4	1.60	19.6
7 CH ₂	1.68 / 1.54	33.8	1.60 / 1.50	34.1
8 C	--	41.2	--	41.3
9 CH	1.66	48.7	1.65	48.7
10 C	--	38.2	--	38.2
11 CH ₂	1.97 / 1.86	24.3	1.95 / 1.84	24.4
12 CH	5.44	124.6	5.46	124.7
13 C	--	141.6	--	141.5
14 C	--	45.5	--	45.6
15 CH	5.54	128.7	5.54	128.6
16 CH	5.67	137.5	5.66	137.8
17 C	--	33.6	--	33.9
18 CH	3.92 (<i>dd, J</i> 13.1 / 2.78 Hz)	43.4	2.93 (<i>dd, J</i> 13.1 / 2.78 Hz)	43.5
19 CH ₂	1.58 / 1.10	44.6	1.56 / 1.09	44.6
20 C	--	31.3	--	31.3
21 CH ₂	1.26	35.6	1.26	35.7
22 CH ₂	1.81	33.6	1.80	33.8
23 CH ₃	0.86 (<i>s</i>)	16.9	0.85 (<i>s</i>)	17.0
24 CH ₃	1.06 (<i>s</i>)	28.4	1.06 (<i>s</i>)	28.5
25 CH ₃	0.88 (<i>s</i>)	16.0	0.88 (<i>s</i>)	16.1
26 CH ₃	0.64 (<i>s</i>)	18.3	0.63 (<i>s</i>)	18.3
27 CH ₃	1.17 (<i>s</i>)	25.0	1.16 (<i>s</i>)	25.1
28 C=O	--	176.1	--	176.1
29 CH ₃	0.90 (<i>s</i>)	33.6	0.90 (<i>s</i>)	33.9
30 CH ₃	0.93 (<i>s</i>)	23.5	0.93 (<i>s</i>)	23.6
Ara at C-28				
1 CH	5.39 (<i>d, J</i> 4.4 Hz)	94.4	5.46 (<i>d, J</i> 5.4 Hz)	94.4
2 CH	3.80	75.4	3.78	75.4
3 CH	3.51	72.5	3.76	72.5
4 CH	3.81	68.1	3.82	67.4
5 CH ₂	3.56/3.86	64.9	5.51/3.88	64.0
Rha				
1 CH	5.12 (<i>s</i>)	101.4	5.06 (<i>s</i>)	101.7
2 CH	3.95	72.3	3.79	72.7
3 CH	3.76	80.8	3.86	71.7
4 CH	3.62	79.2	3.52	80.2
5 CH	3.83	69.0	3.75	69.2
6 CH ₂	1.27 (<i>d, J</i> 6 Hz)	18.5	1.28 (<i>d, J</i> 6 Hz)	18.5
Api				
1 CH	5.19 (<i>d, J</i> 1.7 Hz)	112.5	5.33 (<i>d, J</i> 1.7 Hz)	111.6
2 CH	4.04	79.3	3.92	78.6
3 C	--	80.7	--	80.6
4 CH ₂	3.76/4.05	74.9	3.76/4.00	75.0
5 CH ₂	3.59	65.3	3.58	65.5
Api				
1 CH	5.29 (<i>d, J</i> 1.7 Hz)	112.2	--	--
2 CH	3.95	78.3	--	--
3 C	--	80.4	--	--
4 CH ₂	3.77/4.05	75.1	--	--
5 CH ₂	3.56	65.5	--	--
GlcUA at C-3				
1 CH	4.38 (<i>d, J</i> 7.3 Hz)	107.0	4.38 (<i>d, J</i> 7.3 Hz)	107.0
2 CH	3.24	75.3	3.24	75.5
3 CH	3.37	77.7	3.37	77.9
4 CH	3.51	73.3	3.51	73.4
5 CH	3.76	75.4	3.79	75.4
6 C(OH)=O	--	176.0	--	176.0

resulting alditols were acetylated (acetic anhydride/pyridine, 1 h). The alditol acetates of arabinose, rhamnose and apiose were identified by direct comparison with authentic samples using GC/MS. The monosaccharides were also characterized by TLC on silica gel (butanol/acetic acid/water, 2:1:1). The spots on TLC plates were visualized by spraying with sulphuric-orsinol solution followed by heating. Glucuronic acid, arabinose and rhamnose were identified by comparison with authentic samples.

Results and Discussion

The HRES/ESI MS (positive mode) of chiococcasaponin I (**1**) exhibited a pseudomolecular ion peak [M+Na]⁺ at *m/z* 1195.5513 and the negative mode HRES/ESI-MS exhibited a pseudomolecular ion peak [M-H]⁻ at *m/z* 1171.5546, both peaks being consistent with the molecular formula C₅₇H₈₈O₂₅. Acid hydrolysis of **1** afforded the aglycone and the following monosaccharides: glucuronic acid, rhamnose, arabinose and apiose which were identified by GC/MS and TLC analysis. Due to the fact that apiose is a branched-chain sugar, incomplete acetylation is often observed for it when standard alditol acetylation conditions are utilized. This leads to the observation of two peaks for the partially acetylated apiose alditols in the GC/MS analyses. This behavior has already been described by Harris.⁹ The aglycone had its structure confirmed as 3β-hydroxyolean-12, 15-dien-28-oic acid by comparison of its ¹H and ¹³C NMR spectra with literature data.⁴ This is the first report on the occurrence of a 15,16 unsaturated oleanolic acid saponin. ¹H and ¹³C NMR data are shown in Table 1. 1D ¹H and ¹³C NMR, DEPT and HSQC spectra confirmed the presence of five sugar residues corresponding to anomeric protons/carbons at 4.38/107.0, 5.12/101.4, 5.39/94.4, 5.19/112.5 and 5.29/112.2 ppm respectively. The anomeric proton at 4.38 ppm (*d, J* 7.3 Hz) displayed an array of correlations with signals at 3.24, 3.37, 3.51 and 3.76 ppm in the TOCSY spectrum. These ¹H chemical shifts and the respective correlated carbons (HSQC) at: 75.3, 77.7, 73.3, 75.4 ppm together with the carboxyl group appearing at 176.0 ppm are consistent with the presence of a β-glucopyranosyl residue. A further correlation, observed in the HMBC spectrum between the proton at 4.38 and the carbon at 91.1 ppm (C-3) confirmed the location of the β-glucopyranosyl residue at C-3. Another correlation, between protons at 4.38 and 3.17 (aglycone H-3), observed in the T-ROESY spectrum reinforced the above deduction. The configurations at the anomeric carbons were assigned with the help of a NMR *J*-resolved heteronuclear experiment, on the basis of the measured ¹J_{C,H}: 164 Hz for the signal at 94.4 ppm (C-1, Ara), 170.6 Hz for the signal

at 101.4 ppm (C-1, Rha), 157 Hz for the signal at 107.0 ppm (C-1, GlcUA), 169.8 Hz for the signal at 112.2 ppm (C-1, Api) and 169.8 Hz for the signal at 112.5 ppm (C-1, Api).^{10, 11}

The individual spin systems for the other monosaccharide residues were assigned with the aid of the correlations between the anomeric protons (or methyl signal, for rhamnose) and the remaining protons in 2D TOCSY and COSY spectra. The chemical shifts of the apiofuranosyl residues were identified on the basis of C-3 being quaternary carbons (80.7, 80.4 ppm) and C-4 and C-5 being two methylenes. The assignments were double checked by comparison with literature chemical shift data and from HSQC and HMBC correlations. The sequence of the oligosaccharide chain was determined by a combination of DEPT, HSQC, HMQC, HMBC and T-ROESY experiments. Thus, the correlation observed in the HMBC spectrum between signals at 5.12 (H-1 rhamnose) and 75.4 ppm (C-2 arabinose) indicated the linkage of the rhamnose residue to the position 2 of the arabinose residue. A correlation between signals at 5.12 (H-1 rhamnose) and 3.80 (H-2 arabinose) in the T-ROESY spectrum confirm this glycosidic linkage. The connection of the two terminal apiofuranose residues to positions 3 and 4 of the rhamnose unit was confirmed by the following HMBC correlations: 3.62 ppm (H-4 rhamnose) to 112.2 ppm (C-1, apiofuranose); 3.76 ppm (H-3 rhamnose) to 112.5 (C-1, apiofuranose). Finally, the HMBC correlation between the proton at 5.39 (H-1, arabinose) and C-28 confirms the location of the tetrasaccharide. Thus, Chiococcasaponin I (**1**) is a typical GOTCAB (glucuronide oleanane triterpenic type carboxylic acid 3,28-O-bisdesmoside) according to the classification of Tan *et al.*¹¹

The HR-ESI-MS (positive mode) of chiococcasaponin II (**2**) exhibited a pseudomolecular ion peak $[M+Na]^+$ at m/z 1063.5084 and in the negative mode exhibited a pseudomolecular ion peak $[M-H]^-$ at m/z 1039.5122, both peaks being consistent with the molecular formula $C_{52}H_{80}O_{21}$. Acid hydrolysis of **2** afforded the aglycone 3 β -hydroxyolean-12, 15-dien-28-oic acid and the following monosaccharides: glucuronic acid, rhamnose, arabinose and apiose which were identified by GC/MS and TLC analysis. The hydrolysis reaction suggested the presence of the same sugar residues and aglycon as in **1**. However, a comparison of the ¹³C NMR, HSQC, HMBC and MS data indicated a difference between the sugar chain structures **1** and **2**. Thus, the lack of five signals in the ¹³C NMR

spectrum and the anomeric signal at 5.29 ppm in the ¹H NMR of **2** together with the 132 mass units of difference between Chiococcasaponin I (**1**) and Chiococcasaponin II (**2**) clearly indicates the absence of an apiofuranosyl residue in **2**. It is worth of mention the upfield shift (*ca.* -7 ppm) observed for C-3 in the rhamnose residue of **2** denoting absence of substitution at this site.

Acknowledgments

R.M. Borges acknowledges CNPq for the fellowship. The authors acknowledge Centro Nacional de Ressonância Magnética Nuclear de Macromoleculas Jiri Jonas, IBqM/UFRJ for 400 and 600 MHz NMR spectra.

Supplementary Information

Supplementary data are available free of charge at <http://jbcs.sbq.org.br>, as PDF file.

References

1. Mors, W. B.; Rizzini, C. T.; Pereira, N. A.; *Medicinal Plants of Brazil*, Reference Publications Inc.: Algonac Michigan, 2000.
2. Gazda, V. E.; Gomes-Carneiro, M. R.; Barbi, N. S.; Paumgarten, F. J. R.; *J. Ethnopharmacol.* **2006**, *105*, 187.
3. Abd El-Hafiz, M. A.; Weniger, B.; Quiron, J. C.; Anton, R.; *Phytochemistry* **1991**, *30*, 2029.
4. Bhattacharyya, J.; Cunha, E. V. L.; *Phytochemistry* **1992**, *31*, 2546.
5. Carbonezi, C. A.; Martins, D.; Young, M. C. M.; Lopes, M. N.; Furlan, M.; Rodrigues-Filho, E.; Bolzani, V. S.; *Phytochemistry* **1999**, *51*, 781.
6. Lopes, M. N.; de Oliveira, A. C.; Young, M. C. M.; Bolzani, V. S.; *J. Braz. Chem. Soc.* **2004**, *15*, 468.
7. El-Abadi, N.; Weniger, B.; Lobstein, A.; Quirion, J. C.; Anton, R.; *Planta Med.* **1989**, *55*, 603.
8. Borges-Argáez, R.; Medina Baizabal, L.; May Pat, F.; Waterman, P. G.; Pena-Rodriguez, L. M.; *J. Nat. Prod.* **2001**, *64*, 228.
9. Harris, P. J.; *Carbohydr. Res.* **1992**, *227*, 365.
10. Agrawal, P. K.; Pathak, A. K.; *Phytochem. Anal.* **1996**, *7*, 113.
11. Tan, N.; Zhou, J.; Zhao, S.; *Phytochemistry* **1999**, *52*, 153.

Received: October 14, 2008

Web Release Date: September 25, 2009

Two New Oleanane Saponins from *Chiococca alba* (L.) Hitch.

Ricardo M. Borges,*^a Luzineide W. Tinoco,^a José D. de Souza Filho,^b
Nancy dos Santos Barbi^c and Antonio Jorge R. da Silva^a

^aNúcleo de Pesquisas de Produtos Naturais and ^cFaculdade de Farmácia, Centro de Ciências da Saúde, Universidade Federal do Rio de Janeiro, 21941-590 Rio de Janeiro-RJ, Brazil

^bDepartamento de Química, Instituto de Ciências Exatas, Universidade Federal de Minas Gerais 31270-901 Belo Horizonte-MG, Brazil




Figure S1. ¹³C NMR of chiococcasaponin I.

*e-mail: rmborges@nppn.ufrj.br




Figure S2. ^1H NMR of chiococcasaponin I.




Figure S3. Edited HSQC of chiococcasaponin I.




Figure S4. COSY of chiococcasaponin I.




Figure S5. TOCSY of chiococcasaponin I.




Figure S6. HMBC of chiococcasaponin I.




Figure S7. T-ROESY of chiococcasaponin I.




Figure S8. Mass spectrometry of Chiococcasaponin I (positive mode).





Figure S9. Mass spectrometry of Chiococasaponin I (negative mode).

Chiococcasaponin II:

**Figure S10.** ¹³C NMR of chiococcasaponin II.




Figure S11. ¹H NMR of chiococcasaponin II.




Figure S12. Edited HSQC of chiococcasaponin II.




Figure S13. COSY of chiococcasaponin II.




Figure S14. TOCSY of chiococcasaponin II.




Figure S15. HMBC of chiococcasaponin II.




Figure S16. Mass spectrometry of Chiococcasaponin II (positive mode).




Figure S17. Mass spectrometry of Chiococcasaponin II (negative mode).