

The role of phytochromes in cadmium stress responses in tomato

Lucas Aparecido Gaion^{1*}, Paulo Guilherme Lorevice¹, Carolina Cristina Monteiro¹, Marina Alves Gavassi¹, Víctor D'Amico-Damião², Priscila Lupino Gratão¹, Eduardo Custódio Gasparino¹, Rogério Falleiros Carvalho¹

1.Universidade Estadual Paulista “Júlio de Mesquita Filho” - Faculdade de Ciências Agrárias e Veterinárias - Departamento de Biologia Aplicada à Agropecuária - Jaboticabal (SP), Brazil.

2.Universidade Estadual Paulista “Júlio de Mesquita Filho” - Faculdade de Ciências Agrárias e Veterinárias - Departamento de Produção Vegetal - Jaboticabal (SP), Brazil.

ABSTRACT: It is well known that phytochromes mediate a wide range of photomorphogenic processes in plants. In addition, many studies have demonstrated the involvement of phytochromes as part of abiotic stress signaling responses. However, little is known about cadmium (Cd) stress regulation by phytochromes. Thus, in this study, we used the *phyA* (*far red-insensitive; fri*), *phyB1* (*temporary red-insensitive; tri*) and *phyB2* (*phyB2*) tomato (*Solanum lycopersicum* L.) mutants to investigate the roles of these three phytochromes on Cd stress responses. The plants were grown over a 21-d period in the presence of Cd. We evaluated plant growth, Cd and chlorophyll

content and anatomical changes in the leaves. The results indicated that all genotypes were affected by Cd and showed reduced growth of the shoots and roots, as well as reduced chlorophyll content. The accumulation of Cd was similar for all genotypes, and a higher Cd content was found in roots. Anatomical analysis of the vascular bundles revealed that *fri* and *tri* seem to be more disrupted by Cd. Overall, these results indicate that phytochromes do not determine Cd stress tolerance in tomato plants.

Key words: abiotic stress, heavy metal, mutants, plant anatomy, *Solanum lycopersicum* L.

*Corresponding author: lucas.gaion@yahoo.com.br

Received: Sept. 30, 2016 – Accepted: Fev. 28, 2017



INTRODUCTION

Plants possess photoreceptors that compose sophisticated light-sensing mechanisms that are localized inside and outside the chloroplasts and the nucleus (Hiltbrunner et al. 2005). Among photoreceptors, the phytochrome family stands out as sensors for red (R; ~ 660 nm) and far-red (FR; ~ 730 nm) light (Essen et al. 2008; Yang et al. 2008). The phytochrome holoprotein consists of an apoprotein with a covalently linked linear tetrapyrrole chromophore, and this complex regulates the expression of a large number of light-responsive genes and thus influences many photomorphogenic events (Neff et al. 2000; Quail 2002a,b; Franklin and Quail 2010). In plants, a small family of PHY genes encodes phytochromes. For example, five phytochrome genes are known in *Arabidopsis thaliana* (*PHYA* to *PHYE*) and in tomato (*PHYA*, *PHYB1*, *PHYB2*, *PHYE* and *PHYF*), and three *PHY* genes are known in rice (*Oryza sativa*) (*PHYA* to *PHYC*) (Bae and Choi 2008).

The involvement of phytochromes in controlling plant growth and development is well documented (Franklin 2016; Wit et al. 2016). Additionally, these photoreceptors have been shown to act as key modulators of both biotic and abiotic stresses (Carvalho et al. 2011a; Zao et al. 2014; D'Amico-Damião et al. 2015). These reports have led to a series of studies exploring the molecular and biochemical basis by which phytochromes mediate stresses, such as drought (D'Amico-Damião et al. 2015), salinity (Balestrasse et al. 2008a), high light (Boccalandro et al. 2001) or heavy metals (Cui et al. 2011). For instance, studies of *A. thaliana* under cold conditions demonstrated that light signaling by phytochrome B is involved in plant responses to cold via the expression of C-repeat binding factor (*CBF*) genes (Kim et al. 2002) and to biotic stress via the activation of lipoxygenase (Zao et al. 2014). In addition, rice phytochrome B-deficient mutant (*phyB*) plants were more tolerant to drought than the wild type. These plants exhibited reduced stomatal density and length, resulting in a decreased transpiration rate that enhanced drought tolerance (Liu et al. 2012). Similarly, the *phyB*-deficient mutant of tomato exhibited increased drought tolerance after five days without irrigation (D'Amico-Damião et al. 2015).

Another relevant type of abiotic stress experienced by plants under certain conditions is heavy metals. The increase in the cellular heavy metals are one of the main classes of

abiotic stress agents for living organisms because of their high level of bioaccumulation and toxicity (Verkleij et al. 2009; Nagajyoti et al. 2010). The increase in heavy metal concentrations in the cellular environment can disturb several signaling pathways and cause irreversible damage to biological systems (Verkleij et al. 2009; Rossato et al. 2011). Among heavy metals, Cd represents one of the most damaging metals to plants. At high concentrations, this heavy metal can generate reactive oxygen species (ROS) such as superoxide radicals (O_2^-), singlet oxygen (1O_2), hydrogen peroxide (H_2O_2) and hydroxyl radicals (OH^-), which can disrupt the plant defense system (Gratão et al. 2012; Gratão et al. 2015).

Shen et al. (2011) demonstrated that the overexpression of *BnHO-1*, which is involved in the biosynthesis of heme oxygenase (HO), a key enzyme in the biosynthesis of the chromophore, provided tolerance to mercury (Hg) in *Brassica napus*, indicating that phytochromes are involved in metal-induced signaling pathways. Additionally, Jin et al. (2013) reported involvement of HO-1 in the alleviation of Cd toxicity in root tissues of alfalfa (*Medicago sativa*). Likewise, soybean (*Glycine max* L.) and alfalfa plants exposed to Cd-stressful conditions exhibited the upregulation of heme oxygenase-1 (HO-1) (Balestrasse et al. 2008b; Cui et al. 2011).

However, direct evidence for the involvement of phytochromes with plant responses to heavy metals are still missing. Therefore, in order to investigate whether *phyA* and *phyB* regulate events in Cd-stressful conditions, we assessed phenotypical and anatomical changes in *fri*, *tri* and *phyB2* tomato mutants exposed to elevated cadmium concentrations.

MATERIAL AND METHODS

Plant material and growth conditions

Seeds of the tomato (*S. lycopersicum* L.) mutants *fri* (*far red-insensitive*), *tri* (*temporary red-insensitive*) and *phyB2*, and their wild-type counterpart (WT) (cv. Moneymarker, MM) were germinated in boxes containing a mixture of 1:1 (by volume) commercial potting mix (Plantmax HT Eucatex, Brazil) and vermiculite. To avoid premature plant death by cadmium, seven-day-old plants were transferred to hydroponic conditions with Hoagland's nutrient solution initially containing 50 μM $CdCl_2$. Fourteen days after

→

sowing, the plants received 100 $\mu\text{M}\cdot\text{CdCl}_2$ for seven more days for a total of a 21-day period. Control plants were grown in Hoagland's solution without CdCl_2 . The nutrient solution with or without CdCl_2 was changed weekly, and the pH was maintained at approximately 5.6. The plants were maintained in a greenhouse with an average mean temperature of 27 °C under a 12 h photoperiod.

Cadmium content

Assays for the amount of Cd in the plant tissues were performed following digestion with a mixture of nitric and perchloric acids, according to Malavolta et al. (2011). Cd concentration was measured using flame atomic absorption spectroscopy with a Perkin Elmer spectrometer model 310. Root to shoot Cd transport of tomato genotypes was assessed using a Cd translocation index (TI), which was calculated as the percentage of the total Cd accumulation that was translocated to the shoots (Usman et al. 2012).

Growth analysis

The leaf area was measured using an Image Analysis System (Delta-T Devices, Cambridge, UK). Root area, diameter, length and density were all recorded using a scanner (Hewlett Packard 5c), and the image of each plant was analyzed by Delta-T Scan software. Subsequently, the plants were weighted for fresh mass. Next, roots and shoots were oven-dried at 60 °C for 72 h. The dry weight of roots and shoots was determined using an analytical balance (Adventurer; Ohaus, Shangai, P. R. China).

Anatomical analysis

A series of anatomical analysis were performed by sampling a middle portion of the leaf blade (3 × 3 mm) from selected leaves at the third node in four replications. The samples were fixed in FAA 50 (formaldehyde + acetic acid + 50% alcohol) for approximately 48 h and stored in 70% alcohol, according to Johansen methods (1940).

Permanent slides were prepared using the traditional methods of ethyl dehydration (85, 95 and 100%) for two hours at each concentration and were embedded in paraffin (Johansen 1940). The material was cut using a precision microtome (Leica RM2065) into 10 mm thick slices with disposable steel razors, yielding a transverse

leaf series. The sections were stained with 0.05% toluidine blue (O'Brien et al. 1964) and assembled using synthetic Canada balsam.

The leaf anatomical structures were observed using 10 × and 40 × magnification of a binocular optical microscope (Bel photonics, Bio2 SSI model) and measured using a micrometric objective. The photomicrographs were taken with a trinocular optical microscope (Bel photonics Solaris) coupled with a USB digital camera (5.0 MP Bel). The integrity of the leaf tissue, the thickness of the leaf and palisade, the spongy parenchyma tissues and the diameter of the vascular bundles were evaluated.

Chlorophyll content

Leaf chlorophyll content was determined using a Minolta SPAD-502 m, which measures leaf transmittance at two wavelengths: red (approximately 660 nm) and near-infrared (approximately 940 nm). SPAD readings were taken from the terminal leaf of the fourth leaf from the base of the shoot. The SPAD sensor was randomly placed only on the leaf mesophyll tissue, avoiding the veins.

Chlorophyll fluorescence measurement

The fluorescence emission of chlorophyll (Fv/Fm) was measured using a pulse amplitude modulated fluorometer (FMS2, Hansatech Instruments, Pentney King's Lynn, U.K.). The leaves were exposed to darkness for fifteen minutes and were handled with tweezers before fluorescence estimation, as described by Saijo et al. (2000).

Statistical analysis

The experimental design was randomized with twelve plants from three replicate pots, and the results were expressed as the mean and standard error of the mean (\pm SEM). The means were compared using Tukey's test ($p \leq 0.05$).

RESULTS AND DISCUSSION

After 21 days of experiment, Cd accumulation exhibited a similar pattern in all investigated genotypes (Figure 1a). In non-stressful conditions, there was no Cd accumulation (data not shown), whereas after exposure to Cd stress, high




Figure 1. (a) Cadmium measurements in 21 days-old tomato mutants and WT grown with 100 µM-CdCl₂ in the coordinate axis, values above and below the 0 corresponds to the cadmium shoot and root content, respectively; (b) Cd translocation index. Data are means ± SEM (n = 3). The same lowercase letters on the bars of each panel indicate non-significant differences at p ≤ 0.05 by Tukey's test.

accumulation of Cd was observed in both shoots and roots. This accumulation was more pronounced in roots and was not affected by the genotype (Figure 1). In addition, the translocation index was similar for all genotypes (Figure 1b).

The presence of cadmium in the nutrient solution decreased leaf and root area and the length of roots in all genotypes (Figure 2a,b). Although Cd accumulated mainly in roots, we observed an increase in root diameter in *fri* and *phyB2*, whereas WT and *tri* did not show differences between the control and Cd treatments (Figure 2c). In fact, exposure to Cd inhibited root length, area and density in all genotypes (Figure 2).

After treatment with CdCl₂, WT, *fri* and *tri* exhibited lower shoot fresh weight, while no significant change in root biomass was detected (Figure 3a). However, only WT and *tri* suffered reduction of shoot dry weight after Cd exposure (Figure 3b). Plants cultivated in CdCl₂ did not exhibit a reduction in the root dry weight, but *phyB2*, showed an increase in dry weight after exposure to CdCl₂ (Figure 3b). In addition, Cd exposure caused a strong height reduction in




Figure 2. Leaf and root measurements in 21 days-old tomato mutants and WT grown without or with 100 µM-CdCl₂. (a) in the coordinate axis, values above and below the 0 corresponds to the leaf and root area, respectively; (b) Root length; (c) diameter and (d) and density. Control plants of each genotype did not receive Cd. Data are means ± SE (n = 3). Different lowercase letters on the bars of each panel indicate significant differences at p ≤ 0.05 by Tukey's test.

WT, *fri* and *tri*, while plant height was unaltered by high Cd in *phyB2* (Figure 3c). However, in the absence of Cd, *phyB2* and *fri* had similar heights as WT, whereas *tri* was taller.




Figure 3. Leaf and root measurements in 21 days-old tomato mutants and WT grown without or with 100 μM CdCl₂. (a) in the coordinate axis, values above and below the 0 corresponds to the shoot and root fresh weight, respectively; (b) values above and below the 0 corresponds to the shoot and root dry weight, respectively and (c) shoot height. Control plants of each genotype did not receive Cd. Data are means ± SE (n = 3). Different lowercase letters on the bars of each panel indicate significant differences at p ≤ 0.05 by Tukey's test.

The supply of CdCl₂ significantly decreased chlorophyll content in all genotypes, especially in the phytochrome mutants (Figure 4a). Additionally, there was a clear reduction of chlorophyll fluorescence in WT, *fri* and *tri*, while no significant change was observed in Cd-treated *phyB2* plants (Figure 4b).

All genotypes showed leaf blades with uniseriated adaxial and abaxial epidermis, palisade parenchyma presenting one layer of cells, spongy parenchyma thicker than the palisade parenchyma and evident intercellular spaces. The vascular bundles were trapezoidal and surrounded by parenchyma cells in both the adaxial and abaxial sides of the leaf (Figures 5a-f).




Figure 4. Chlorophyll and fluorescence measurements in 21 days-old tomato mutants and WT grown without or with 100 μM CdCl₂. (a) chlorophyll content and (b) fluorescence. Control plants of each genotype did not receive Cd. Data are means ± SE (n = 3). Different lowercase letters on the bars of each panel indicate significant differences at p ≤ 0.05 by Tukey's test.

The *fri* and *phyB2* plants (without the addition of CdCl₂) had a large amount of parenchyma cells above the vascular bundles, which was not observed in these same genotypes when exposed to Cd, indicating a reduction in the number of cells of this tissue in this area. After the other genotypes were analyzed, we observed that there was no difference in the parenchymal cell density both above and below the vascular bundles. The *tri* genotype usually had a prominent apical cell above the parenchyma of the vascular bundles (Figure 5).

When we looked at the amount of parenchyma cells located below the vascular bundles in the midrib of the analyzed leaves, we also noticed a reduction in the number of these cells in the *fri* and *tri* genotypes supplied with Cd. The vascular bundles of the *fri* genotype showed a smaller diameter in plants undergoing Cd stress, while only small differences could be observed in the diameter of vascular bundles for the remaining genotypes (Table 1). There was a decrease in the number of xylem cells in all genotypes except *phyB2* in Cd stress conditions. The significant decrease in the number of xylem cells in *fri* plants, as well as the large




Figure 5. Anatomical analysis in 21 days-old tomato mutants and WT grown without or with 100 μM -CdCl₂. (a) WT without Cd; (b) WT with Cd; (c) *fri* without Cd; (d) *fri* with Cd; (e) *tri* without Cd; (f) *tri* with Cd; (g) *phyB2* without Cd and (h) *phyB2* with Cd. Scales = 30 μm .

number of tiny cells of xylem in *tri* and *fri* before Cd contact, were clearly noted (Table 1).

It is increasingly evident that phytochromes play a key role in abiotic stress responses (Zilli et al. 2008; Carvalho et al. 2011a; Xie et al. 2011). In this study, we used the *phyA* (*fri*), *phyB1* (*tri*), and *phyB2* (*phyB2*) mutants of tomato to evaluate whether some of these phytochromes were involved in plant responses to Cd stress conditions. D'Amico-Damião et al. (2015) reported that phytochrome B

mutants grow better under drought conditions. Thus, better plant development of these genotypes may be expected in Cd conditions.

However, all plants except *phyB2* exhibited reduced shoot fresh and dry weight and reduced height in the presence of Cd (Figures 2a-c). Although this observation could suggest that *phyB2* was tolerant to Cd stress, these values were similar to those of the WT plants in the same conditions, suggesting that *phyB2* plants grow slower in non-stressful conditions. →

Table 1. Anatomical comparison among 21 days-old tomato mutants and WT grown without or with 100 μM CdCl_2 . *Signs ++, +- and -- indicating, respectively, greater, middle and less integrity, diameter or amount of cells from 21 days-old tomato mutants and WT grown without or with 100 μM CdCl_2 .

Characteristics	CdCl_2 (μM)	WT	<i>fri</i>	<i>tri</i>	<i>phyB2</i>
Parenchymal integrity	0	--*	+-	++	--
	100	++	--	--	--
Parenchyma above the vascular bundles	0	+-	++	+-	++
	100	+-	+-	+-	+-
Parenchyma below the vascular bundles	0	++	++	++	+-
	100	++	+-	+-	+-
Diameter of the vascular bundles	0	+-	++	+-	+-
	100	+-	--	+-	+-
Number of xylem cells	0	++	++	++	+-
	100	+-	--	+-	+-

On the other hand, Cd induced a larger root diameter in all photomorphogenic mutants (Figure 1c), especially *fri*, which had a larger root diameter than WT, but there was no increase in root dry weight (Figure 2b). These results can be explained by the impairment of root elongation and phytochrome involvement in root development. In fact, it has been observed that phytochromes triggered the expression of *ELONGATION HYPOCOTYLS 5 (HY5)* and, in turn, the HY5 protein acted as a transcription factor and promoted root growth in response to light (Lee et al. 2016). In addition, it was verified that plants treated with Cd exhibited an enlargement of the roots and a reduction in root elongation (Fusconi et al. 2007; Maksimovic et al. 2007). Changes in root development in response to Cd exposure could be due to negative effects on the root cell microtubule cytoskeleton, which plays a crucial role during root development (Fusconi et al. 2007). However, more studies are needed to clarify these issues.

Additionally, *tri* was taller than WT in both control and stressful conditions, but similar shoot dry weight was verified, indicating no difference in biomass accumulation (Figure 2). Indeed, it was demonstrated that *phyB1* plays a key role in stem elongation and shade avoidance responses (Schrager-Lavelle et al. 2016). Thus, *tri*, a *phyB1*-deficient mutant, exhibited a pronounced elongation of the stem due to impaired light perception (Figure 2c).

The Cd accumulation and translocation index were similar in all genotypes (Figure 1). Despite this similarity, *fri* plants showed a reduced number of xylem cells and a reduced diameter of the vascular bundles compared to WT plants after Cd exposure (Table 1). The same result

was observed by Auge et al. (2012), whose hand-cut stem cross-sections showed a lower xylem vessel number and transversal area in *fri* plants under high evaporative demand. This reduction in xylem caused a reduction in the water supply to the leaves (Auge et al. 2012) and was expected to affect the transport of Cd from the roots to the shoot. In fact, Cd enters into the root system through the symplast or apoplast and eventually reaches the xylem, where it can be translocated to shoots (Lux et al. 2011). Furthermore, Cd translocation in the roots is impaired by the barriers in the exo- and endoderm (Redjala et al. 2011). Therefore, the destabilization of the root tissue facilitates the diffusion of Cd and its loading in the xylem, thus affecting Cd transport from the roots to the shoot (Lux et al. 2011). In addition, Cd accumulation in the shoot causes severe damage to plant physiological processes such as photosynthesis, water relations, mineral metabolism and leaf morphology (Lopez-Chuken and Young 2010; Gill et al. 2012; Asgher et al. 2015). In our study, we observed Cd in the shoots of all genotypes (Figure 1), and there was also a clear disruption in the xylem and parenchymal tissues (Figure 5 and Table 1), as well as a reduction in chlorophyll content and fluorescence, independent of genotype (Figure 4), which were most likely due to instability in the chloroplast membrane due to the Cd accumulation.

In order to assess how Cd exposure affects plant growth and development, the anatomical structure of leaves was investigated. So, Cd accumulation and the loss of phytochrome function in the mutants induced clear cell disorganization caused by stress conditions. In addition, we observed increased cell and vascular bundle disorganization in all genotypes, but mainly in mutants (Figure 5 and Table 1). More recently, it has been demonstrated that phytochromes play a pivotal role in cell division and cell number in the leaf blade (Tsukaya et al. 2002; Nishihama et al. 2015). For example, the *phyB* mutant of *Arabidopsis* exhibited lower leaf size and cell number in the leaf mesophyll compared to the WT (Tsukaya et al. 2002). In addition, several studies have proven that Cd has prejudicial effects on cell ultrastructure and leaf anatomy (Ali et al. 2015; Plusquin et al. 2015; Vaculik et al. 2015). For instance, in Cd stress conditions, ethylene- and auxin-insensitive tomato mutants showed reduced mesophyll cell size, smaller palisade parenchyma cells and disorganization of the chloroplasts (Gratão et al. 2009).

In fact, our results could indicate the control of these responses by phytochromes. Clear anatomical differences were observed between the tested phytochrome mutants. For instance, *fri* and *phyB2* exhibited more parenchyma cells above the vascular bundles, while *tri* had a prominent apical cell in the parenchyma above the vascular bundles. However, under Cd stress conditions, both parenchymal integrity and parenchyma cells above or below the vascular bundles were severely reduced in *fri* compared to WT plants (Figure 5 and Table 1).

CONCLUSION

Despite mounting evidence that phytochromes are involved in plant responses to several abiotic stresses (Carvalho et al. 2011b; Auge et al. 2012), none of the three phytochrome mutants assessed in the present study exhibited higher tolerance to Cd. Thus, taken together, these results indicate that phytochromes do not integrate the pathway of Cd stress response in tomato plants.

REFERENCES

- Ali, B., Deng, X., Hu, X., Gill, R. A., Ali, S., Wang, S. and Zhou, W. (2015). Deteriorative effects of cadmium stress on antioxidant system and cellular structure in germinating seeds of *Brassica napus* L. *Journal of Agricultural Science and Technology*, 17, 63-74.
- Asgher, M., Khan, M. I., Anjum, N. A. and Khan, N. A. (2015). Minimising toxicity of cadmium in plants — role of plant growth regulators. *Protoplasma*, 252, 399-413. <https://doi.org/10.1007/s00709-014-0710-4>.
- Auge, G. A., Rugnone, M. L., Cortés, L. E., González, C. V., Zarlavsky, G., Bocalandro, H. E. and Sánchez, R. A. (2012). Phytochrome A increase tolerance to high evaporative demand. *Physiologia Plantarum*, 151, 228-235. <https://doi.org/10.1111/j.1399-3054.2012.01625.x>.
- Bae, G. and Choi, G. (2008). Decoding of light signals by plant phytochromes and their interacting proteins. *Annual Review of Plant Biology*, 59, 281-311. <https://doi.org/10.1146/annurev.arplant.59.032607.092859>.
- Balestrasse, K. B., Yannarelli, G. G., Noriega, G. O., Batlle, A. and Tomaro, M. L. (2008b). Hemeoxygenase and catalase gene expression in nodules and roots of soybean plants subjected to cadmium stress. *Biometals*, 21, 433-441. <https://doi.org/10.1007/s10534-008-9132-0>.
- Balestrasse, K. B., Zilli, C. G. and Tomaro, M. L. (2008a). Signal transduction pathways and haem oxygenase induction in soybean leaves subjected to salt stress. *Redox Report*, 13, 255-262. <https://doi.org/10.1179/135100008x308966>.
- Bocalandro, H. E., Mazza, C. A., Mazzella, M. A., Casal, J. J. and Ballaré, C. L. (2001). Ultraviolet B radiation enhances a phytochrome-B mediated photomorphogenic response in *Arabidopsis*. *Plant Physiology*, 126, 780-788. <https://doi.org/10.1104/pp.126.2.780>.
- Carvalho, R. F., Aidar, S. T., Azevedo, R. A., Dodd, I. C. and Peres, L. E. P. (2011b). Enhanced transpiration rate in the *high pigment 1* tomato mutant and its physiological significance. *Plant Biology*, 13, 546-550. <https://doi.org/10.1111/j.1438-8677.2010.00438.x>.
- Carvalho, R. F., Campos, M. L. and Azevedo, R. A. (2011a). The role of phytochrome in stress tolerance. *Journal of Integrative Plant Biology*, 53, 920-929. <https://doi.org/10.1111/j.1744-7909.2011.01081.x>.
- Cui, W. T., Fu, G. Q., Wu, H. H. and Shen, W. B. (2011). Cadmium-induced *heme oxygenase-1* gene expression is associated with the depletion of glutathione in the roots of *Medicago sativa*. *Biometals*, 24, 93-103. <https://doi.org/10.1007/s10534-010-9377-2>.
- D'Amico-Damião, V., Cruz, F. J. R., Gavassi, M. A., Santos, D. M. M., Melo, H. C. and Carvalho, R. F. (2015). Photomorphogenic modulation of water stress in tomato (*Solanum lycopersicum* L.): the role of phytochromes A, B1, and B2. *Journal of Horticultural Science and Biotechnology*, 90, 25-30. <https://doi.org/10.1080/14620316.2015.11513149>.
- Essen, L. O., Mailliet, J. and Hughes, J. (2008). The structure of a complete phytochrome sensory module in the Pr ground state. *Proceedings of the National Academy of Science of the U.S.A.*, 105, 14709-14714. <https://doi.org/10.1073/pnas.0806477105>.
- Franklin, K. A. (2016). Photomorphogenesis: plants feel blue in the shade. *Current Biology*, 26, 1275-1276. <https://doi.org/10.1016/j.cub.2016.10.039>.
- Franklin, K. A. and Quail, P. H. (2010). Phytochrome functions in *Arabidopsis* development. *Journal of Experimental Botany*, 61, 11-24. <https://doi.org/10.1093/jxb/erp304>.
- Fusconi, A., Gallo, C. and Camusso, W. (2007). Effects of cadmium on root apical meristems of *Pisum sativum* L.: Cell viability, cell

- proliferation and microtubule pattern as suitable markers for assessment of stress pollution. *Mutation Research*, 632, 9-19. <https://doi.org/10.1016/j.mrgentox.2007.03.012>.
- Gill, S. S., Khan, N. A. and Tuteja, N. (2012). Cadmium at high dose perturbs growth, photosynthesis and nitrogen metabolism while at low dose it up regulates sulfur assimilation and antioxidant machinery in garden cress (*Lepidium sativum* L.). *Plant Science*, 182, 112-120. <https://doi.org/10.1016/j.plantsci.2011.04.018>.
- Gratão, P. L., Monteiro, C. C., Carvalho, R. F., Tezotto, T., Piotto, F. A., Peres L. E. P. and Azevedo, R. A. (2012). Biochemical dissection of *diageotropica* and *Never ripe* tomato mutants to Cd-stressful conditions. *Plant Physiology and Biochemistry*, 56, 79-96. <https://doi.org/10.1016/j.plaphy.2012.04.009>.
- Gratão, P. L., Monteiro, C. C., Rossi, M. L., Martinelli, A. P., Peres, L. E. P., Medici, L. O., Lea, P. J. and Azevedo, R. A. (2009). Differential ultrastructural changes in tomato hormonal mutants exposed to cadmium. *Environmental and Experimental Botany*, 67, 387-394. <https://doi.org/10.1016/j.envexpbot.2009.06.017>.
- Gratão, P. L., Monteiro, C. C., Tezotto, T., Carvalho, R. F., Alves, L. R., Peters, L. P. and Azevedo, R. A. (2015). Cadmium stress antioxidant responses and root-to-shoot communication in grafted tomato plants. *Biometals*, 28, 803-816. <https://doi.org/10.1007/s10534-015-9867-3>.
- Hiltbrunner, A., Viczian, A., Bury, E., Tscheuschler A., Kircher, S., Toth, R., Honsberger, A., Nagy, F., Fankhauser, C. and Schafer, E. (2005). Nuclear accumulation of the phytochrome A photoreceptor requires FHY1. *Current Biology*, 15, 2125-2130. <https://doi.org/10.1016/j.cub.2005.10.042>.
- Jin, Q. J., Zhu, K. K., Xie, Y. J. and Shen, W. B. (2013). *Heme oxygenase-1* is involved in ascorbic acid-induced alleviation of cadmium toxicity in root tissues of *Medicago sativa*. *Plant and Soil*, 366, 605-616. <https://doi.org/10.1007/s11104-012-1451-9>.
- Johansen, D. A. (1940). *Plant microtechnique*. New York: McGraw-Hill Book Co. Inc.
- Kim, H., Kim, Y., Park, J. and Kim, J. (2002). Light signalling mediated by phytochrome plays an important role in cold-induced gene expression through the C-repeat/dehydration responsive element (C/DRE) in *Arabidopsis thaliana*. *The Plant Journal*, 29, 693-704. <https://doi.org/10.1046/j.1365-313x.2002.01249.x>.
- Lee, H-J., Ha, J-H., Kim, S-G., Choi, H-K., Kim, Z. H., Han, Y-J., Kim, J-I., Oh, Y., Fragoso, V., Shin, K., Hyeon, T., Choi, H-G., Oh, K-H., Baldwin, I. T. and Park, C-M. (2016). Stem-piped light activates phytochrome B to trigger light responses in *Arabidopsis thaliana* roots. *Science Signaling*, 9, 1-8. <https://doi.org/10.1126/scisignal.aaf6530>.
- Liu, J., Zhang, F., Zhou, J., Chen, F., Wang, B. and Xie X. (2012). Phytochrome B control of total leaf area and stomatal density affects drought tolerance in rice. *Plant Molecular Biology*, 78, 289-300. <https://doi.org/10.1007/s11103-011-9860-3>.
- Lopez-Chuken, U. J. and Young, S. D. (2010). Modelling sulphate-enhanced cadmium uptake by *Zea mays* from nutrient solution under conditions of constant free Cd²⁺ ion activity. *Journal of Environmental Sciences*, 22, 1080-1085. [https://doi.org/10.1016/s1001-0742\(09\)60220-5](https://doi.org/10.1016/s1001-0742(09)60220-5).
- Lux, A., Martinka, M., Vaculik, M. and White, P. J. (2011). Root responses to cadmium in the rhizosphere: a review. *Journal of Experimental Botany*, 62, 21-37. <https://doi.org/10.1093/jxb/erq281>.
- Maksimovic, I., Kastori, R., Krstic, L. and Lukovic, J. (2007). Steady presence of cadmium and nickel affects root anatomy, accumulation and distribution of essential ions in maize seedlings. *Biologia Plantarum*, 51, 589-592. <https://doi.org/10.1007/s10535-007-0129-2>.
- Malavolta, E., Vitti, G. C. and Oliveira, S. A. (2011). *Avaliação do estado nutricional de plantas: princípios e aplicações*, 2 ed. Piracicaba: Associação Brasileira para Pesquisa de Potassa e do Fosfato.
- Nagajyoti, P. C., Lee, K. D. and Sreekanth, T. V. M. (2010). Heavy metals, occurrence and toxicity for plants: a review. *Environmental Chemistry Letters*, 8, 199-216. <https://doi.org/10.1007/s10311-010-0297-8>.
- Neff, M. M., Fankhauser, C. and Chory, J. (2000). Light: an indicator of time and place. *Genes and Development*, 14, 257-271.
- Nishihama, R., Ishizaki, K., Hosaka, M., Matsuda, Y., Kubota, A. and Kohchi, T. (2015). Phytochrome-mediated regulation of cell division and growth during regeneration and sporeling development in the liverwort *Marchantia polymorpha*. *Journal of Plant Research*. <https://doi.org/10.1007/s10265-015-0724-9>.
- O'Brien, T. P., Feder, N. and Mccully, M. M. E. (1964). Polychromatic staining of plant cell wall by toluidine blue. *Protoplasma*, 59, 367-373. <https://doi.org/10.1007/bf01248568>.
- Plusquin, M., Mulder, K., Belleghem, F. V., DeGheselle, O., Pirotte, N., Willems, M., Cuypers, A., Salvenmoser, W., Laudurner, P., Artois, T. and Smeets, K. (2015). Toxic effects of cadmium on flatworm stem cell dynamics: a transcriptomic and ultrastructural elucidation of underlying mechanisms. *Environmental Toxicology*, 31, 1217-1228. <https://doi.org/10.1002/tox.22129>.

- Quail, P. H. (2002a). Photosensory perception and signalling in plant cells: new paradigms? *Current Opinion in Cell Biology*, 14, 180-188. [https://doi.org/10.1016/s0955-0674\(02\)00309-5](https://doi.org/10.1016/s0955-0674(02)00309-5).
- Quail, P. H. (2002b). Phytochrome photosensory signalling networks. *Nature Reviews Molecular Cell Biology*, 3, 85-93. <https://doi.org/10.1038/nrm728>.
- Redjala, T., Zelko, I., Sterckeman, T., Legué, V. and Lux, A. (2011). Relationship between root structure and root cadmium uptake in maize. *Environmental and Experimental Botany*, 71, 241-248. <https://doi.org/10.1016/j.envexpbot.2010.12.010>.
- Rossato, L. V., Nicoloso, F. T., Farias, J. G., Cargnelluti, D, Tabaldi, L. A., Antes, F. G., Dressler, V. L., Morsch, V. M. and Schetinger, M. R. C. (2011). Effects of lead on the growth, lead accumulation and physiological responses of *Pluchea agittalis*. *Ecotoxicology*, 21, 111-123. <https://doi.org/10.1007/s10646-011-0771-5>.
- Saijo, Y., Hata, S., Kyojuka, J., Shimamoto, K. and Izui, K. (2000). Over-expression of a single Ca²⁺-dependent protein kinase confers both cold and salt/drought tolerance on rice plants. *The Plant Journal*, 23, 319-327. <https://doi.org/10.1046/j.1365-313x.2000.00787.x>.
- Schrager-Lavelle, A., Herrera, L. A., and Maloof, J. N. (2016). Tomato phyE is required for shade avoidance in the absence of phyB1 and phyB2. *Frontiers in Plant Science*, 7, 1275. <https://doi.org/10.3389/fpls.2016.01275>.
- Shen, Q., Jiang, M., Li, H., Che, L. L. and Yang, Z. M. (2011). Expression of a *Brassica napus* heme oxygenase confers plant tolerance to mercury toxicity. *Plant Cell and Environment*, 34, 752-763. <https://doi.org/10.1111/j.1365-3040.2011.02279.x>.
- Tsukaya, H., Kozuka, T. and Kim, G. T. (2002). Genetic control of petiole length in *Arabidopsis thaliana*. *Plant Cell Physiology*, 43, 1221-1228. <https://doi.org/10.1093/pcp/pcf147>.
- Usman, A. R. A., Lee, S. S., Awad, Y. M., Lim, K. J., Yang, J. E. and Ok, Y. S. (2012). Soil pollution assessment and identification of hyperaccumulating plants in chromated copper arsenate (CCA) contaminated sites. Korea. *Chemosphere*, 87, 872-878. <https://doi.org/10.1016/j.chemosphere.2012.01.028>.
- Vaculik, M., Pavlovic, A. and Lux, A. (2015). Silicon alleviates cadmium toxicity by enhanced photosynthetic rate and modified bundles sheath's cell chloroplast ultrastructure in maize. *Ecotoxicology and Environmental Safety*, 120, 66-73. <https://doi.org/10.1016/j.ecoenv.2015.05.026>.
- Verkleij, J. A. C., Golan-Goldhirsh, A., Antosiewicz, D. M., Schwitzguebel, J. P. and Schroder, P. (2009). Dualities in plant tolerance to pollutants and their uptake and translocation to the upper plant parts. *Environmental and Experimental Botany*, 67, 10-22. <https://doi.org/10.1016/j.envexpbot.2009.05.009>.
- Wit, M., Galvão, V. C. and Fankhauser, C. (2016). Light-mediated hormonal regulation of plant growth and development. *Annual Review of Plant Biology*, 67, 513-537. <https://doi.org/10.1146/annurev-arplant-043015-112252>.
- Xie, Y. J., Cui, W. T., Yuan, X. X., Shen, W. B. and Yang, Q. (2011). *Heme oxygenase-1* is associated with wheat salinity acclimation by modulating reactive oxygen species homeostasis. *Journal of Integrative Plant Biology*, 53, 653-670. <https://doi.org/10.1111/j.1744-7909.2011.01052.x>.
- Yang, X., Kuk, J. and Moffat, K. (2008). Crystal structure of *Pseudomonas aeruginosa* bacteriophytochrome: photoconversion and signal transduction. *Proceedings of the National Academy of Sciences of the U.S.A.*, 105, 14715-14720. <https://doi.org/10.1073/pnas.0806718105>.
- Zao, Y., Zhou, J. and Ding, D. (2014). Phytochrome B-mediated activation of lipoxygenase modulates an excess red light-induced defence response in *Arabidopsis*. *Journal of Experimental Botany*, 65, 4907-4918. <https://doi.org/10.1093/jxb/eru247>.
- Zilli, C. G., Balestrasse, K. B., Yannarelli, G. G., Polizio, A. H., Santa-Cruz, D. M. and Tomaro, M. L. (2008). Hemeoxygenase up-regulation under salt stress protects nitrogen metabolism in nodules of soybean plants. *Environmental and Experimental Botany*, 64, 83-89. <https://doi.org/10.1016/j.envexpbot.2008.03.005>.