

What do we know about Neotropical trap-nesting bees?

Synopsis about their nest biology and taxonomy

Camila Cristina Ferreira da Costa^{1,2} & Rodrigo Barbosa Gonçalves^{1,3}

¹ Universidade Federal do Paraná (UFPR), Departamento de Zoologia (DZOO). Curitiba, PR, Brasil.

² ORCID: <http://orcid.org/0000-0003-1950-7525>. E-mail: ccfdacosta@gmail.com (corresponding author)

³ ORCID: <http://orcid.org/0000-0001-5623-0938>. E-mail: goncalvesrb@gmail.com

Abstract. Cavity-nesting bees are enigmatic because they are difficult to observe in the wild, hence trap-nests (man-made cavities) provide the means by which these bees may be studied. Trap-nests is an efficient methodology to study these bees and are common worldwide. These traps have been used for a variety of reasons, including inventories, to examine pollen load, to study habitat disturbance, and bee conservation. However Neotropical trap-nesting bees' taxonomy and biology are still poorly known and here we provide a review about these subjects. We searched for trap-nest bee studies in the Neotropical Region using Google Scholar and ISI Web of Science at any time in the past to December 2017. We found 109 independent studies, most of which were from Brazil (87 studies), followed by Argentina (10 studies), and other countries had fewer than five studies each. A total of 140 species, 24 genera, 10 tribes and three subfamilies were reported in trap-nests. Nest architecture was described for only 49 species. Taxonomy is only well-known for 14 genera, somewhat known for seven and is essentially unavailable for three genera. Construction material, closing plug and cell shape are similar among species in the same tribes and genera. Vestibular and intercalary cells, and the preliminary plug are variable, even at the specific level. Apinae is the most studied group with available data for all genera recorded in trap-nests. Colletinae is the least-studied group and nothing is known for their nesting biology. Megachilinae is intermediate, with some studies of taxonomy and nesting. We suggest that further trap-nest studies should provide more detailed information on nest architecture and construction materials, including explicit mention of structures that are absent. All Neotropical bees need more taxonomic studies, but some, such as *Hylaeus* and *Megachile*, require more attention since *Hylaeus* is essentially unknown and *Megachile* is very common on trap-nests.

Key-Words. Bee hotels; Diversity; Methodology; Nesting behavior; Systematics.

INTRODUCTION

Bee nests ordinarily comprises brood cells and associated structures and are often in burrows in the soil, aboveground cavities or free-standing (Michener, 2007). Most bees and apoid wasps excavate underground nests and this form of nesting is primitive in the superfamily (Melo, 1999; Hedtke *et al.*, 2013; Branstetter *et al.*, 2017). While we do not yet have a phylogenetic reconstruction of substrate preference for all bee species, apparently aboveground nesting arose independently several times. Four of the seven bee main lineages have species that nest in cavities and there are some reversals to soil nesting (Almeida, 2008).

Aboveground substrates are variable and cavity-nesting bees are likely to be an artificial ecological grouping. The use of existing tunnels in deadwood is common and bees often excavate decomposing wood and soft pith in stems and galls for nests (Sheffield *et al.*, 2011). Other exam-

ples of natural substrates include snail shells (Gess & Gess, 2008), rock surfaces (Eickwort, 1975) and man-made cavities, such as in brick walls (Santos *et al.*, 2016), metal frames (Sheffield *et al.*, 2011), farm tractor radiators (Sheffield, 2017) and door locks (RBG *pers. obs.*).

As a consequence of the wide variety of nesting substrates, artificial nesting substrates (trap-nests, nest-boxes, bee hotels) can be used to trap these cavity nesting bees and wasps (Krombein, 1967; Maclvor & Packer, 2015). These traps are often made of bundles of hollow stems, paper or cardboard tubes (Camillo *et al.*, 1995; Araújo *et al.*, 2016) and holes drilled in wood (Krombein, 1967; Buschini, 2006; see Maclvor, 2017 for a review). Characteristics of the entrance diameter, nest length, color and also placement of nests all influence bee selection and use of traps-nest (Krombein, 1967; Maclvor & Packer, 2015). Also, traps can be placed in aggregates of greater density to improve the likelihood of use. Studies tend

Pap. Avulsos Zool., 2019; v.59: e20195926

<http://doi.org/10.11606/1807-0205/2019.59.26>

<http://www.revistas.usp.br/paz>

<http://www.scielo.br/paz>

Edited by: Kelli dos Santos Ramos

Received: 29/08/2018

Accepted: 15/04/2019

Published: 15/05/2019

ISSN On-Line: 1807-0205

ISSN Printed: 0031-1049

ISNI: 0000-0004-0384-1825



to develop their own type of trap and so a wide variety of traps and their dispositions have been used, making comparisons of these studies very difficult.

As a consequence of the successful use of traps, studies are common worldwide, with over 1,300 results in Google (January 2018). Traps are used for many reasons, including to sample and monitor cavity nesting species and their predators (Araujo *et al.*, 2017; Oliveira & Gonçalves, 2017), to compare habitats among different regions (Araújo *et al.*, 2016), to examine altitudinal gradients (Perillo *et al.*, 2017) and vertical stratification (Morato, 2001b; Stangler *et al.*, 2015, 2016), to detect responses to fragmentation (Stangler *et al.*, 2015, 2016; Rocha-Filho *et al.*, 2017) and urbanization (Pereira-Peixoto *et al.*, 2014), to promote pollination and pollinator conservation (MacIvor & Packer, 2015) and to study the nest biology and behavior of particular groups (Rocha-Filho & Garófalo, 2016a,b; Moure-Oliveira *et al.*, 2017).

Despite of the large number of trap-nest studies, the Neotropical bee fauna taxonomy and diversity is still poorly known (Silveira *et al.*, 2002). Here we summarize the available information on biology and taxonomy of trap-nesting bees in this region. A synopsis is important to provide direction for future studies because further coordinated efforts will be important to produce comparable data and robust advances in this research field.

MATERIAL AND METHODS

To summarize the trap-nesting studies, we searched using Google Scholar and ISI Web of Science through the end of 2017. We used the following search terms: (Ninhos armadilha OR Nidos trampa OR Trap-nest OR Trap-nest bees) AND (Neotropical OR *countries names*). We included all countries from Chile to Mexico as search terms. The literature cited along any retrieved study was also used to find additional references. The following criteria was used to select the studies for this work: (1) used trap-nesting methods (understood here as any artificial cavity that was built by the researcher in which bees nested); (2) a primary reference, revision studies were not included; and (3) published in a peer reviewed journal or as an academic thesis or dissertation (other gray literature such as abstracts and conference reports were not included).

Data for nesting behavior, other biological details at higher taxonomic levels and the number of species in the world follow Michener (2007). The number of Neotropical species follows the online version of Moure's Bee Catalogue (Moure *et al.*, 2013). Nesting behaviors of species and genera were gathered from the original studies. Terminology for nest architecture follows Krombein (1967) as illustrated in Fig. 1. Trap-nest biology knowledge is considered "available" when there is at least one published description with details on the architecture using the trap-nest methodology, and "unavailable" in the absence of this information. Species identification was taken from the original studies and, if necessary, revised following Moure's Bee Catalogue (Moure *et al.*, 2013). We adopt the single-family classification for bees following

Table 1. A summary of trap-nesting bee genera from the Neotropical region. Notes: ¹morphospecies are excluded for most genera; ²only recorded as morphospecies; ³available for nest description of at least one species; ⁴sufficient for taxa with published taxonomical revision, moderate for taxa only identified by taxonomists, insufficient when no revision is available.

Taxon	Number of recorded species ¹	Trap-nest description ³	Taxonomic knowledge ⁴
Apinae			
Centridini			
<i>Centris</i>	14	Available	Moderate
Euglossini			
<i>Eufriesea</i>	8	Available	Sufficient
<i>Euglossa</i>	14	Available	Moderate
Tetrapediini			
<i>Tetrapedia</i>	7	Available	Insufficient
Xylocopini			
<i>Xylocopa</i>	6	Available	Moderate
Colletinae			
Colletini			
<i>Colletes</i>	1	Unavailable	Insufficient
<i>Rhynchocolletes</i>	1 ²	Available	Sufficient
Hylaeini			
<i>Hylaeus</i>	1	Unavailable	Insufficient
Megachilinae			
Anthidiini			
<i>Anthidium</i>	4	Available	Sufficient
<i>Anthidulum</i>	1 ²	Unavailable	Sufficient
<i>Anthodioctes</i>	5	Available	Sufficient
<i>Carlotala</i>	1	Available	Sufficient
<i>Ctenanthidium</i>	1	Available	Sufficient
<i>Dicranthidium</i>	3	Unavailable	Sufficient
<i>Duckeanthidium</i>	1	Available	Moderate
<i>Epanthidium</i>	5	Available	Sufficient
<i>Hypanthidium</i>	1	Unavailable	Moderate
<i>Loyolanthidium</i>	1 ²	Unavailable	Moderate
<i>Nananthidium</i>	1	Unavailable	Sufficient
<i>Saranthidium</i>	2	Unavailable	Sufficient
Lithurgini			
<i>Microthurga</i>	1	Unavailable	Sufficient
<i>Trichothurgus</i>	1	Available	Sufficient
Megachilini			
<i>Megachile</i>	33	Available	Insufficient
Osmiini			
<i>Heriades</i>	1 ²	Unavailable	Moderate

Melo & Gonçalves (2005). Identification at morphospecies ("spp.") was also included in Table S1 but not counted in Table 1 except for the genera only recorded for undetermined species. We opted to include all records of primary references except for a single morphospecies of *Neofidelia* (Veddeler *et al.*, 2010). This refers to soil nesting species that are probably incorrectly identified. Taxonomy for each genus was evaluated at the species level and was considered "sufficient" when published taxonomical revision with identification keys is available, "moderate" when species are relatively well known and partial (regional or subgeneric) revisions are available, but much of the identification relies on taxonomists, and "insufficient" otherwise. Identification resources were gathered from Michener (2007), Moure *et al.* (2013) and




Figure 1. A generalized bee trap-nest architecture. At the left is the first cell made. The dark grey indicates food objects for the larvae (white). On the right is the last cell built with presence of a vestibular cell being variable.

published studies as described above. For the distributions of genera we followed Moure *et al.* (2013) and for species we used information from the original trap-nest studies (Table S1). States or provinces are informed of Argentina, Brazil and Mexico.

RESULTS

Our literature search resulted in a total of 109 independent studies, 87 from Brazil followed by Argentina (10), Costa Rica (5), Ecuador (2), Colombia (2), Mexico (1), Jamaica (1) and Trinidad and Tobago (1) (Table S1 summarizes trap-nesting bee literature). These studies comprised 140 species, 24 genera, 10 tribes and three subfamilies of trap-nesting bees in the Neotropical region. Nest architecture was described for 49 species and another 65 species were reported without descriptions (Tables 1 and 2). Fourteen genera had sufficient taxonomic descriptions, seven moderate and three insufficient.

APINAE. Although this group lacks a phylogenetic consensus (Cardinal *et al.*, 2010; Hedtke *et al.*, 2013; Martins *et al.*, 2014; Bossert *et al.*, 2019), wood nesting arose from soil nesting at least seven times following those topol-

ogies. All tribes of wood cavity nesting bees except the Tapinotaspidini were sampled using trap-nests, and Centridini, Euglossini and Tetrapediini were often found in these Neotropical studies.

Centridini. Traditionally the tribe comprised *Centris* and *Epicharis* (Bossert *et al.*, 2019), but it can be paraphyletic (*e.g.*, Martins & Melo, 2016). *Centris* included 251 species (Michener, 2007), mostly Neotropical (224 species) (Moure *et al.*, 2013), use floral oils mixed with other material for nest construction and protection (Vinson *et al.*, 1996). Nesting in existing cavities arose two times in this tribe in *C. (Xanthemisia)* and the clade *C. (Hemisiella) + C. (Heterocentris)* (Martins & Melo, 2016) and according with Moure *et al.* (2013) they comprise 35 described species. Only *C. (Hemisiella)* and *C. (Heterocentris)* were reported using trap nests (Vinson *et al.*, 2010; Vélez *et al.*, 2017). These subgenera use different material for nest construction; for example, *Heterocentris* use wood chips and *Hemisiella* use sand (Vinson *et al.*, 2010). Otherwise, nest characteristics are similar among these subgenera (Table 2).

Centris nests entrance diameter varies from 4.8-14 mm (Drummond *et al.*, 2008; Vinson *et al.*, 2010; Carvalho *et al.*,

Table 2. A summary on nest architecture of trap-nesting bee genera from Neotropical region. Data was summarized through the end of 2017. (?) There is not information or is dubious; (P) present, (O) occasional, (A) absent. The genera without published information about nest architecture we did not list in table.

Taxon	Material	Entrance diameter (mm)	Cell arrangement	Number of brood cells	Brood cells shape	Vestibular cells	Closing plug	Empty space	Preliminary plug	Intercalary cells
<i>Anthidium</i>	trichomes, plants parts, detritus	5-11	linear	1-19	?	?	P	?	?	?
<i>Anthodiocetes</i>	resin, mud, wood chips	5-11	linear	2-11	?	O	P	O	O	O
<i>Carlotala</i>	resin, clay, sand	6-10	?	3-6	?	P	?	?	P	?
<i>Centris</i>	oil, wood chips or sand	4.8-14	linear or irregular	1-16	cylindrical or oval	O	P	P	?	O
<i>Ctenanthidium</i>	resin	4	linear	?	?	?	P	P	?	?
<i>Duckeanthidium</i>	resin, saliva	11-13	?	1-3	?	?	?	?	?	?
<i>Epanthidium</i>	resin, mud, sand	?	linear	2-9	?	O	?	?	?	?
<i>Eufriesea</i>	resin, wood chips	15-25	linear or irregular	2-4	oval	O	?	A	A	A
<i>Euglossa</i>	resin	11-22	linear or cluster	2-14	oval	?	P	P	A	A
<i>Megachile</i>	leaves, petals, mud rocks	6-27	linear	1-16	cylindrical	O	?	?	O	?
<i>Tetrapedia</i>	oil, soil, sand	3-12	linear	1-9	?	P	P	P	P	?
<i>Trichothurgus</i>	wood chips, pollen	8-11	?	no partition	?	?	A	?	?	?
<i>Xylocopa</i>	wood particles	12-23	linear	1-6	barrel	?	A	?	?	?

2016; Vélez *et al.*, 2017). Nests usually comprise linear brood cells (Drummond *et al.*, 2008; Vinson *et al.*, 2010; Carvalho *et al.*, 2016; Vélez *et al.*, 2017), with two rows of cells in *Centris tarsata* Smith, 1874 (Aguiar & Garófalo, 2004). Nests have from one to 16 circular to oval brood cells (Aguiar & Garófalo, 2004; Aguiar *et al.*, 2006; Buschini & Wolff, 2006; Drummond *et al.*, 2008; Vinson *et al.*, 2010; Carvalho *et al.*, 2016; Moure-Oliveira *et al.*, 2017; Vélez *et al.*, 2017). Vestibular and intercalary cells are occasionally found (Buschini & Wolff, 2006; Vinson *et al.*, 2010; Vélez *et al.*, 2017). A closing plug and an empty space in the distal end of the cell rows are common (Aguiar & Garófalo, 2004; Aguiar *et al.*, 2006). The genus is mostly Neotropical with few species reaching North America (Moure *et al.*, 2013). Subgenera of *Centris* may be identified by Silveira *et al.* (2002) and Michener (2007). Some *C. (Hemisiella)* and *C. (Heterocentris)* species may be identified by Thiele (2003) and Vivallo & Vélez (2016). Keys to *Centris* species from Central and North America and for species from Argentina are available Snelling (1984) and Roig-Alsina (2000), respectively.

Euglossini. Orchid bees are mostly Neotropical, comprising five genera: *Aglae* (1 Neotropical species), *Eufriesea* (67 Neotropical species), *Euglossa* (128 Neotropical species and 6 subgenera), *Eulaema* (33 Neotropical species and 2 subgenera), and *Exaerete* (8 Neotropical species). This tribe is unique in its elongate tongue and males collect orchid fragrances. *Aglae* and *Exaerete* are cleptoparasites of *Eufriesea* and *Eulaema*. Most species nest in existing cavities in which they do not build storage cells, unlike the other corbiculate bees (Michener, 2007). Taxonomy of the group is mostly based on males while females are identified by experts.

Eufriesea nests are built with wood chips and plant resins (Viana *et al.*, 2001; Kamke *et al.*, 2008). Entrance diameter varies from 15-25 mm (Viana *et al.*, 2001; Kamke *et al.*, 2008). Cell orientation may be horizontal as in *Eufriesea mussitans*, (Fabricius, 1787) (Viana *et al.*, 2001) to irregular as in *Eufriesea smaragdina* (Perty, 1833) (Kamke *et al.*, 2008). In the latter, cells are still built sequentially (Kamke *et al.*, 2008). Usually 2-4 oval and smooth brood cells are built having internal divisions of resin, with occasional vestibular cells (Viana *et al.*, 2001; Kamke *et al.*, 2008). *Eufriesea* is distributed from Argentina to Mexico (Moure *et al.*, 2013). Kimsey (1982) provided a key to the males.

Euglossa have nest architecture in the subgenera *Euglossa* s.s. and *Glossura* similar to *Eufriesea*, with brood cell divisions and closing plugs of resins only (Garófalo *et al.*, 1998; Peruquetti, 1998; Augusto & Garófalo, 2004; Parra-H & Nates-Parra, 2009). Entrance diameters varied from 11-22 mm. Brood cells may be linear (vertical or horizontal) or clustered (Garófalo *et al.*, 1998; Peruquetti, 1998; Augusto & Garófalo, 2004, 2009), and the usually oval cells vary from 4-14 per nest (Garófalo *et al.*, 1998; Peruquetti, 1998; Augusto & Garófalo, 2004). Preliminary plugs were not mentioned, while a distal empty space is typical (Parra-H & Nates-Parra, 2009). Females may nest

alone or with other, usually sister, females, and may reuse empty cells (Silva *et al.*, 2016). If with others, a dominant female remains at the nest while others forage (Augusto & Garófalo, 2004, 2009, 2011; Freiria *et al.*, 2017). *Euglossa* is found from Mexico to Argentina (Moure *et al.*, 2013). Most subgenera can be identified using Silveira *et al.* (2002), while no complete revision of the genus is available. Males from São Paulo (Brazil) can be identified using keys provided by Rebêlo & Moure (1995), males and females of *E. (Glossura)* from Atlantic forest with Faria-Jr. & Melo (2007) and *E. (Euglossa)* in the *E. analis* group using Faria & Melo (2012).

Tetrapediini. Traditionally including two genera, the cleptoparasite *Coelioxoides* Cresson and *Tetrapedia* (Michener, 2007), recent phylogenetic hypotheses place *Coelioxoides* within the cleptoparasitic clade of Apinae (Cardinal *et al.*, 2010; Hedtke *et al.*, 2013). The *Tetrapedia* position in Apinae is uncertain, Martins *et al.* (2014) suggested that the genus is related to Ctenoplectrini, and Bossert *et al.* (2019) consider both as related with Xylocopini. In both genera include species that nest aboveground. *Tetrapedia*, comprising 28 species, is found from Mexico to Argentina (Moure *et al.*, 2013). There is no available key for species identification. They collect floral oils and nest in existing cavities (Alves-dos-Santos *et al.*, 2002). Cell partitions and closing plug are of soil or sand mixed with floral oils (Alves-dos-Santos *et al.*, 2002; Camillo, 2005; Menezes *et al.*, 2012; Rocha-Filho & Garófalo, 2016a). Entrance diameter varies between 3-12 mm (Alves-dos-Santos *et al.*, 2002; Camillo, 2005; Menezes *et al.*, 2012; Rocha-Filho & Garófalo, 2016a). Cells are linear and horizontal (Alves-dos-Santos *et al.*, 2002; Camillo, 2005; Menezes *et al.*, 2012; Rocha-Filho & Garófalo, 2016a), or linear and vertical (Camillo, 2005). Brood cells vary from 1-9 (Alves-dos-Santos *et al.*, 2002; Camillo, 2005; Menezes *et al.*, 2012; Rocha-Filho & Garófalo, 2016a). Vestibular cells, preliminary plugs and distal empty spaces are present (Rocha-Filho & Garófalo, 2016a).

Xylocopini. Among the most speciose lineages of aboveground nesting bees, most nest in cavities, with one reversal to ground-nesting in *X. (Proxylocopa)* (Leys *et al.*, 2002). The tribe has four lineages, of which only Allodapina is not Neotropical (Michener, 2007). *Ceratina*, with 199 species, has never been reported in trap-nests. Also, *Manuelia* (3 species) in Argentina and Chile has not been reported in trap-nests (see Daly *et al.*, 1987; Flores-Prado *et al.*, 2008 for detailed biology). *Xylocopa* (111 species) in the Neotropical region (Moure *et al.*, 2013) includes trap-nesting only in the subgenus *Neoxylocopa* (Table 1, Table S1). Phylogenetic relationships among these four lineages vary by study, yet Allodapina and Ceratinina are consistently considered to be sister groups (Flores-Prado *et al.*, 2008; Cardinal *et al.*, 2010; Martins *et al.*, 2014).

Xylocopa nests comprise fine wood particles that the bees excavate (Marchi & Melo, 2010; Pereira & Garófalo, 2010; Lucia *et al.*, 2017) and cell partitions are of wood

particles mixed with saliva (Pereira & Garófalo, 2010; Lucia *et al.*, 2017). Females of *Xylocopa* and *Ceratina* remove nest partitions and the allodapines lack cell partitions (Michener, 2007). Entrance diameter varies from 12-23 mm (Marchi & Melo, 2010) with 1-6 brood cells (Marchi & Melo, 2010; Pereira & Garófalo, 2010). Brood cells are barrel-shaped (Pereira & Garófalo, 2010), as in species of *Manuelia* (Flores-Prado *et al.*, 2008). Cells are aligned (Marchi & Melo, 2010; Pereira & Garófalo, 2010). Females may reuse nests and nest cooperatively with sisters (Camillo & Garófalo, 1989). Guarding behavior, recognition and tolerance of nest males are found too in all xylocopines (Flores-Prado *et al.*, 2010). *Xylocopa* is distributed in all Neotropical region. Subgenera are identified following Silveira *et al.* (2002) and Michener (2007). Species from São Paulo (Brazil) are identified following the key in Marchi & Alves-dos-Santos (2013).

COLLETINAE. Known for polyester brood-cell lining, wood nesting probably arose once in the subfamily along with multiple reversals to soil nesting (Almeida, 2008). Of the Neotropical aboveground nesting lineages, only Xeromelissini was not sampled with trap-nests. Nesting behavior in this subfamily was revised by Almeida (2008).

Colletini. This tribe comprises three Neotropical genera, *Hemicotelles* (2 species), *Rhynchocolletes* (12 species) and *Xanthocolletes* (11 species) plus the cosmopolitan *Colletes* (330 species, 108 Neotropical species) (Michener, 2007). The nesting substrates of *Hemicotelles* and *Xanthocolletes* are unknown (Michener, 2007). One species of *Rhynchocolletes* was recently sampled with trap-nests (Diniz, 2010). *Colletes* was sampled only once with trap-nests in Neotropical region. Most species nests in soil, some in stem pith or existing cavities (Almeida, 2008). Soil nesting *Colletes* have linear cells and they lack basitibial and pygidial plates, for this the cavity nesting behavior may be primitive (Almeida, 2008). Otherwise, trap-nest biology in Neotropical species is unknown. Ferrari & Silveira (2015) provided a key to species of Colletini of Minas Gerais (Brazil) and Ferrari (2017) for *Colletes* of Chile.

Hylaeini. *Hylaeus* comprises 624 species worldwide of which 111 are Neotropical (Michener, 2007). They nest in existing cavities in a variety of substrates, including wood, pith, rock and soil (Michener, 2007; Almeida, 2008). Nothing is known of the Neotropical species and no identification key is available.

MEGACHILINAE. Most lineages of this subfamily include species that nest above ground, except near root lineages (Gonzalez *et al.*, 2012). Nests may be in the soil, in burrows in the wood, in plant stems, in other cavities, or may be free-standing constructs. Materials used to construct their nest are variable (petals and leaves, resin, nectar, saliva, others). The tribes Anthidiini, Megachilini and Osmiini have been observed carrying material for nesting (Michener, 2007) and all these tribes, along with the Lithurgini, used trap-nests in the Neotropical region.

Anthidiini. This tribe comprises 677 species worldwide and 339 Neotropical species, with 38 (Moure *et al.*, 2013) genera in the neotropics. See Martins *et al.* (2015) for a list of Danuncia Urban's publications that include comparative notes and keys to the species. Anthidiines nest in existing cavities or build exposed nests, while few species excavate soil nests (Michener, 2007). Nests comprise a wide variety of materials, including resin, leaf and flower pieces, plant fibers and pebbles (Michener, 2007). A total of 26 species and 12 genera were reported using trap-nests (Tables 1 and S1). Nest architecture is known for: *Anthidium* (50 Neotropical species), *Anthodioctes* (43 species), *Carloticola* (2 Neotropical species) *Ctenanthidium* (4 species), *Duckeanthidium* (4 species), *Epanthidium* (23 species), while nothing is known for half of the genera: *Anthidulum* (7 species), *Dicrananthidium* (8 species), *Hypanthidium* (20 species), *Loyolanthidium* (8 Neotropical species), *Nananthidium* (13 species) and *Saranthidium* (10 species) (Moure *et al.*, 2013, Table 1, Table S1).

Anthidium nests partitions and plugs include plant trichomes and other material such as fruits, seeds, leaves, small rocks, wood chips and detritus (Vitale *et al.*, 2017). Entrance diameter varies from 5-11 mm (Vitale *et al.*, 2017). Brood cells are linear, except for *Anthidium vigintipunctatum* Friese, 1908, which cells were perpendicular or oblique. Brood cells vary from 1-19 (Vitale *et al.*, 2017). The genus occurs from Argentina to Mexico (Moure *et al.*, 2013). Gonzalez & Griswold (2013) provided a key to species.

Anthodioctes cells and closing plug are made with resins, sometimes mixed with mud or wood chips (Morato, 2001a; Alves-dos-Santos, 2004; Camarotti-de-Lima & Martins, 2005). Entrance diameter ranges from 5-11 mm (Morato, 2001a; Camarotti-de-Lima & Martins, 2005). The 2-11 brood cells are linear, some with vestibules and preliminary plugs (Morato, 2001a; Alves-dos-Santos, 2004; Camarotti-de-Lima & Martins, 2005). *Anthodioctes megachiloides* Holmberg 1903 nests sometimes have intercalary cells and distal empty space (Alves-dos-Santos, 2004). The genus occurs from Mexico to Argentina (Moure *et al.*, 2013). Identification of its species is possible following Urban (1999, 2002, 2003, 2004).

Carloticola nests and cells partitions are of clay or sand mixed with resin (Mello, 2014). Entrance diameter varies from 6-10 mm and nests have 3-6 brood cells. Vestibular cells are filled with flower buds (Asteraceae or Malpighiaceae) and a preliminary plug is present (Mello, 2014). The genus occurs in Argentina, Brazil and Paraguay (Moure *et al.*, 2013). Identification follows Moure & Urban (1990).

Ctenanthidium, a single nest of *Ctenanthidium bifasciatum* Urban, 1993 was described by Alvarez *et al.* (2015). Resin covered the inner walls, brood cells and partitions. The entrance was 4.0 mm in diameter. The nest had three serial brood cells between an empty space in the distal end of the nest and a closing plug. The genus is found in Argentina, Bolivia, Brazil and Uruguay (Moure *et al.*, 2013). Identification follows Urban (1991).

Duckeanthidium cells are made with glandular substance and plant resins (Thiele, 2002). Entrance diameter varies from 11-13 mm. Nests have 1-3 brood cells, 15-45 mm from the preliminary plug. Partitions and closing plug were extremely hard, with a resin-like material in the closing plug. The genus occurs from Costa Rica to Brazil (Moure *et al.*, 2013). Comparative notes about taxonomy of the species can be found in Michener (2002) and Urban (2004).

Epanthidium nests are of resin mixed with mud or sand (Gomes, 2016) with 2-9 linear brood cells, occasionally with vestibules. The genus occurs from Mexico to Argentina (Moure *et al.*, 2013) and information to identify its species can be found in Urban (1992, 2006, 2011).

Lithurgini. This tribe comprises two South American genera, *Microthurge* (4 species) and *Trichothurgus* (14 species) plus two genera with Neotropical species, *Lithurgopsis* (5 Neotropical species) and *Lithurgus* (1 Neotropical species), these species nest in dead, dry, decomposing wood, making nests without cell linings and often without partitions (Michener, 2007; Moure *et al.*, 2013). *Microthurge corumbae* (Cockerell, 1901) can reuse nests made by one or more females in which guarding occurs (Garófalo *et al.*, 1992). *Trichothurgus laticeps* (Friese, 1906) build nests with wood particles and pollen (Vitale & Vázquez, 2017). Entrance diameter varying from 8-11 mm and nests are without a closing plug and most are without cell partitions. Females excavate wood with mandibles (Vitale & Vázquez, 2017). *Trichothurgus* occurs in Argentina, Chile and Peru (Moure *et al.*, 2013), and an identification key is provided by Michener (1983).

Megachilini. This tribe has two Neotropical genera: *Megachile* and the cleptoparasite *Coelioxys*. *Megachile* comprises 1,093 species, of which 431 are Neotropical (Michener, 2007; Moure *et al.*, 2013), with 32 Neotropical subgenera. Nests are built, often with petals and leaves, in existing cavities in soil, wood, and man-made objects, while some may be free-standing construction (Michener, 2007). Seventeen subgenera and 33 species were studied using trap-nests. *Megachile* builds with leaves, petals, soil, mud and pebbles (Torretta & Durante, 2011; Torretta *et al.*, 2012, 2014, Marques & Gaglianone, 2013; Sabino & Antonini, 2017). Entrance diameter varies from 6-27 mm (Torretta & Durante, 2011; Torretta *et al.*, 2012, 2014; Marques & Gaglianone, 2013; Sabino & Antonini, 2017). One to 16 brood cells tend to be cylindrical and linear (Teixeira *et al.*, 2011; Cardoso & Silveira, 2012; Marques & Gaglianone, 2013; Rocha-Filho & Garófalo, 2016b; Sabino & Antonini, 2017). Preliminary plugs and vestibular cells are uncommon (Torretta & Durante, 2011; Teixeira *et al.*, 2011; Cardoso & Silveira, 2012; Torretta *et al.*, 2012, 2014; Marques & Gaglianone, 2013; Rocha-Filho & Garófalo, 2016b; Sabino & Antonini, 2017). Two species were observed sharing the same nest (Cardoso & Silveira, 2012). *Megachile* occurs from Mexico to Argentina (Moure *et al.*, 2013). The *Megachile* subgenera can be identified following Michener (2007) and Silveira *et al.* (2002). A key to the Neotropical species is not available.

Osmiini. Twenty genera are recognized in this tribe, four of which are found in Mexico and Central America (*Ashmeadiella*, *Atoposmia*, *Heriades* and *Osmia*; Michener, 2007). Osmiini nest in the soil and in existing cavities (Rozen *et al.*, 2010; Rozen-Jr. & Praz, 2016), using a variety of materials, including petals, mud, pebbles and resins (Praz *et al.*, 2008). Osmiini from the New World tend to nest aboveground more than those from the Old World (Praz *et al.*, 2008), however, the group lacks phylogenetic consensus which is necessary for a proper reconstruction of nesting behavior (Praz *et al.*, 2008; Gonzalez *et al.*, 2012). One *Heriades* morphospecies was reported in trap-nests in the Neotropical region without nest information (Roubik & Villanueva-Gutiérrez, 2009). An identification subgenera's key is provided by Michener (2007).

DISCUSSION

We reviewed 109 trap-nesting studies in the Neotropical region that included information for 140 species and 24 genera. Garófalo *et al.* (2004) listed 57 species for Brazil while we report 90 Brazilian species demonstrating rapid advancement in the study of this group of bees. Some of them are studied more often than others. The Apinae are the most studied, with data for all genera sampled in trap-nests. Conversely, trap-nesting Colletinae have not been reported, and the taxonomy of this subfamily has received little attention (see below). Megachilinae are intermediate, with many nests of the Anthidiini being properly described. Their taxonomy is relatively well-understood (Table 1).

We found that most studies were from Brazil and they were restricted to about half of the states, mostly southeastern (Freitas *et al.*, 2009). The other Neotropical countries continue to be very poorly studied. Argentina, Ecuador and Colombia have several studies and may be considered to have quite understanding of their trap-nesting bees biodiversity. In Central America, Costa Rica is better sampled (*e.g.*, Coville & Coville, 1980; Thiele, 2005; Vinson *et al.*, 2010), followed by Panama.

The details of nest architecture are similar to most of the taxa with respect to construction material, closing plug and cell shape (Table 2), while other structures may vary among lineages and intraspecifically. For example, entrance diameter is correlated with female body size ensuring the fit of brood cells (Krombein, 1967; Maclvor, 2017). Diameter of trap-nests also seems to be due to female choice, in part, and varies widely among studies (Coville, 1982; Maclvor, 2017). The number of brood cells is also variable and associated with resource availability, sex ratio of progeny and of course, nest length (Coville, 1982; Morato & Martins, 2006). Vestibular and intercalary cells and the preliminary plug also vary widely among trap-nesting bees (Table 2). Vestibular cells vary from occasional to common (Krombein, 1967; Asís *et al.*, 2007), and while they have been suggested to be a defense against parasites (Krombein, 1967; Coville & Coville, 1980), their true purpose has not been experimentally tested (Asís *et al.*, 2007). Also, vestibular cells often remain unmentioned in studies and so their presence or

absence in those studies is unclear. As an important nest feature, we recommend that presence and absence of the vestibular cells always be clearly stated. Intercalary cells and the preliminary plug are seldom observed, they may be used in the defense against parasites or to change the conformation of the inner end of the boring, respectively (Krombein, 1967; O'Neill, 2001).

Taxonomic information was sufficient for only 14 genera, more than half of which are in the Anthidiini. This well-known tribe was studied extensively by Urban (see bibliography in Martins *et al.*, 2015). Essentially all the remaining groups are poorly studied taxonomically (Table 1). Relatively well known genera include *Centris*, and *Xylocopa*, whose species can often be identified by specialists, but comprehensive keys are not yet available. *Hylaeus* and *Megachile* remain poorly studied with many unnamed species (Moure *et al.*, 2013). *Tetrapedia* is also poorly studied but not so diverse as *Hylaeus* and *Megachile* (Moure *et al.*, 2013).

Taxonomical impediment is an important issue in many insect studies (Oliveira *et al.*, 2011; Begum *et al.*, 2011; Jordaens *et al.*, 2013), including in trap-nesting bee studies Araujo *et al.*, 2017; Matos *et al.*, 2016; Iantas *et al.*, 2017). Reliable identification is necessary when using bees as ecological indicators (Tscharrntke *et al.*, 1998) even if morphospecies can be used in some diversity metrics and analyses (Magurran, 2004; Tylianakis *et al.*, 2007; Pereira-Peixoto *et al.*, 2014; Araújo *et al.*, 2016; Iantas *et al.*, 2017). Studies of phylogenetic and beta diversity, temporal and spatial distribution patterns and nesting behaviors are all hampered by problems with species identification (Faith, 1992; Magurran, 2004; Mittelbach *et al.*, 2007; Faith, 2015).

CONCLUSIONS

Herein we summarized biological and taxonomic knowledge of trap-nest bees in the Neotropical region. Priorities for future research must be settled to fill the more important gaps. For example, researchers should clearly provide details of nest architecture, including clear statements about the presence and absence of structures that can be considered as characters. We propose that researchers provide details of building material, entrance diameter (if oval, diameter along both axes), cell arrangement, number and shapes of brood cells, presence/absence of vestibular and intercalary cells, preliminary and closing plugs and the back empty space. Some of Neotropical trap-nesting groups require further taxonomical work, but especially *Hylaeus* and *Megachile* seeing of the absence of modern taxonomic studies. *Megachile* has an additional requirement since its high abundance in trap-nests studies.

ACKNOWLEDGMENTS

Camila C.F. Costa is funded by CAPES (*Coordenação de Aperfeiçoamento de Pessoal de Nível Superior*) and

Rodrigo B. Gonçalves by CNPq (*Conselho Nacional de Desenvolvimento Científico e Tecnológico*). We thank Carlos Garófalo, Gabriel Melo and John Lattke for comments on a previous version of the manuscript. Special thanks to Anderson Lepeco for the nest drawing. James J. Roper, reviewed the English and provided additional suggestions.

REFERENCES

- Aguiar, C.M.L. & Garófalo, C.A. 2004. Nesting biology of *Centris* (*Hemisiella*) *tarsata* Smith (Hymenoptera, Apidae, Centridini). *Revista Brasileira de Zoologia*, 21(3): 477-486. DOI
- Aguiar, C.M.L.; Garófalo, C.A. & Almeida, G.F. 2006. Biologia de nidificação de *Centris* (*Hemisiella*) *trigonoides* Lepeletier (Hymenoptera, Apidae, Centridini). *Revista Brasileira de Zoologia*, 23(2): 323-330. DOI
- Almeida, E.A.B. 2008. Colletidae nesting biology (Hymenoptera: Apoidea). *Apidologie*, 39(1): 16-29. DOI
- Alvarez, L.J.; Lucia, M.; Durante, S.P.; Lilian, M. & Gonzalez, V.H. 2015. Description of the male, new geographical record, and biological notes on the small resin bee *Hypanthidioides* (*Ctenanthidium*) *bifasciata* (Urban, 1993) (Hymenoptera: Megachilidae). *Pan-Pacific Entomologist*, 91(2): 184-188. DOI
- Alves-dos-Santos, I. 2004. Biologia de nidificação de *Anthodioctes megachiloides* Holmberg (Anthidiini, Megachilidae, Apoidea). *Revista Brasileira de Zoologia*, 21(4): 739-744. DOI
- Alves-dos-Santos, I.; Melo, G.A.R. & Rozen, J.G. 2002. Biology and Immature Stages of the Bee Tribe Tetrapediini (Hymenoptera: Apidae). *American Museum Novitates*, (3377): 1-45.
- Araujo, G.J.; Fagundes, R. & Antonini, Y. 2017. Trap-Nesting Hymenoptera and their network with parasites in Recovered Riparian Forests Brazil. *Neotropical Entomology*, 47(1): 26-36.
- Araújo, P.C.S.; Lourenço, A.P. & Raw, A. 2016. Trap-nesting bees in Montane Grassland (Campo Rupestre) and Cerrado in Brazil: Collecting generalist or specialist nesters. *Neotropical Entomology*, 45(5): 482-489.
- Asís, J.D.; Benítez, A.; Tormos, J.; Gayubo, S.F. & Tomé, M.A.M. 2007. The significance of the vestibular cell in trap nesting wasps (Hymenoptera: Crabronidae): Does its presence reduce mortality? *Journal of Insect Behavior*, 20(2): 289-305. DOI
- Augusto, S.C. & Garófalo, C.A. 2004. Nesting biology and social structure of *Euglossa* (*Euglossa*) *townsendi* Cockerell (Hymenoptera, Apidae, Euglossini). *Insectes Sociaux*, 51(4): 400-409. DOI
- Augusto, S.C. & Garófalo, C.A. 2009. Bionomics and sociological aspects of *Euglossa fimbriata* (Apidae, Euglossini). *Genetics and Molecular Research*, 8(2): 525-538. DOI
- Augusto, S.C. & Garófalo, C.A. 2011. Task allocation and interactions among females in *Euglossa carolina* nests (Hymenoptera, Apidae, Euglossini). *Apidologie*, 42(2): 162-173. DOI
- Begum, S.; Anis, S.B.; Farooqi, M.K.; Rehmat, T. & Fatma, J. 2011. Aphelinid parasitoids (Hymenoptera; Aphelinidae) of whiteflies (Homoptera: Aleyrodidae) from India. *Biology and Medicine*, 3(2 special issue): 222-231.
- Bossert, S.; Murray, E.A.; Almeida, E.A.B.; Brady, S.G.; Blaimer, B.B. & Danforth, B.N. 2019. Combining transcriptomes and ultraconserved elements to illuminate the phylogeny of Apidae. *Molecular Phylogenetics and Evolution*, 130(2019): 121-131. DOI
- Branstetter, M.G.; Danforth, B.N.; Pitts, J.P.; Faircloth, B.C.; Ward, P.S.; Buffington, M.L.; Gates, M.W.; Kula, R.R. & Brady, S.G. 2017. Phylogenomic insights into the evolution of stinging wasps and the origins of ants and bees. *Current Biology*, 27(7): 1019-1025. DOI
- Buschini, M.L.T. 2006. Species diversity and community structure in trap-nesting bees in Southern Brazil. *Apidologie*, 37: 58-66. DOI

- Buschini, M.L.T. & Wolff, L.L. 2006. Nesting biology of *Centris (Hemisiella) tarsata* Smith in Southern Brazil (Hymenoptera, Apidae, Centridini). *Brazilian Journal of Biology*, 66(4): 1091-1101. [DOI](#)
- Camarotti-de-Lima, M. de F. & Martins, C.F. 2005. Biologia de nidificação e aspectos ecológicos de *Anthodiocetes lunatus* (Smith) (Hymenoptera: Megachilidae, Anthidiini) em área de tabuleiro nordestino, PB. *Neotropical Entomology*, 34(3): 375-380.
- Camillo, E. 2005. Nesting biology of four *Tetrapedia* species in trap nests (Hymenoptera: Apidae: Tetrapedini). *Revista de Biologia Tropical*, 53: 175-186.
- Camillo, E. & Garófalo, C.A. 1989. Social organization in reactivated nests of three species of *Xylocopa* (Hymenoptera, Anthophoridae) in southeastern Brazil. *Insectes Sociaux*, 36(2): 92-105.
- Camillo, E.; Garófalo, C.A.; Serrano J.C. & Mucilo, G. 1995. Diversidade e abundância sazonal de abelhas e vespas solitárias em ninhos-armadilha (Hymenoptera, Apocrita, Aculeata). *Revista Brasileira de Entomologia*, 39: 459-470.
- Cardinal, S.; Straka, J. & Danforth, B.N. 2010. Comprehensive phylogeny of apid bees reveals the evolutionary origins and antiquity of cleptoparasitism. *Proceedings of the National Academy of Sciences*, 107(37): 16207-16211. [DOI](#)
- Cardoso, C.F. & Silveira, F.A. 2012. Nesting biology of two species of *Megachile (Moureapis)* (Hymenoptera: Megachilidae) in a semideciduous forest reserve in southeastern Brazil. *Apidologie*, 43(1): 71-81. [DOI](#)
- Carvalho, G.C.A.; Carreira, L.M.M.; Rêgo, M.M.C. & Albuquerque, P.M.C. 2016. Nesting habits of *Centris (Hemisiella) dichrotricha* (Hymenoptera: Apidae) in the Northern Cerrado of Brazil. *Revista de Biologia Tropical*, 64(3): 1041-1056. [DOI](#)
- Coville, R.E. 1982. *Wasps of the genus Trypoxylon subgenus Trypargilum in North America*. Berkeley, University of California Press.
- Coville, R.E. & Coville, P.L. 1980. Nesting biology and male behavior of *Trypoxylon (Trypargilum) tenocitlan* in Costa Rica (Hymenoptera: Sphecidae). *Annals of the Entomological Society of America*, 73(1): 110-119. [DOI](#)
- Daly, H.V.; Michener, C.D.; Moure, J.S. & Sakagami, S.F. 1987. The relictual bee genus *Manuelia* and its relation to other Xylocopinae (Hymenoptera: Apoidea). *Pan-Pacific Entomologist*, 63: 102-124.
- Diniz, M.E. dos R. 2010. *Biologia de Nidificação de Rhynchocolletes sp. (Hymenoptera; Apoidea; Colletidae; Colletinae)*. (Undergraduated Thesis). Universidade Estadual do Centro-Oeste, Guarapuava, Brazil.
- Drummond, P.; Silva, F.O. & Viana, B.F. 2008. Ninhos de *Centris (Heterocentris) terminata* Smith (Hymenoptera: Apidae, Centridini) em Fragmentos de Mata Atlântica Secundária, Salvador, BA. *Neotropical Entomology*, 37(3): 239-246.
- Eickwort, G.C. 1975. Gregarious nesting of the mason bee *Hoplitis anthocopoides* and the evolution of parasitism and sociality among megachilid bees. *Evolution*, 29(1): 142-150. [DOI](#)
- Faith, D.P. 1992. Systematics and conservation: on predicting the feature diversity of subsets of taxa. *Cladistics*, 8: 361-373. [DOI](#)
- Faith, D.P. 2015. Phylogenetic diversity, functional trait diversity and extinction: avoiding tipping points and worst-case losses. *Philosophical Transactions of the Royal Society of London, Series B, Biological Sciences*, 370(1662): 1-10. [DOI](#)
- Faria, L.R.R. & Melo, G.A.R. 2012. Species of *Euglossa* of the analis group in the Atlantic forest (Hymenoptera, Apidae). *Zoologia (Curitiba)*, 29(4): 349-374. [DOI](#)
- Faria-Jr.; L.R.R. & Melo, G.A.R. 2007. Species of *Euglossa (Glossura)* in the Brazilian Atlantic forest, with taxonomic notes on *Euglossa stellfeldi* Moure (Hymenoptera, Apidae, Euglossina). *Revista Brasileira de Entomologia*, 51(3): 275-284. [DOI](#)
- Ferrari, R.F. & Silveira, F.A. 2015. The species of Colletinae (Hymenoptera: Colletidae) in the Brazilian state of Minas Gerais. *Zootaxa*, 3914(3): 245-274.
- Ferrari, R.R. 2017. Taxonomic revision of the species of *Colletes* Latreille, 1802 (Hymenoptera: Colletidae: Colletinae) found in Chile. *Zootaxa*, 4364(1): 1-137.
- Flores-Prado, L.; Chiappa, E. & Niemeyer, H.M. 2008. Nesting biology, life cycle, and interactions between females of *Manuelia postica*, a solitary species of the Xylocopinae (Hymenoptera: Apidae). *New Zealand Journal of Zoology*, 35(1): 93-102. [DOI](#)
- Flores-Prado, L.; Flores, S.V. & McAllister, B.F. 2010. Phylogenetic relationships among tribes in Xylocopinae (Apidae) and implications on nest structure evolution. *Molecular Phylogenetics and Evolution*, 57(1): 237-244.
- Freiria, G.A.; Garófalo, C.A. & Del Lama, M.A. 2017. The primitively social behavior of *Euglossa cordata* (Hymenoptera, Apidae, Euglossini): a view from the perspective of kin selection theory and models of reproductive skew. *Apidologie*, 48(4): 523-532. [DOI](#)
- Freitas, B.M.; Imperatriz-Fonseca, V.L.; Medina, L.M.; Kleinert, A. de M.P.; Galetto, L.; Nates-Parra, G. & Quezada-Euán, J.J.G. 2009. Diversity, threats and conservation of native bees in the Neotropics. *Apidologie*, 40(3): 332-346. [DOI](#)
- Garófalo, C.A.; Camillo, E.; Augusto, S.C.; Jesus, B.M.V. & Serrano, J.C. 1998. Nest structure and communal nesting in *Euglossa (Glossura) annexans* Dressler (Hymenoptera, Apidae, Euglossini). *Revista Brasileira de Zoologia*, 15(3): 589-596. [DOI](#)
- Garófalo, C.A.; Camillo, E.; Campos, M.J.O. & Serrano, J.C. 1992. Nest re-use and communal nesting in *Microthurga corumbae* (Hymenoptera, Megachilidae), with special reference to nest defense. *Insectes Sociaux*, 39(3): 301-311.
- Garófalo, C.A.; Martins, C.F. & Alves-dos-Santos, I. 2004. The Brazilian solitary bee species caught in trap nests. In: Freitas, B.M. & Pereira, J.O. *Solitary Bees conservation, rearing and management for pollination*. Fortaleza, Imprensa Universitária. p. 77-84.
- Gess, S.K. & Gess, F.W. 2008. Patterns of usage of snail shells for nesting by wasps (Vespidae: Masarinae and Eumeninae) and bees (Megachilidae: Megachilinae) in Southern Africa. *Journal of Hymenoptera Research*, 17(1): 86-109.
- Gomes, A.M.S. 2016. *Bionomia e compartimento de nidificação da abelha Epanthidium tigrinum (Hymenoptera-Megachilidae) em ninhos-armadilha*. (Master Dissertation). Universidade Federal do Ceará, Fortaleza, Brazil.
- Gonzalez, V. & Griswold, T. 2013. Wool carder bees of the genus *Anthidium* in the Western Hemisphere (Hymenoptera: Megachilidae): diversity, host plant associations, phylogeny, and biogeography. *Zoological Journal of the Linnean Society*, 168(2): 221-425. [DOI](#)
- Gonzalez, V.H.; Griswold, T.; Praz, C.J. & Danforth, B.N. 2012. Phylogeny of the bee family Megachilidae (Hymenoptera: Apoidea) based on adult morphology. *Systematic Entomology*, 37(2): 261-286. [DOI](#)
- Hedtke, S.M.; Patiny, S. & Danforth, B.N. 2013. The bee tree of life: A supermatrix approach to apoid phylogeny and biogeography. *BMC Evolutionary Biology*, 13(138): 1-13. [DOI](#)
- Iantas, J.; Gruchowski-Woitowicz, F.C. & Tunes-Buschini, M.L. 2017. Habitat modification and alpha-beta diversity in trap nesting bees and wasps (Hymenoptera: Aculeata) in southern Brazil. *Tropical Zoology*, 30(2): 83-96. [DOI](#)
- Jordaens, K.; Sonet, G.; Richet, R.; Dupont, E.; Braet, Y. & Desmyter, S. 2013. Identification of forensically important *Sarcophaga* species (Diptera: Sarcophagidae) using the mitochondrial COI gene. *International Journal of Legal Medicine*, 127(2): 491-504.
- Kamke, R.; Zillikens, A.; Heinle, S. & Steiner, J. 2008. Natural enemies and life cycle of the orchid bee *Eufriesea smaragdina* (Hymenoptera: Apidae) reared from trap nests. *Journal of the Kansas Entomological Society*, 81(2): 101-109. [DOI](#)
- Kimsey, L.S. 1982. *Systematics of bees of the genus Eufriesea (Hymenoptera, Apidae)*. Berkeley, University of California Press.

- Krombein, K. 1967. *Trap-nesting wasps and bees. Life-histories, nests and associates*. Washington. Smithsonian Press.
- Lucia, M.; Telleria, M.C.; Ramello, P.J. & Abrahamovich, A.H. 2017. Nesting ecology and floral resource of *Xylocopa augusti* Lepeletier de Saint Fargeau (Hymenoptera, Apidae) in Argentina. *Agricultural and Forest Entomology*, 19(3): 281-293. DOI
- MacIvor, J.S. 2017. Cavity-nest boxes for solitary bees: a century of design and research. *Apidologie*, 48(3): 311-327. DOI
- MacIvor, J.S. & Packer, L. 2015. "Bee hotels" as tools for native pollinator conservation: A premature verdict? *PLoS ONE*, 10(3): 1-13.
- Magurran, A.E. 2004. *Measuring Biological Diversity*. Oxford, Blackwell Science.
- Marchi, P. & Alves-dos-Santos, I. 2013. As abelhas do gênero *Xylocopa* Latreille (Xylocopini, Apidae) do Estado de São Paulo, Brasil. *Biota Neotropica*, 13(2): 249-269.
- Marchi, P. & Melo, G.A.R. 2010. Biologia de nidificação de *Xylocopa* (*Neoxylocopa*) *frontalis* (Olivier) (Hymenoptera, Apidae, Xylocopini) em Morretes, Paraná. *Oecologia Australis*, 14(1): 210-231. DOI
- Marques, M.F. & Gaglianone, M.C. 2013. Biologia de nidificação e variação altitudinal na abundância de *Megachile* (*Melanosarus*) *nigripennis* Spinola (Hymenoptera, Megachilidae) em um inselbergue na mata Atlântica, Rio de Janeiro. *Bioscience Journal*, 29(1): 198-208.
- Martins, A.C. & Melo, G.A.R. 2016. The New World oil-collecting bees *Centris* and *Epicharis* (Hymenoptera, Apidae): Molecular phylogeny and biogeographic history. *Zoologica Scripta*, 45(1): 22-33. DOI
- Martins, A.C.; Melo, G.A.R. & Renner, S.S. 2014. The corbiculate bees arose from New World oil-collecting bees: Implications for the origin of pollen baskets. *Molecular Phylogenetics and Evolution*, 80: 88-94.
- Martins, A.C.; Santos, L.M. & Molin, A.D. 2015. Danúncia Urban: a life devoted to Entomology. In: Aguiar, A.J.C. Gonçalves, R.B. & Ramos, K.S. *Ensaio sobre as abelhas da Região Neotropical: Homenagem aos 80 anos de Danúncia Urban*. Editora UFPR, Curitiba, p. 11-46.
- Matos, M.C.B.; Silva, S.S. & Teodoro, A.V. 2016. Seasonal population abundance of the assembly of solitary wasps and bees (Hymenoptera) according to land-use in Maranhão state, Brazil. *Revista Brasileira de Entomologia*, 60: 171-176.
- Mello, B.N. da S. 2014. *Nicho de abelhas Megachilidae (Hymenoptera, Apoidea) em floresta ombrófila na Mata Atlântica: dinâmica temporal, relações tróficas e uso de recursos de nidificação*. (Master Dissertation). Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, Brazil.
- Melo, G.A.R. 1999. Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on the crabronid wasps. *Scientific Papers, Natural History Museum of the University of Kansas*, 14: 1-55.
- Melo, G.A.R. & Gonçalves, R.B. 2005. Higher-level bees classifications (Hymenoptera, Apoidea, Apidae *sensu lato*). *Revista Brasileira de Zoologia*, 22(1): 153-159. DOI
- Menezes, G.B.; Gonçalves-Esteves, V.; Bastor, E.M.A.F.; Augusto, S.C. & Gaglianone, M.C. 2012. Nesting and use of pollen resources by *Tetrapedia diversipes* Klug (Apidae) in Atlantic Forest areas (Rio de Janeiro, Brazil) in different stages of regeneration. *Revista Brasileira de Entomologia*, 56(1): 86-94. DOI
- Michener, C.D. 1983. The classification of the *Lithurginae* (Hymenoptera: Megachilidae). *Pan-Pacific Entomologist*, 59(1-4): 176-187.
- Michener, C.D. 2002. *Duckeanthidium*, a genus new to Central America, with generic synonymy and a new species (Hymenoptera: Megachilidae). *Journal of the Kansas Entomological Society*, 75(4): 233-240.
- Michener, C.D. 2007. *The Bees of the World*. Baltimore, The Johns Hopkins University Press.
- Mittelbach, G.G.; Schemske, D.W.; Cornell, H.V.; Allen, A.P.; Brown, J.M.; Bush, M.B.; Harrison, S.P.; Hurlbert, A.H.; Knowlton, N.; Lessios, H.A.; McCain, C.M.; McCune, A.R.; McDade, L.A.; McPeck, M.A.; Near, T.J.; Price, T.D.; Ricklefs, R.E.; Roy, K.; Sax, D.F.; Schluter, D.; Sobel, J.M. & Turelli, M. 2007. Evolution and the latitudinal diversity gradient: Speciation, extinction and biogeography. *Ecology Letters*, 10(4): 315-331. DOI
- Morato, E.F. 2001a. Biologia e ecologia de *Anthodiocetes moratoi* Urban (Hymenoptera, Megachilidae, Anthidiini) em matas contínuas e fragmentos na Amazônia Central, Brasil Anthidiinae. *Revista Brasileira de Entomologia*, 18(3): 729-736. DOI
- Morato, E.F. 2001b. Efeitos da fragmentação florestal sobre vespas e abelhas solitárias na Amazônia Central. II. Estratificação vertical. *Revista Brasileira de Zoologia*, 18(3): 737-747.
- Morato, E.F. & Martins, R.P. 2006. An overview of proximate factors affecting the nesting behavior of solitary wasps and bees (Hymenoptera: Aculeata) in preexisting cavities in wood. *Neotropical Entomology*, 35(3): 285-298.
- Moure, J.S. & Urban, D. 1990. *Carlaticola* gen. nov. e *Carlaticola trichura* sp. nov. da Argentina (Apoidea, Megachilidae, Anthidiinae). *Acta Biológica Paranaense*, 19(1-4): 89-99.
- Moure, J.S.; Urban, D. & Melo, G.A.R. 2013. *Catalogue of Bees (Hymenoptera, Apoidea) in the Neotropical Region – online version*. Available at: <http://moure.cria.org.br/catalogue>. Access in: 10/04/2018.
- Moure-Oliveira, D.; Rocha-Filho, L.C.; Ferreira-Caliman, M.J. & Garófalo, C.A. 2017. Nesting dynamic and sex allocation of the oil-collecting bee *Centris* (*Heterocentris*) *analís* (Fabricius, 1804) (Apidae: Centridini). *Journal of Natural History*, 51(19-20): 1-18.
- O'Neill, K. 2001. *Solitary Wasps: Behavior and Natural History*. Cornell University Press, Ithaca.
- Oliveira, A.R.; Farinha, A.; Rebelo, M.T. & Dias, D. 2011. Forensic entomology: Molecular identification of blowfly species (Diptera: Calliphoridae) in Portugal. *Forensic Science International: Genetics Supplement Series*, 3(1): 439-440.
- Oliveira, P.S. & Gonçalves, R.B. 2017. Trap-nesting bees and wasps (Hymenoptera, Aculeata) in a Semideciduous Seasonal Forest Fragment, Southern Brazil. *Papéis Avulsos de Zoologia*, 57(13): 149-156. DOI
- Parra-H, A. & Nates-Parra, G. 2009. La arquitectura de nidos de *Euglossa* (*Euglossa*) *hemichlora* (Hymenoptera: Apidae: Euglossini). *Revista Colombiana de Entomología*, 35(2): 283-285. <http://www.scielo.org.co/pdf/rcen/v35n2/v35n2a26.pdf>.
- Pereira, M. & Garófalo, C.A. 2010. Biologia da nidificação de *Xylocopa frontalis* e *Xylocopa griseescens* (Hymenoptera, Apidae, Xylocopini) em ninhos-armadilha. *Oecologia Australis*, 14(1): 193-209.
- Pereira-Peixoto, M.H.; Pufal, G.; Martins, C.F. & Klein, A.-M. 2014. Spillover of trap-nesting bees and wasps in an urban-rural interface. *Journal of Insect Conservation*, 18(5): 815-826.
- Perillo, L.N.; Neves, F.D.S.; Antonini, Y. & Martins, R.P. 2017. Compositional changes in bee and wasp communities along Neotropical mountain altitudinal gradient. *PLoS ONE*, 12(7): 1-14.
- Peruquetti, R.C. 1998. Notes on adults of *Euglossa townsendi* Cockerell (Apidae: Euglossini) reared from a trap nest. *Anais da Sociedade Entomológica do Brasil*, 27(2): 309-311. DOI
- Praz, C.J.; Müller, A.; Danforth, B.N.; Griswold, T.L.; Widmer, A. & Dorn, S. 2008. Phylogeny and biogeography of bees of the tribe Osmiini (Hymenoptera: Megachilidae). *Molecular Phylogenetics and Evolution*, 49: 185-197.
- Rebêlo, J.M.M. & Moure, J.S. 1995. As espécies de *Euglossa* Latreille do nordeste de São Paulo (Apidae, Euglossinae). *Revista Brasileira de Zoologia*, 12(3): 445-466.
- Rocha-Filho, L.C. da & Garófalo, C.A. 2016a. Natural history of *Tetrapedia diversipes* (Hymenoptera: Apidae) in an atlantic semideciduous forest

- remnant surrounded by coffee crops, *Coffea arabica* (Rubiaceae). *Annals of the Entomological Society of America*, 109(2): 183-197. [DOI](#)
- Rocha-Filho, L.C. da & Garófalo, C.A. 2016b. Nesting biology of *Megachile (Chrysosarus) guaranitica* and high mortality caused by its cleptoparasite *Coelioxys bertonii* (Hymenoptera: Megachilidae) in Brazil. *Austral Entomology*, 55(1): 25-31. [DOI](#)
- Rocha-Filho, L.C. da, Rabelo, L.S.; Augusto, S.C. & Garófalo, C.A. 2017. Cavity-nesting bees and wasps (Hymenoptera: Aculeata) in a semi-deciduous Atlantic forest fragment immersed in a matrix of agricultural land. *Journal of Insect Conservation*, 21(4): 727-736.
- Roig-Alsina, A. 2000. Claves para las especies argentinas de *Centris* (Hymenoptera, Apidae), con descripción de nuevas especies y notas sobre distribución. *Revista del Museo Argentino de Ciencias Naturales, Nueva Serie*, 2(2): 171-193.
- Roubik, D.W. & Villanueva-Gutiérrez, R. 2009. Invasive Africanized honey bee impact on native solitary bees: A pollen resource and trap nest analysis. *Biological Journal of the Linnean Society*, 98(1): 152-160. [DOI](#)
- Rozen, J.G.; Özbek, H.; Ascher, J.S.; Sedivy, C.; Praz, C.; Monfared, A. & Müller, A. 2010. Nests, Petal Usage, Floral Preferences, and Immatures of *Osmia (Ozbekosmia) avosetta* (Megachilidae: Megachilinae: Osmiini), Including Biological Comparisons with Other Osmiine Bees. *American Museum Novitates*, (3680): 1-22.
- Rozen-Jr.; J.G. & Praz, C.J. 2016. Mature Larvae and Nesting Biologies of Bees Currently Assigned to the Osmiini (Apoidea: Megachilidae). *American Museum Novitates*, 27(3864): 1-46.
- Sabino, W. de O. & Antonini, Y. 2017. Nest architecture, life cycle, and natural enemies of the neotropical leafcutting bee *Megachile (Moureapis) maculata* (Hymenoptera: Megachilidae) in a Montane Forest. *Apidologie*, 48(4): 450-460. [DOI](#)
- Santos, A.K.G.; Warkentin, M.; Silva, J.V.; Barbosa, R.N. & Mouga, M.D.S. 2016. Nesting patterns of bees (Apidae) in brick walls in southern Brazil. *Acta Biologica Catarinense*, 3(2): 80-90. [DOI](#)
- Sheffield, C.S. 2017. Unusual nesting behavior in *Megachile (Eutricharaea) rotundata* (Hymenoptera: Megachilidae). *Journal of Melittology*, (69): 1-6. [DOI](#)
- Sheffield, C.S.; Ratti, C.; Packer, L. & Griswold, T. 2011. Leafcutter and mason bees of the genus *Megachile* Latreille (Hymenoptera: Megachilidae) in Canada and Alaska. *Canadian Journal of Arthropod Identification*, (18): 1-107.
- Silva, C.I. da; do Nascimento Castro, M.M.; dos Santos, I.A. & Garófalo, C.A. 2016. High prevalence of *Miconia chamissois* (Melastomataceae) pollen in brood cell provisions of the orchid bee *Euglossa townsendi* in São Paulo State, Brazil. *Apidologie*, 47(6): 855-866.
- Silveira, F.A.; Melo, G.A.R. & Almeida, E.A.B. 2002. *Abelhas brasileiras: Sistemática e identificação*. Belo Horizonte, Probio/PNUD. 253p.
- Snelling, R.R. 1984. Studies on the taxonomy and distribution of American Centridini bees (Hymenoptera: Anthophoridae). *Contributions in Science Natural History Museum of Los Angeles*, 347: 1-69.
- Stangler, E.S.; Hanson, P.E. & Steffan-Dewenter, I. 2015. Interactive effects of habitat fragmentation and microclimate on trap-nesting Hymenoptera and their trophic interactions in small secondary rainforest remnants. *Biodiversity and Conservation*, 24(3): 563-577. [DOI](#)
- Stangler, E.S.; Hanson, P.E. & Steffan-Dewenter, I. 2016. Vertical diversity patterns and biotic interactions of trap-nesting bees along a fragmentation gradient of small secondary rainforest remnants. *Apidologie*, 47(4): 527-538. [DOI](#)
- Teixeira, F.M.; Schwartz, T.A.C. & Gaglianone, M.C. 2011. Biologia da nidificação de *Megachile (Moureapis) benigna* Mitchell. *EntomoBrasilis*, 4(3): 92-99.
- Thiele, R. 2002. Nesting biology and seasonality of *Duckeanthidium thielei* Michener (Hymenoptera: Megachilidae), an oligolectic Rainforest bee. *Journal of the Kansas Entomological Society*, 75(4): 274-282.
- Thiele, R. 2003. A review of Central American *Centris* (*Heterocentris*) and evidence for male dimorphism in *C. labrosa* (Hymenoptera, Apidae). *Deutsche Entomologische Zeitschrift*, 50(2): 237-242.
- Thiele, R. 2005. Phenology and nest site preferences of wood-nesting bees in a Neotropical lowland rain forest. *Studies on Neotropical Fauna and Environment*, 40(1): 39-48.
- Torretta, J.P. & Durante, S.P. 2011. Nesting ecology of *Megachile (Sayapis) mendozana* Cockerell and its synonymy with *Megachile (Sayapis) santiaguensis* Durante (Hymenoptera: Megachilidae). *Zootaxa*, 3008: 61-68.
- Torretta, J.P.; Durante, S.P. & Basilio, A.M. 2014. Nesting ecology of *Megachile (Chrysosarus) catamarcensis* Schrottky (Hymenoptera: Megachilidae), a *Prosopis*-specialist bee. *Journal of Apicultural Research*, 53(5): 590-598.
- Torretta, J.P.; Durante, S.P.; Colombo, M.G. & Basilio, A.M. 2012. Nesting biology of the leafcutting bee *Megachile (Pseudocentron) gomphrenoides* (Hymenoptera: Megachilidae) in an agro-ecosystem. *Apidologie*, 43(6): 624-633. [DOI](#)
- Tscharntke, T.; Gathmann, A. & Steffan-Dewenter, I. 1998. Bioindication using trap-nesting bees and wasps and their natural enemies: community structure and interactions. *Journal of Applied Ecology*, 35(5): 708-719.
- Tylianakis, J.M.; Tscharntke, T. & Lewis, O.T. 2007. Habitat modification alters the structure of tropical host-parasitoid food webs. *Nature*, 445: 202-205. [DOI](#)
- Urban, D. 1991. *Ctenanthidium*, gen. nov. de *Dianthidiini* com quatro espécies novas da América do Sul (Hymenoptera, Megachilidae). *Revista Brasileira de Zoologia*, 8(1-4): 85-93.
- Urban, D. 1992. Espécies novas de *Epanthidium* Moure (Hymenoptera, Megachilidae, Anthidiinae). *Acta Biológica Paranaense*, 21(1-4): 1-21.
- Urban, D. 1999. Espécies novas e notas sobre *Anthodioctes* Holmberg (Hymenoptera, Apoidea, Megachilidae). *Revista Brasileira de Zoologia*, 16(1): 135-169.
- Urban, D. 2002. Espécies novas de *Anthodioctes* Holmberg e notas sobre alguns *Dianthidiini* (Hymenoptera, Megachilidae). *Revista Brasileira de Zoologia*, 19(2): 151-158.
- Urban, D. 2003 (2005). Novos *Dianthidiini* neotropicais (Hymenoptera, Megachilidae). *Acta Biológica Paranaense*, 32 (1-4): 95-106.
- Urban, D. 2004. Espécies novas e notas sobre *Dianthidiini* (Hymenoptera, Megachilidae) do Peru. *Revista Brasileira de Entomologia*, 48(3): 347-352. [DOI](#)
- Urban, D. 2006. Espécies novas de *Epanthidium* Moure e de *Austrostelis* Michener & Griswold (Hymenoptera, Apidae, Megachilinae). *Revista Brasileira de Entomologia*, 50(1): 43-48.
- Urban, D. 2011. *Melostelis* gen. nov., espécies novas e notas complementares sobre *Anthidiini* (Hymenoptera, Apidae). *Revista Brasileira de Entomologia* 55 (2): 219-225.
- Vélez, D.; Vivallo, F. & Silva, D.P. 2017. Nesting biology and potential distribution of an oil-collecting Centridine Bee from South America. *Apidologie*, 48(2): 181-193. [DOI](#)
- Viana, B.F.; Neves, E.L. & Silva, F.O. 2001. Aspectos da biologia de nidificação de *Euplusia mussitans* (Fabricius) (Hymenoptera, Apidae, Euglossini). *Revista Brasileira de Zoologia*, 18(4): 1081-1087.
- Vinson, S.B.; Frankie, G. & Cónsoli, R. 2010. Description, comparison and identification of nests of cavity-nesting *Centris* bees (Hymenoptera: Apidae: Centridini) in Guanacaste Province, Costa Rica. *Journal of the Kansas Entomological Society*, 83(1): 25-46. [DOI](#)
- Vinson, S.B.; Frankie, G.W. & Williams, H.J. 1996. Chemical ecology of bees of the genus *Centris* (Hymenoptera: Apidae). *Florida Entomologist*, 79(2): 109-129.
- Vitale, N. & Vázquez, D.P. 2017. Ecology and nesting biology of the wood-boring bee *Trichothurgus laticeps* (Hymenoptera: Megachilidae) in a Monte desert reserve in mid-western Argentina. *Apidologie*, 48(1): 31-40. [DOI](#)
- Vitale, N.; Gonzalez, V.H. & Vázquez, D.P. 2017. Nesting ecology of sympatric species of wool carder bees (Hymenoptera: Megachilidae: *Anthidium*) in South America. *Journal of Apicultural Research*, 56(5): 497-509. [DOI](#)
- Vivallo, F. & Vélez, D. 2016. A synopsis of the subgenus *Centris* (*Hemisiella*) Moure, 1945 (Hymenoptera: Apidae: Centridini) in Colombia, with description of a new species. *Zootaxa*, 4162(1): 107-133.

APPENDICES

Table S1. A summary of trap nesting bee species from Neotropical region. Data was summarized through the end of 2017. (Available) It is when there is at least one published description with details on the architecture using trap-nest methodology; (unavailable) is when information is absence. Abbreviations follow: Argentina: BA = Buenos Aires, LP = La Pampa and MZ = Mendoza. Brazil: AC = Acre, AM = Amazonas, BA = Bahia, CE = Ceará, MA = Maranhão, MG = Minas Gerais, PB = Paraíba, PE = Pernambuco, PR = Paraná, RJ = Rio de Janeiro, RN = Rio Grande do Norte, SC = Santa Catarina and SP = São Paulo. Mexico: QR = Quintana Roo.

Taxon	Distribution records (States or provinces from Argentina, Brazil and Mexico are indicated under parenthesis)	References for records	Trap nest description	References for nest descriptions
APINAE				
Centridini				
<i>Centris (Hemisiella) crassipes</i> Smith, 1874	Jamaica	1	Available	1
<i>Centris (Hemisiella) dichrotricha</i> (Moore, 1945)	Brazil (AC, AM, MA)	2–4	Available	4
<i>Centris (Hemisiella) facialis</i> Mocsáry, 1899	Colombia	5	Unavailable	
<i>Centris (Hemisiella) merrillae</i> Cockerell, 1919	Brazil (AC), Trinidad and Tobago	3, 6	Available	6
<i>Centris (Hemisiella) nitida</i> Smith, 1874	Brazil (AM), Costa Rica	7–9	Available	8, 9
<i>Centris (Hemisiella) tarsata</i> Smith, 1874	Argentina (BA); Brazil (BA, CE, MA, MG, PB, PE, PR, RJ, RN, SC, SP)	10–42	Available	33, 43, 44
<i>Centris (Hemisiella) trigonoides</i> Lepeletier, 1841	Brazil (AM, MG, RN); Costa Rica; Colombia	2, 5, 8, 16, 21	Available	8, 9, 45
<i>Centris (Hemisiella) vittata</i> Lepeletier, 1841	Brazil (MA, MG, SP); Costa Rica	3, 7, 8, 15, 19, 26, 28, 36, 46, 47	Available	8, 9, 47, 48
<i>Centris (Hemisiella) sp.</i>	Brazil (AC)	3	—	
<i>Centris (Heterocentris) adunca</i> Moore, 2003	Brazil (AC)	3	Unavailable	
<i>Centris (Heterocentris) analis</i> (Fabricius, 1804)	Brazil (AC, AM, BA, MG, PB, PE, PR, RJ, SP); Costa Rica; Colombia; Mexico (QR)	2, 3, 5, 7, 8, 12, 13, 15, 22, 24, 25, 27, 28, 32, 36, 38–40, 42, 46, 49–55	Available	8, 54
<i>Centris (Heterocentris) bicornuta</i> Mocsáry, 1899	Brazil (AM, MA); Costa Rica	2, 7, 8, 19, 46	Available	8, 9
<i>Centris (Heterocentris) difformis</i> Smith, 1854	Costa Rica	46	Unavailable	
<i>Centris (Heterocentris) labrosa</i> Friese, 1899	Brazil (SP); Costa Rica	25, 36, 46, 50	Unavailable	
<i>Centris (Heterocentris) terminata</i> Smith, 1874	Brazil (AC, AM, BA, MA, MG, PB)	2, 3, 12, 14, 18, 19, 22, 27, 56, 57	Available	58
<i>Centris (Heterocentris) spp.</i>	Brazil (AC, CE, MG, RJ)	3, 14, 18, 27	—	
<i>Centris spp.</i>	Brazil (SP); Ecuador	25, 59, 60	—	
Euglossini				
<i>Eufriesea auriceps</i> (Friese, 1899)	Brazil (SP)	25, 36	Unavailable	
<i>Eufriesea mussitans</i> (Fabricius, 1787)	Brazil (BA)	35	Unavailable	
<i>Eufriesea purpurata</i> (Mocsáry, 1896)	Brazil (AM)	2	Unavailable	
<i>Eufriesea smaragdina</i> (Perty, 1833)	Brazil (SC)	61	Available	62
<i>Eufriesea surinamensis</i> (Linnaeus, 1758)	Brazil (SP)	25	Unavailable	
<i>Eufriesea theresiae</i> (Mocsáry, 1908)	Brazil (AM)	2	Unavailable	
<i>Eufriesea violacea</i> (Blanchard, 1840)	Brazil (MG, SC, SP)	15, 37, 63	Available	63
<i>Eufriesea violascens</i> (Mocsáry, 1898)	Brazil (SP)	36	Unavailable	
<i>Eufriesea sp.</i>	Brazil (MG)	26	—	
<i>Euglossa (Euglossa) amazonica</i> Dressler, 1982	Brazil (CE)	18	Unavailable	
<i>Euglossa (Euglossa) anodorhynchi</i> Nemésio, 2006	Brazil (SP)	30, 64	Unavailable	
<i>Euglossa (Euglossa) avicula</i> Dressler, 1982	Brazil (AC, MG)	3, 65	Unavailable	
<i>Euglossa (Euglossa) cordata</i> (Linnaeus, 1758)	Brazil (BA, MA, MG, RN, SP)	17, 21, 32, 35, 36, 39, 65–67	Available	68
<i>Euglossa (Euglossa) fimbriata</i> Moore, 1968	Brazil (SP)	69	Unavailable	
<i>Euglossa (Euglossa) gairanii</i> Dressler, 1982	Brazil (AM, MA)	2, 17, 67	Unavailable	
<i>Euglossa (Euglossa) hemichlora</i> Cockerell, 1917	Colombia	70	Available	70
<i>Euglossa (Euglossa) melanotricha</i> Moore, 1967	Brazil (MG, SP)	12, 36, 65	Unavailable	
<i>Euglossa (Euglossa) modestior</i> Dressler, 1982	Brazil (AC)	3	Unavailable	
<i>Euglossa (Euglossa) pleosticta</i> Dressler, 1982	Brazil (CE, SP)	15, 18, 36	Unavailable	
<i>Euglossa (Euglossa) townsendi</i> Cockerell, 1904	Brazil (CE, MG, SP)	16, 19, 25, 26, 28, 36, 65, 66	Available	71
<i>Euglossa (Euglossa) truncata</i> Rebêlo & Moore, 1996	Brazil (SP)	36, 66	Unavailable	
<i>Euglossa (Euglossa) variabilis</i> Friese, 1899	Ecuador	59	Unavailable	
<i>Euglossa (Glossura) annectans</i> Dressler, 1982	Brazil (SC, SP)	61, 72, 73	Available	73
<i>Euglossa spp.</i>	Brazil (BA, MA)	35, 74	—	
Tetrapediini				
<i>Tetrapedia amplitarsis</i> Friese, 1899	Brazil (SP)	36	Available	75
<i>Tetrapedia curvitaris</i> Friese, 1899	Brazil (BA, MG, SP)	15, 25, 28, 36, 39, 75	Available	75
<i>Tetrapedia diversipes</i> Klug, 1810	Brazil (BA, CE, MG, PB, PE, RJ, SP)	15, 22, 24, 25, 27, 28, 32, 36, 38, 42, 75, 76	Available	75, 77, 78
<i>Tetrapedia garofaloi</i> Moore, 1999	Brazil (SP)	36, 75	Available	75
<i>Tetrapedia maura</i> Cresson, 1878	Costa Rica	46, 50	Unavailable	
<i>Tetrapedia ornata</i> (Spinola, 1853)	Brazil (AM)	2	Unavailable	
<i>Tetrapedia rugulosa</i> Friese, 1899	Brazil (MG, SP)	25, 28, 36, 75	Unavailable	
<i>Tetrapedia spp.</i>	Argentina (BA); Brazil (AC, MG, PE, SP); Ecuador	3, 12, 14–16, 29, 36, 40, 41, 60	—	
Xylocopini				
<i>Xylocopa (Neoxylocopa) augusti</i> Lepeletier, 1841	Argentina (BA); Brazil (PR)	34, 79	Available	79
<i>Xylocopa (Neoxylocopa) frontalis</i> (Olivier, 1789)	Brazil (BA, MG, SP)	28, 35, 39, 80–82	Available	82, 83
<i>Xylocopa (Neoxylocopa) griseescens</i> Lepeletier, 1841	Brazil (BA, MG, SP)	28, 39, 80–82	Available	82

Taxon	Distribution records (States or provinces from Argentina, Brazil and Mexico are indicated under parenthesis)	References for records	Trap nest description	References for nest descriptions
<i>Xylocopa (Neoxylocopa) suspecta</i> Moure & Camargo, 1988	Brazil (MA, MG)	28, 74, 80	Unavailable	
<i>Xylocopa (Schoenherria) subcyanea</i> Pérez, 1901	Brazil (BA, MG)	23, 35	Unavailable	
<i>Xylocopa (Schoenherria) varians</i> Smith, 1874	Costa Rica	46	Unavailable	
<i>Xylocopa</i> sp.	Brazil (MG)	29	—	
Colletinae				
Colletini				
<i>Colletes rufipes</i> Smith, 1879	Brazil (SP)	25	Unavailable	
<i>Colletes</i> spp.	Brazil (PE, PR)	24, 34	—	
<i>Rhynchocolletes</i> sp.	Brazil (PR)	34	Available	34
Hylaeini				
<i>Hylaeus transversus</i> (Vachal, 1909)	Brazil (SP)	15	Unavailable	
<i>Hylaeus</i> spp.	Brazil (PB, RN, PR, SC, SP); Costa Rica	11, 21, 22, 37, 38, 50, 84	—	
Megachilinae				
Anthidiini				
<i>Anthidium andinum</i> Jörgensen, 1912	Argentina (MZ)	85	Available	85
<i>Anthidium decaspilum</i> Moure, 1957	Argentina (MZ)	85	Available	85
<i>Anthidium rubripes</i> Friese, 1908	Argentina (MZ)	85	Available	85
<i>Anthidium vigintipunctatum</i> Friese, 1908	Argentina (LP, MZ)	41, 85, 86	Available	85
<i>Anthidium</i> spp.	Costa Rica, Ecuador	7, 59	—	
<i>Anthidulum</i> sp.	Brazil (SP)	36	Unavailable	
<i>Anthodioctes claudii</i> Urban, 1999	Brazil (PR)	34	Unavailable	
<i>Anthodioctes lunatus</i> (Smith, 1854)	Brazil (AC, PE)	3, 24	Available	87
<i>Anthodioctes manauara</i> Urban, 1999	Brazil (AM)	88	Unavailable	
<i>Anthodioctes megachilooides</i> Holmberg, 1903	Brazil (MG, SP)	15, 25, 29, 30, 38	Available	89
<i>Anthodioctes moratoi</i> Urban, 1999	Brazil (AM)	2	Available	88
<i>Anthodioctes</i> spp.	Brazil (AC, BA, SP), Mexico (QR)	3, 36, 39, 49	—	
<i>Carloticola paraguayensis</i> (Schrottky, 1908)	Brazil (RJ, SP)	36, 90	Available	90
<i>Ctenanthidium bifasciatum</i> Urban, 1993	Argentina (BA)	91	Available	91
<i>Dicranthidium arenarium</i> (Ducke, 1907)	Brazil (BA, PB, PE, RN)	21, 22, 24, 39	Unavailable	
<i>Dicranthidium luciae</i> Urban, 1993	Brazil (BA)	39	Unavailable	
<i>Dicranthidium seabrai</i> Urban, 2002	Brazil (SP)	92	Unavailable	
<i>Dicranthidium</i> spp.	Brazil (MA, PE)	17, 40	—	
<i>Duckeanthidium thielei</i> Michener, 2002	Costa Rica	46, 50	Available	93
<i>Duckeanthidium</i> sp.	Brazil (AM)	2	Unavailable	
<i>Epanthidium autumnale</i> (Schrottky, 1909)	Brazil (PR, SC)	11	Unavailable	
<i>Epanthidium erythrocephalum</i> (Schrottky, 1902)	Brazil (SP)	36	Unavailable	
<i>Epanthidium maculatum</i> Urban, 1995	Brazil (MG)	26, 28	Unavailable	
<i>Epanthidium nectarinioides</i> (Schrottky, 1902)	Brazil (PR, SC, SP)	11, 34, 36	Unavailable	
<i>Epanthidium tigrinum</i> (Schrottky, 1905)	Brazil (CE, MG, PB, PE, RN, SP)	15, 16, 19, 21, 22, 24, 26, 32, 36, 92, 94	Available	94
<i>Epanthidium</i> spp.	Brazil (BA, MG, SP)	12, 36, 39	—	
<i>Hypanthidium maranhense</i> Urban, 1998	Brazil (MA, PE)	24, 74	Unavailable	
<i>Loyolanthidium</i> sp.	Mexico (QR)	49	Unavailable	
<i>Nananthidium gualanense</i> (Cockerell, 1912)	Costa Rica	50	Unavailable	
<i>Saranthidium marginatum</i> Moure & Urban, 1994	Brazil (SP)	25	Unavailable	
<i>Saranthidium muscifforme</i> (Schrottky, 1902)	Brazil (SP)	36, 92	Unavailable	
Lithurgini				
<i>Microthurgus corumbae</i> (Cockerell, 1901)	Brazil (SP)	95	Unavailable	
<i>Trichothurgus laticeps</i> (Friese, 1906)	Argentina (MZ)	96	Available	96
Megachilini				
<i>Megachile (Acentron)</i> sp.	Argentina (BA)	41	Unavailable	
<i>Megachile (Austromegachile) facialis</i> Vachal, 1909	Brazil (SP, MG)	15, 23, 29	Unavailable	
<i>Megachile (Austromegachile) fiebrigi</i> Schrottky, 1908	Brazil (PR)	13, 34	Unavailable	
<i>Megachile (Austromegachile) orbiculata</i> Mitchell, 1930	Brazil (AC, AM)	2, 3	Unavailable	
<i>Megachile (Austromegachile) sejuncta</i> Cockerell, 1927	Brazil (MA)	74	Unavailable	
<i>Megachile (Austromegachile) susurrans</i> Haliday, 1836	Brazil (PR)	13	Unavailable	
<i>Megachile (Austromegachile) trigonaspis</i> Schrottky, 1913	Brazil (PR)	97	Unavailable	
<i>Megachile (Austromegachile)</i> spp.	Argentina (BA), Brazil (AC, CE, MG, PR)	3, 14, 18, 41, 97	—	
<i>Megachile (Callomegachile) rufipennis</i> (Fabricius, 1793)	Jamaica	1	Available	1
<i>Megachile (Chrysosarus) catamarcensis</i> Schrottky, 1908	Argentina (LP)	41, 86, 98	Available	98
<i>Megachile (Chrysosarus) guaranítica</i> Schrottky, 1908	Brazil (PR, SP)	13, 15, 36	Available	99
<i>Megachile (Chrysosarus) jenseni</i> Friese, 1906	Argentina (BA)	41	Unavailable	
<i>Megachile (Chrysosarus) pseudanthidioides</i> Moure, 1943	Brazil (SC)	100	Available	100
<i>Megachile (Chrysosarus) ruficornis</i> Smith, 1853	Brazil (AC)	3	Unavailable	
<i>Megachile (Chrysosarus)</i> spp.	Argentina (BA), Brazil (AC, BA, CE, MG, PB, PE, PR, RN, SC, SP)	3, 11, 14, 21, 22, 24, 30, 37, 39, 41	—	
<i>Megachile (Dasymegachile)</i> sp.	Brazil (PR)	34	Unavailable	
<i>Megachile (Eutricharaea) concinna</i> Smith, 1879	Argentina (BA), Jamaica	1, 101	Available	1, 101

Taxon	Distribution records (States or provinces from Argentina, Brazil and Mexico are indicated under parenthesis)	References for records	Trap nest description	References for nest descriptions
<i>Megachile (Grafella)</i> sp.	Brazil (SC)	61	—	
<i>Megachile (Melanosarus) brasiliensis</i> Dalla Torre, 1896	Brazil (PR)	34	Unavailable	
<i>Megachile (Melanosarus) nigripennis</i> Spinola, 1841	Brazil (RJ)	102	Available	102
<i>Megachile (Melanosarus)</i> spp.	Brazil (MG, SC)	12, 61	—	
<i>Megachile (Moureapis) benigna</i> Mitchell, 1930	Brazil (MG, RJ)	103, 104	Available	104, 105
<i>Megachile (Moureapis) maculata</i> Smith, 1853	Brazil (MG, PR)	14, 97, 104, 106	Available	104, 106
<i>Megachile (Moureapis) pleuralis</i> Vachal, 1909	Brazil (PR, SC)	11	Unavailable	
<i>Megachile (Moureapis)</i> spp.	Brazil (AC, CE, MG, PR)	3, 14, 19, 34, 97, 107	—	
<i>Megachile (Neochelynia) brethesi</i> Schrottky, 1909	Brazil (MA, MG)	32, 74	Unavailable	
<i>Megachile (Neochelynia) paulista</i> (Schrottky, 1920)	Brazil (AC)	3	Unavailable	
<i>Megachile (Pseudocentron) curvipes</i> Smith, 1853	Brazil (AC, MA, SP)	3, 15, 74	Unavailable	
<i>Megachile (Pseudocentron) gomphrenoides</i> Vachal, 1909	Argentina (BA)	41, 108	Available	108
<i>Megachile (Pseudocentron) inscita</i> Mitchell, 1930	Brazil (BA)	39	Unavailable	
<i>Megachile (Pseudocentron) nudiventris</i> Smith, 1853	Brazil (SC)	61	Unavailable	
<i>Megachile (Pseudocentron)</i> spp.	Argentina (BA); Brazil (AC, MG, PE, RN, SP)	3, 21, 24, 25, 29, 32, 38, 41	—	
<i>Megachile (Pseudomegachile) lanata</i> (Fabricius, 1775)	Jamaica	1	Available	1
<i>Megachile (Ptilosarus) bertonii</i> Schrottky, 1908	Brazil (MG)	29	Unavailable	
<i>Megachile (Ptilosarus) leucostomella</i> Cockerell, 1927	Brazil (AC)	3	Unavailable	
<i>Megachile (Ptilosarus)</i> spp.	Brazil (AC, SP)	3, 30	—	
<i>Megachile (Ptilosaroides) neoxanthoptera</i> Cockerell, 1933	Brazil (SP)	25, 36	Unavailable	
<i>Megachile (Rhyssomegachile)</i> sp.	Brazil (AM)	2	Unavailable	
<i>Megachile (Sayapis) cylindrica</i> Friese, 1906	Brazil (MG, PB, PE)	24, 32	Available	24
<i>Megachile (Sayapis) mendozana</i> Cockerell, 1907	Argentina (BA)	109	Available	109
<i>Megachile (Sayapis) planula</i> Vachal, 1909	Brazil (SP)	15	Unavailable	
<i>Megachile (Sayapis) zaptlana</i> Cresson, 1878	Jamaica, Mexico (QR)	1, 49	Available	1
<i>Megachile (Sayapis)</i> spp.	Brazil (CE, PB, RN)	19, 21, 22, 57	—	
<i>Megachile (Tylomegachile) orba</i> Schrottky, 1913	Brazil (AC)	3	Unavailable	
<i>Megachile</i> spp.	Brazil (AM, BA, MA, MG, PR, SP); Costa Rica; Ecuador	2, 7, 12, 17, 25, 26, 36, 50, 59, 60, 74, 107	—	
Osmiini				
<i>Heriades</i> spp.	Mexico (QR)	49	—	

REFERENCES

- Jayasingh, D.B. & Freeman, B.E. 1980. Trap-nesting solitary aculeates (Insecta: Hymenoptera) in St. Catherine Jamaica. *Caribbean Journal of Science*, 15: 69-78.
- Morato, E.F. 2001. Efeitos da fragmentação florestal sobre vespas e abelhas solitárias na Amazônia Central. II. Estratificação vertical. *Revista Brasileira de Zoologia*, 18: 737-747.
- Morato, E.F. 2004. *Efeitos da sucessão florestal sobre a nidificação de vespas e abelhas solitárias*. (Doctoral Thesis). Belo Horizonte, Universidade Federal de Minas Gerais.
- Carvalho, G.C.A.; Carreira, L.M.M.; Rêgo, M.M.C. & Albuquerque, P.M.C. 2016. Nesting habits of *Centris (Hemisiella) dichrotricha* (Hymenoptera: Apidae) in the Northern Cerrado of Brazil. *Revista de Biología Tropical*, 64: 1041-1056.
- Marroquín, H.A.S. 2017. *Diversidad y variación temporal de abejas del género Centris (Hymenoptera: Apidae: Centridini) en el ecosistema de bosque seco Desierto de la Tatacoa (Neiva-Huila)*. (Master dissertation) Colombia, Universidad Militar Nueva Granada. Bogotá.
- Velez, D.; Vivallo, F. & Silva, D.P. 2017. Nesting biology and potential distribution of an oil-collecting Centridine Bee from South America. *Apidologie*, 48: 181-193.
- Cooperband, M.F.; Wharton, R.A.; Frankie, G.W. & Vinson, S.B. 1999. New host and distribution records for *Leucospis* (Hymenoptera: Leucospidae) associated primarily with nests of *Centris* (Hymenoptera: Anthophoridae) in the dry forests of Costa Rica. *Journal of Hymenoptera Research*, 8: 154-164.
- Vinson, S.B.; Frankie, G. & Cònsoli, R. 2010. Description, Comparison and Identification of Nests of Cavity-Nesting *Centris* Bees (Hymenoptera: Apidae: Centridini) in Guanacaste Province, Costa Rica. *Journal of the Kansas Entomological Society*, 83: 25-46.
- Morato, E.F.; Garcia, M.V.B. & Campos, L.A.D.O. 1999. Biología de *Centris* Fabricius (Hymenoptera, Anthophoridae, Centridini) em matas contínuas e fragmentos na Amazônia Central. *Revista Brasileira de Zoologia*, 16: 1213-1222.
- Mendes, F.N. & Rêgo, M.M.C. 2006. Nidificação de *Centris (Hemisiella) tarsata* Smith (Hymenoptera, Centridini) em ninhos-armadilha no Nordeste do Maranhão, Brasil. *Revista Brasileira de Entomologia*, 51: 382-388.
- Iantas, J.; Gruchowski-Woitowicz, F.C. & Buschini, M.L.T. 2017. Habitat modification and alpha-beta diversity in trap nesting bees and wasps (Hymenoptera: Aculeata) in southern Brazil. *Tropical Zoology*, 30: 83-96.
- Araujo, G.J.; Fagundes, R. & Antonini, Y. 2017. Trap-nesting Hymenoptera and their network with parasites in Recovered Riparian Forests Brazil. *Neotropical Entomology*, 47: 26-36.
- Oliveira, P.S. & Gonçalves, R.B. 2017. Trap-nesting bees and wasps (Hymenoptera, Aculeata) in a Semideciduous Seasonal Forest Fragment, Southern Brazil. *Papéis Avulsos de Zoologia*, 57: 149-156.
- Perillo, L.N.; Neves, F.S.; Antonini, Y. & Martins, R.P. 2017. Compositional changes in bee and wasp communities along Neotropical mountain altitudinal gradient. *PLoS One*, 12: 1-14.
- Rocha-Filho, L.C.; Rabelo, L.S.; Augusto, S.C. & Garófalo, C.A. 2017. Cavity-nesting bees and wasps (Hymenoptera: Aculeata) in a semi-deciduous Atlantic forest fragment immersed in a matrix of agricultural land. *Journal of Insect Conservation*, 21: 727-736.

16. Araújo, P.C.S.; Lourenço, A.P. & Raw, A. 2016. Trap-nesting bees in Montane Grassland (Campo Rupestre) and Cerrado in Brazil: collecting generalist or specialist nesters. *Neotropical Entomology*, 45: 482-489.
17. Gostinski, L.F.; Carvalho, G.C.A.; Rêgo, M.M.C. & Albuquerque, P.M.C. 2016. Species richness and activity pattern of bees (Hymenoptera, Apidae) in the restinga area of Lençóis Maranhenses National Park, Barreirinhas, Maranhão, Brazil. *Revista Brasileira de Entomologia*, 60: 319-327.
18. Brasil, M.O.G. 2015. Aspectos da biologia de nidificação de abelhas solitárias no Maciço de Baturité, Ceará. Doctoral Thesis, Universidade Federal do Ceará.
19. Carvalho, G.C.A. 2014. Comunidade de abelhas solitárias (Hymenoptera: Apidae) que nidificam em ninhos-armadilha em uma área de Cerrado no Parque Estadual do Mirador, Formosa da Serra Negra, Maranhão. Masters Dissertation, Universidade Federal do Maranhão.
20. Gonçalves, L.; Silva, C.I. & Buschini, M.L.T. 2012. Collection of pollen grains by *Centris (Hemisiella) tarsata* Smith (Apidae: Centridini): Is *C. tarsata* an oligolectic or polylectic species? *Zoological Studies*, 51: 195-203.
21. Melo, R.R. & Zanella, S.C.V. 2010. Sazonalidade na nidificação de abelhas e vespas solitárias em ninhos-armadilha em área de Caatinga. *Sitentibus Série Ciências Biológicas*, 10: 258-266.
22. Martins, C.F.; Ferreira, R.P. & Carneiro, L.T. 2012. Influence of the orientation of nest entrance, shading, and substrate on sampling trap-nesting bees and wasps. *Neotropical Entomology*, 41: 105-111.
23. Pires, E.P.; Pompeu, D.C. & Souza-Silva, M. 2012. Nidificação de vespas e abelhas solitárias (Hymenoptera: Aculeata) na Reserva Biológica Boqueirão, Ingaí, Minas Gerais. *Journal of Biosciences*, 28: 302-311.
24. Santos, A.S.S. 2011. Nidificação de abelhas e vespas solitárias e biologia reprodutiva de *Megachile dentipes* Vachal (Hymenoptera, Megachilidae) em ninhos-armadilha. Masters Dissertation, Universidade Federal da Paraíba.
25. Gazola, A.L. & Garófalo, C.A. 2009. Trap-nesting bees (Hymenoptera: Apoidea) in forest fragments of the State of São Paulo, Brazil. *Genetics and Molecular Research*, 8: 607-22.
26. Mesquita, T.M.D.S. 2009. Diversidade de abelhas solitárias (Hymenoptera, Apoidea) que nidificam em ninhos-armadilha em áreas de Cerrado. Masters Dissertation, Universidade Federal de Uberlândia.
27. Menezes, G.B. 2011. Abelhas coletoras de óleos florais na Reserva Biológica União, RJ: Composição e diversidade de espécies, nidificação em ninhos-armadilha e utilização de fontes polínicas. Masters Dissertation, Universidade Estadual do Norte Fluminense Darcy Ribeiro.
28. Carvalho, S.M. 2011. Diversidade de abelhas e vespas solitárias (Hymenoptera, Apoidea) que nidificam em ninhos-armadilha disponibilizados em áreas de Cerrado e em fragmentos próximos de Mata Estacional Semidecidual, MG. Masters Dissertation, Universidade Federal de Uberlândia.
29. Loyola, R.D. & Martins, R.P. 2008. Habitat structure components are effective predictors of trap-nesting Hymenoptera diversity. *Basic and Applied Ecology*, 9: 735-742.
30. Marchi, P. 2008. Biologia de nidificação de abelhas solitárias em áreas de mata atlântica. Doctoral Thesis, Universidade Federal do Paraná.
31. Viana, B.F.; Neves, E.L. & Silva, F.O. 2001. Aspectos da biologia de nidificação de *Euplusia mussitans* (Fabricius) (Hymenoptera, Apidae, Euglossini). *Revista Brasileira de Zoologia*, 18: 1081-1087.
32. Aguiar, A.J.C. & Martins, C.F. 2002. Abelhas e vespas solitárias em ninhos-armadilha na Reserva Biológica Guaribas (Mamanguape, Paraíba, Brasil). *Revista Brasileira de Zoologia*, 19: 101-116.
33. Aguiar, C.M.L. & Garófalo, C.A. 2004. Nesting biology of *Centris (Hemisiella) tarsata* Smith (Hymenoptera, Apidae, Centridini). *Revista Brasileira de Zoologia*, 21: 477-486.
34. Buschini, M.L.T. 2006. Species diversity and community structure in trap-nesting bees in Southern Brazil. *Apidologie*, 37: 58-66.
35. Viana, B.F.; Silva, F.O. & Kleinert, A.M.P. 2001. Diversidade e sazonalidade de abelhas solitárias (Hymenoptera: Apoidea) em dunas litorâneas no nordeste do Brasil. *Neotropical Entomology*, 30: 245-251.
36. Camillo, E.; Garófalo, C.A.; Serrano, J.C. & Muchillo, G. 1995. Diversidade e abundância sazonal de abelhas e vespas solitárias em ninhos-armadilha (Hymenoptera, Apocrita, Aculeata). *Revista Brasileira de Entomologia*, 39: 459-470.
37. Krug, C. & Alves-dos-Santos, I.O. 2008. Uso de diferentes métodos para amostragem da fauna de abelhas (Hymenoptera: Apoidea), um estudo em Floresta Ombrofila Mista em Santa Catarina. *Neotropical Entomology*, 37: 265-278.
38. Alves-dos-Santos, I. 2003. Trap-nesting bees and wasps on the university campus in São Paulo, Southeastern Brazil (Hymenoptera: Aculeata). *Journal of the Kansas Entomological Society*, 76: 328-334.
39. Aguiar, C.M.L.; Garófalo, C.A. & Almeida, G.F. 2005. Trap-nesting bees (Hymenoptera, Apoidea) in areas of dry semideciduous forest and Caatinga, Bahia, Brazil. *Revista Brasileira de Zoologia*, 22: 1030-1038.
40. Oliveira, R. & Schlindwein, C. 2009. Searching for a manageable pollinator for Acerola orchards: the solitary oil-collecting bee *Centris analis* (Hymenoptera: Apidae: Centridini). *Journal of Economic Entomology*, 102: 265-273.
41. Torretta, J.P.; Molina, G.A.R. & Aquino, D.A. 2017. Life cycles and host-parasitoid relationships of five species of *Leucospis* wasps in Argentina (Hymenoptera: Leucospidae). *Journal of Natural History*, 51: 1727-1742.
42. Pina, W.D.C. & Aguiar, C.M.L. 2011. Trap-nesting bees (Hymenoptera: Apidae) in orchards of acerola (*Malpighia emarginata*) in a semiarid region of Brazil. *Sociobiology*, 58: 379-392.
43. Silva, F.O.; Viana, B.F. & Neves, E.L. 2001. Biologia e Arquitetura de Ninhos de *Centris (Hemisiella) tarsata* Smith (Hymenoptera: Apidae: Centridini). *Neotropical Entomology*, 30: 541-545.
44. Buschini, M.L.T. & Wolff, L.L. 2006. Nesting biology of *Centris (Hemisiella) tarsata* Smith in Southern Brazil (Hymenoptera, Apidae, Centridini). *Brazilian Journal of Biology*, 66: 1091-1101.
45. Aguiar, C.M.L.; Garófalo, C.A. & Almeida, G.F. 2006. Biologia de nidificação de *Centris (Hemisiella) trigonoides* Lepeletier (Hymenoptera, Apidae, Centridini). *Revista Brasileira de Zoologia*, 23: 323-330.
46. Thiele, R. 2005. Phenology and nest site preferences of wood-nesting bees in a Neotropical lowland rain forest. *Studies on Neotropical Fauna and Environment*, 40: 39-48.
47. Pereira, M.; Garófalo, C.A.; Camillo, E. & Serrano, J.C. 1999. Nesting biology of *Centris (Hemisiella) vittata* Lepeletier in southeastern Brazil (Hymenoptera, Apidae, Centridini). *Apidologie*, 30: 327-338.

48. Ramos, M.; de Albuquerque, P. & Rêgo, M. 2010. Nesting behavior of *Centris (Hemisiella) vittata* Lepeletier (Hymenoptera: Apidae) in an area of the Cerrado in the northeast of the state of Maranhão, Brazil. *Neotropical Entomology*, 39: 379-383.
49. Roubik, D.W. & Villanueva-Gutiérrez, R. 2009. Invasive Africanized honey bee impact on native solitary bees: A pollen resource and trap nest analysis. *Biological Journal of the Linnean Society*, 98: 152-160.
50. Stangler, E.S.; Hanson, P.E. & Steffan-Dewenter, I. 2015. Interactive effects of habitat fragmentation and microclimate on trap-nesting Hymenoptera and their trophic interactions in small secondary rainforest remnants. *Biodiversity and Conservation*, 24: 563-577.
51. Moure-Oliveira, D.; Rocha-Filho, L.C.; Ferreira-Caliman, M.J. & Garófalo, C.A. 2017. Nesting dynamic and sex allocation of the oil-collecting bee *Centris (Heterocentris) analis* (Fabricius, 1804) (Apidae: Centridini). *Journal of Natural History*, 51: 1-18.
52. Couto, R.M. & Camillo, E. 2007. Influência da temperatura na mortalidade de imaturos de *Centris (Heterocentris) analis* (Hymenoptera, Apidae, Centridini). *Iheringia, Série Zoologia*, 97: 51-55.
53. Dórea, M. da C.; Aguiar, C.M.L.; Figueroa, L.E.R.; Lima e Lima, L.C. & Santos, F.A.R. 2010. Polen residual em ninhos de *Centris analis* (Hymenoptera, Apidae, Centridini) em uma área de vegetação de Caatinga no Brasil. *Oecologia Australis*, 14: 232-237.
54. Jesus, B.M.V. & Garófalo, C.A. 2000. Nesting behaviour of *Centris (Heterocentris) analis* (Fabricius) in southeastern Brazil (Hymenoptera, Apidae, Centridini). *Apidologie*, 31: 503-515.
55. Alonso, J.D.S.; Silva, J.F. & Garófalo, C.A. 2012. The effects of cavity length on nest size, sex ratio and mortality of *Centris (Heterocentris) analis* (Hymenoptera, Apidae, Centridini). *Apidologie*, 43: 436-448.
56. Rocha-Filho, L.C.; Morato, E.F. & Melo, G.A.R. 2009. New host records of *Aglaomelissa duckei* and a compilation of host associations of Ericroidini bees (Hymenoptera: Apidae). *Zoologia, Curitiba*, 26: 299-304.
57. Viana, B.F.; Melo, A.M.C. & Drumond, P.D. 2006. Variação na estrutura do habitat afetando a composição de abelhas e vespas solitárias em remanescentes florestais urbanos de Mata Atlântica no Nordeste do Brasil. *Sitientibus, série Ciências Biológicas*, 6: 282-295.
58. Drummond, P.; Silva, F.O. & Viana, B.F. 2008. Ninhos de *Centris (Heterocentris) terminata* Smith (Hymenoptera: Apidae, Centridini) em fragmentos de Mata Atlântica Secundária, Salvador, BA. *Neotropical Entomology*, 37: 239-246.
59. Tylianakis, J.M.; Klein, A.M.; Lozada, T. & Tscharntke, T. 2006. Spatial scale of observation affects alpha, beta and gamma diversity of cavity-nesting bees and wasps across a tropical land-use gradient. *Journal of Biogeography*, 33: 1295-1304.
60. Vedder, D.; Tylianakis, J.; Tscharntke, T. & Klein, A.M. 2010. Natural enemy diversity reduces temporal variability in wasp but not bee parasitism. *Oecologia*, 162: 755-62.
61. Kamke, R.; Zillikens, A. & Steiner, J. 2011. Species richness and seasonality of bees (Hymenoptera, Apoidea) in a restinga area in Santa Catarina, southern Brazil. *Studies on Neotropical Fauna and Environment*, 46: 35-48.
62. Kamke, R.; Zillikens, A.; Heinle, S. & Steiner, J. 2008. Natural enemies and life cycle of the orchid bee *Eufriesea smaragdina* (Hymenoptera: Apidae) reared from trap nests. *Journal of the Kansas Entomological Society*, 81: 101-109.
63. Perqueti, R.C. & Campos, L.A.O. 1997. Aspectos da biologia de *Euplusia violacea* (Blanchard) (Hymenoptera, Apidae, Euglossini). *Revista Brasileira de Zoologia*, 14: 91-97.
64. Melo, G.A.R.; Faria, L.R.R.; Marchi, P. & Carvalho, C.J.B. 2008. Small orchid bees are not safe: parasitism of two species of *Euglossa* (Hymenoptera: Apidae: Euglossina) by conopid flies (Diptera: Conopidae). *Revista Brasileira de Zoologia*, 25: 573-575.
65. Perqueti, R.C.; Campos, L.A.O.; Coelho, C.D.P.; Abrantes, C.V.M. & Lisboa, L.C.O. 1999. Abelhas Euglossini (Apidae) de áreas de Mata Atlântica: abundância, riqueza e aspectos biológicos. *Revista Brasileira de Zoologia*, 16: 101-118.
66. Freiria, G.A.; Rocha-Filho, L.C.; Del Lama, M.A. & Garófalo, C.A. 2017. Surrounded by foes: Parasitism in *Euglossa* spp. (Apidae, Euglossini) caused by six species of natural enemies. *Journal of Apicultural Research*, 56: 255-261.
67. Silva, O.; Rego, M.M.C.; Albuquerque, P.M.C. & Ramos, M.C. 2009. Abelhas Euglossina (Hymenoptera: Apidae) em área de restinga do nordeste do Maranhão. *Neotropical Entomology*, 38: 186-196.
68. Garófalo, C.A. 1985. Social structure of *Euglossa cordata* nests (Hymenoptera: Apidae: Euglossini). *Entomologia Generalis*, 11: 77-83.
69. Augusto, S.C. & Garófalo, C.A. 2009. Bionomics and sociological aspects of *Euglossa fimbriata* (Apidae, Euglossini). *Genetics and Molecular Research*, 8: 525-538.
70. Parra-H, A. & Nates-Parra, G. 2009. La arquitectura de nidos de *Euglossa (Euglossa) hemichlora* (Hymenoptera: Apidae: Euglossini). *Revista Colombiana de Entomología*, 35: 283-285.
71. Augusto, S.C. & Garófalo, C.A. 2004. Nesting biology and social structure of *Euglossa (Euglossa) townsendi* Cockerell (Hymenoptera, Apidae, Euglossini). *Insectes Sociaux*, 51: 400-409.
72. Cortopassi-Laurino, M.; Zillikens, A. & Steiner, J. 2009. Pollen sources of the orchid bee *Euglossa annectans* Dressler 1982 (Hymenoptera: Apidae, Euglossini) analyzed from larval provisions. *Genetics and Molecular Research*, 8: 546-556.
73. Garófalo, C.A.; Camillo, E.; Augusto, S.C.; Jesus, B.M.V. & Serrano, J.C. 1998. Nest structure and communal nesting in *Euglossa (Glossura) annectans* Dressler (Hymenoptera, Apidae, Euglossini). *Revista Brasileira de Zoologia*, 15: 589-596.
74. Matos, M.C.B.; Silva, S.S. & Teodoro, A.V. 2016. Seasonal population abundance of the assembly of solitary wasps and bees (Hymenoptera) according to land-use in Maranhão state, Brazil. *Revista Brasileira de Entomologia*, 60: 171-176.
75. Camillo, E. 2005. Nesting biology of four *Tetrapedia* species in trap nests (Hymenoptera: Apidae: Tetrapediini). *Revista de Biologia Tropical*, 53: 175-186.
76. Cavalcante, A.M. 2016. Nicho trófico e aspectos bionômicos da abelha solitária *Tetrapedia diversipes* Klug (Hymenoptera, Apidae, Tetrapediini). Masters Dissertation, Universidade Federal do Ceará.
77. Alves-dos-Santos, I.; Melo, G.A.R. & Rozen, J.G. 2002. Biology and immature stages of the bee tribe Tetrapediini (Hymenoptera: Apidae). *American Museum Novitates*, 1-45.
78. Menezes, G.B.; Gonçalves-Esteves, V.; Bastos, E.M.A.F.; Augusto, S.C. & Gaglianone, M.C. 2012. Nesting and use of pollen resources by *Tetrapedia diversipes* Klug (Apidae) in Atlantic Forest areas (Rio de Janeiro, Brazil) in different stages of regeneration. *Revista Brasileira de Entomologia*, 56: 86-94.
79. Lucia, M.; Telleria, M.C.; Ramello, P.J. & Abrahamovich, A.H. 2017. Nesting ecology and floral resource of *Xylocopa augusti* Lepeletier de Saint Fargeau (Hymenoptera, Apidae) in Argentina. *Agricultural and Forest Entomology*, 19: 281-293.

80. Alves-Chaves, T.M. 2009. Nidificação de *Xylocopa* sp. (Apidae, Xylocopini) em ninhos-armadilha em áreas de cerrado do Triângulo Mineiro. Masters Dissertation, Universidade Federal de Uberlândia.
81. Junqueira, C.N.; Hogendoorn, K. & Augusto, S.C. 2012. The use of trap-nests to manage carpenter bees (Hymenoptera: Apidae: Xylocopini), pollinators of passion fruit (Passifloraceae: *Passiflora edulis* f. *flavicarpa*). *Annals of the Entomological Society of America*, 105: 884-889.
82. Pereira, M. & Garófalo, C.A. 2010. Biologia da nidificação de *Xylocopa frontalis* e *Xylocopa griseescens* (Hymenoptera, Apidae, Xylocopini) em ninhos-armadilha. *Oecologia Australis*, 14: 193-209.
83. Marchi, P. & Melo, G.A.R. 2010. Biologia de nidificação de *Xylocopa* (*Neoxylocopa*) *frontalis* (Olivier) (Hymenoptera, Apidae, Xylocopini) em Morretes, Paraná. *Oecologia Australis*, 14: 210-231.
84. Macedo, A.C.C.; Cordeiro, G.D. & Alves-dos-Santos, I. 2012. Entering behavior of *Gasteruption brachychaetum* Schrottky (Hymenoptera, Gasteruptionidae) into a nest of *Hylaeus Fabricius* (Hymenoptera, Colletidae). *Revista Brasileira de Entomologia*, 56: 325-328.
85. Vitale, N.; Gonzalez, V.H. & Vázquez, D.P. 2017. Nesting ecology of sympatric species of wool carder bees (Hymenoptera: Megachilidae: *Anthidium*) in South America. *Journal of Apicultural Research*, 56: 497-509.
86. Torretta, J.P. 2015. Host-parasite relationships and life cycles of cuckoo wasps in agro-ecosystems in Argentina (Hymenoptera: Chrysididae: Chrysidini). *Journal of Natural History*, 49: 1641-1651.
87. Camarotti-de-Lima, M.F. & Martins, C.F. 2005. Biologia de nidificação e aspectos ecológicos de *Anthodioctes lunatus* (Smith) (Hymenoptera: Megachilidae, Anthidiini) em área de tabuleiro nordestino, PB. *Neotropical Entomology*, 34: 375-380.
88. Morato, E.F. 2001. Biologia e ecologia de *Anthodioctes moratoi* Urban (Hymenoptera, Megachilidae, Anthidiini) em matas contínuas e fragmentos na Amazônia Central, Brasil Anthidiinae. *Revista Brasileira de Entomologia*, 18: 729-736.
89. Alves-dos-Santos, I. 2004. Biologia de nidificação de *Anthodioctes megachiloides* Holmberg (Anthidiini, Megachilidae, Apoidea). *Revista Brasileira de Zoologia*, 21: 739-744.
90. Mello, B.N.S. 2014. *Nicho de abelhas Megachilidae (Hymenoptera, Apoidea) em floresta ombrófila na Mata Atlântica: dinâmica temporal, relações tróficas e uso de recursos de nidificação*. Masters Dissertation, Universidade Estadual do Norte Fluminense Darcy Ribeiro.
91. Alvarez, L.J.; Lucia, M.; Durante, S.P.; Passarelli, L.M. & Gonzales, V.H. 2015. Description of the male, new geographical record, and biological notes on the small resin bee *Hypanthidioides* (Megachilidae). *Pan-Pacific Entomologist*, 91: 184-188.
92. Cordeiro, G.D.; Nascimento, A.L.O.; Alves-dos-Santos, I. & Garófalo, C.A. 2015. The cleptoparasitic bee *Austrostelis iheringi* (Schrottky, 1910) (Anthidiini, Megachilidae): New geographical records and potential hosts. *Journal of the Kansas Entomological Society*, 88: 453-456.
93. Thiele, R. 2002. Nesting biology and seasonality of *Duckeanthidium thielei* Michener (Hymenoptera: Megachilidae), an oligolectic Rainforest bee. *Journal of the Kansas Entomological Society*, 75: 274-282.
94. Gomes, A.M.S. 2016. *Bionomia e compartamento de nidificação da abelha Epanthidium tigrinum (Hymenoptera-Megachilidae) em ninhos-armadilha*. (Master Dissertation). Universidade Federal do Ceará, Fortaleza, Brazil.
95. Garófalo, C.A.; Camillo, E.; Campos, M.J.O. & Serrano, J.C. 1992. Nest re-use and communal nesting in *Microthurga corumbae* (Hymenoptera, Megachilidae), with special reference to nest defense. *Insectes Sociaux*, 39: 301-311.
96. Vitale, N. & Vázquez, D.P. 2017. Ecology and nesting biology of the wood-boring bee *Trichothurgus laticeps* (Hymenoptera: Megachilidae) in a Monte desert reserve in mid-western Argentina. *Apidologie*, 48: 31-40.
97. Woiski, T.D. 2009. *Estrutura da comunidade de vespas e abelhas solitárias em um fragmento urbano de floresta ombrófila mista*. Masters Dissertation, Universidade Federal do Paraná.
98. Torretta, J.P.; Durante, S.P. & Basilio, A.M. 2014. Nesting ecology of *Megachile (Chrysosarus) catamarcensis* Schrottky (Hymenoptera: Megachilidae), a *Prosopis*-specialist bee. *Journal of Apicultural Research*, 53: 590-598.
99. Rocha-Filho, L.C. & Garófalo, C.A. 2016. Nesting biology of *Megachile (Chrysosarus) guaranítica* and high mortality caused by its cleptoparasite *Coelioxys bertonii* (Hymenoptera: Megachilidae) in Brazil. *Austral Entomology*, 55: 25-31.
100. Zillikens, A. & Steiner, J. 2004. Nest architecture, life cycle and cleptoparasite of the neotropical leaf-cutting bee *Megachile (Chrysosarus) pseudanthidioides* Moure (Hymenoptera: Megachilidae). *Journal of the Kansas Entomological Society*, 77: 193-202.
101. Alvarez, L.J.; Lucia, M.; Aquino, D.A.; Ramello, P.J. & Abrahamovich, A.H. 2016. Nesting biology and associated insect enemies of the exotic leaf cutter bee *Megachile (Eutricharaea) concinna* (Hymenoptera: Megachilidae) in Argentina. *Journal of Apicultural Research*, 8839: 1-9.
102. Marques, M.F. & Gaglianone, M.C. 2013. Biologia de nidificação e variação altitudinal na abundância de *Megachile (Melanosarus) nigripennis* spinola (Hymenoptera, Megachilidae) em um inselbergue na mata Atlântica, Rio de Janeiro. *Bioscience Journal*, 29: 198-208.
103. Teixeira, F.M. 2011. *Aculeata (Insecta, Hymenoptera) em ninhos-armadilha em diferentes tipos fitofisionômicos de Mata Atlântica no estado do Rio de Janeiro*. Doctoral Thesis, Universidade Estadual do Norte Fluminense Darcy Ribeiro.
104. Cardoso, C.F. & Silveira, F.A. 2012. Nesting biology of two species of *Megachile (Moureapis)* (Hymenoptera: Megachilidae) in a semideciduous forest reserve in southeastern Brazil. *Apidologie*, 43: 71-81.
105. Teixeira, F.M.; Schwartz, T.A.C. & Gaglianone, M.C. 2011. Biologia da nidificação de *Megachile (Moureapis) benigna* Mitchell. *EntomoBrasilis*, 4: 92-99.
106. Sabino, W.O. & Antonini, Y. 2017. Nest architecture, life cycle, and natural enemies of the neotropical leafcutting bee *Megachile (Moureapis) maculata* (Hymenoptera: Megachilidae) in a montane forest. *Apidologie*, 48: 450-460.
107. Buggenhagen, T.C. 2016. *Aspectos ecológicos e biológicos de vespas e abelhas (Insecta, Hymenoptera) que nidificam em cavidades preexistentes*. Masters Dissertation, Universidade Estadual do Centro-Oeste.
108. Torretta, J.P.; Durante, S.P.; Colombo, M.G. & Basilio, A.M. 2012. Nesting biology of the leafcutting bee *Megachile (Pseudocentron) gomphrenoides* (Hymenoptera: Megachilidae) in an agro-ecosystem. *Apidologie*, 43: 624-633.
109. Torretta, J.P. & Durante, S.P. 2011. Nesting ecology of *Megachile (Sayapis) mendozana* Cockerell and its synonymy with *Megachile (Sayapis) santiaguensis* Durante (Hymenoptera: Megachilidae). *Zootaxa*, 3008: 61-68.