

## PRENYLATED XANTHONES FROM *Rheedia acuminata*

Maria das G. B. ZOGHBI<sup>1</sup>, Patrícia D. de D. B. LIMA, Maria do P. Socorro C. da CUNHA<sup>1</sup>, Angelo da C. PINTO<sup>2</sup>

**ABSTRACT** — Pyranojacareubin; 1,5-dihydroxy-6',6'-dimethyl-2H-pyran(2',3':6,7) -6'',6''-dimethyl-2H,4H-pyran(2'',3'':2,3)xanthone and a new xanthone 1,6-dihydroxy-5-methoxy-6',6'-dimethyl-2H-pyran(2',3':3,2)-7-(3,3-dimethylprop-2-enyl)xanthone were isolated from the ether extract of the root bark of *Rheedia acuminata* together with friedelin and friedelanol.

**Key words:** Guttiferae, *Rheedia*, Xanthones.

Xantonas Preniladas de *Rheedia acuminata* (GUTTIFERAE)

**RESUMO** — Piranojacareubina; 1,5-diidroxi-6',6'-dimetil-2H-pirano(2',3':6,7)-6'',6''-dimetil-2H,4H(2'',3'':2,3)xantona e uma xantona inédita 1,6-diidroxi-5-metoxi-6',6'-dimetil-2H-pirano(2',3':3,2)-7-(3,3-dimetilprop-2-enil) xantona foram isoladas do extrato etéreo da casca da raiz de *Rheedia acuminata* além de friedelina e friedelanol.

**Palavras-chave:** Guttiferae, *Rheedia*, Xantonas.

### INTRODUCTION

The family Guttiferae numbers over 1000 species, which occur widely in temperate regions. Xanthones or the related benzophenones have been found in all their major and several minor genera (BENNETT & LEE, 1989). Prenylated xanthones are widely distributed in the Guttiferae, and the genus *Rheedis* has been shown to be rich with them. They have been isolated from the *R. benthamiana* (DELLE MONACHE *et al.*, 1981), *R. gardneriana* (DELLE MONACHE *et al.*, 1983) and *R. brasiliensis* (DELLE MONACHE *et al.*, 1984); we now report the isolation of three prenylated xanthones from the *R. acuminata*, which are present in the ether extract of the roots. Two xanthones were iden-

tified as pyranojacareubin (1) and 1,5-dihydroxy-6',6'-dimethyl-2H-pyran(2',3':6,7)-6'',6''-dimethyl-2H,4H-pyran(2'',3'':2,3)- xanthone (2), while the third one, (3), was novel and was named *acuminatine*. The triterpenes friedelin and friedelanol also were isolated.

### IDENTIFICATION OF CONSTITUENTS

Compound 1, yellow needles (hexane-acetone), mp 261-263°, [lit. 259.5-260.5° (Et<sub>2</sub>O)], C<sub>23</sub>H<sub>20</sub>O<sub>6</sub> (M<sup>+</sup> at m/z 392) two 2,2-dimethyl-2H-pyran rings and two separated aromatic protons (<sup>1</sup>H NMR evidence) exhibited IR and MS spectra data, which were in agreement with those reported in the literature for pyranojacareubin

<sup>1</sup> Instituto Nacional de Pesquisas da Amazônia, Coordenação de Pesquisas em Produtos Naturais, Caixa Postal 478, 69011-970 - Manaus, Amazonas, Brasil.

<sup>2</sup> Instituto de Química, Universidade Federal do Rio de Janeiro, Caixa Postal 1573, 20001-970, Rio de Janeiro, RJ, Brasil.

(DELLEMONACHE *et al.*, 1984). Hydrogenation of 1 furnished 4 (mp 234-235°).

Compound 2, yellow crystals (hexane-acetone), mp 212-215° [lit. 202-204°], C<sub>23</sub>H<sub>22</sub>O<sub>6</sub> at m/z 394), was isolated in a mixture with pyranojacareubin. The <sup>1</sup>H NMR spectrum showed beside signals of the pyranojacareubin system, two triplets (δ 2.90, 2H, J = 7.0 Hz and 1.91, 2H, J = 7.0 Hz) that were assigned to H-4' and H-5', respectively (DELLEMONACHE *et al.*, 1984). The <sup>13</sup>C NMR chemical shifts are presented in Table 1, along with reported shifts for the

closely related geronxanthone B (CHANG *et al.*, 1989). The chemical shifts were assigned on the basis of the PND and DEPT spectra. The presence of the two carbonyl groups (δ 180.4 and 180.1) and two methylenic carbons (δ 32.5 and 21.7) and the signals peaks at 394 (M<sup>+</sup>), 379 (M<sup>+</sup>-Me) and 323 (M<sup>+</sup>-Me-C<sub>4</sub>H<sub>8</sub>) in the mass spectrum permitted identification of the 1,5-dihydroxy-6',6'-dimethyl-2H-pyran(2',3':6,7)-6'', 6''-dimethyl-2H, 4H(2'',3'':2,3)xanthone.

Compound 3, yellow crystals (hexane-Et<sub>2</sub>O, mp 152-154°, C<sub>24</sub>H<sub>24</sub>O<sub>6</sub> (M<sup>+</sup> at m/z 408) showed characteristic

**Table 1.** <sup>13</sup>C NMR chemical shifts of 1,2 and 3, in CDCl<sub>3</sub>

	1	2	ref.6	3	ref.8
C-1	157.6	157.6	163.1	157.8	157.8
C-2	104.4	113.4	113.5	104.7	104.4
C-3	160.3	156.7	161.8	160.3	159.8
C-4	95.2	95.2	95.7	95.0	94.0
C-4a	156.7#	157.0#	155.8	156.6	156.1
C-4b	144.7#	142.8#	144.7	147.8	155.6
C-5	132.2	132.2	131.7	133.9	101.6
C-6	145.0#	146.4#	144.9	152.6	154.5
C-7	117.7#	118.1#	117.6	125.6	142.7
C-8	121.2	121.2	121.4	120.5	136.9
C-8a	114.5	114.5	113.6	114.0	112.1
C-8b	103.1	103.1	103.8	103.1	103.6
C-9	180.4#	180.1#	180.5	180.1	181.8
C-4'	127.3	32.5	—	127.4	126.9
C-5'	116.1	21.7	—	115.5	115.6
C-1''	—	—	—	25.8	26.5
C-2''	—	—	—	121.0	123.1
C-3''	—	—	—	133.2	131.8
C-4''	127.5	127.5	114.5	17.8	18.1
C-5''	115.4#	113.8#	130.8	28.1	25.6
C-6'	78.8#	78.0#	—	78.2	77.8
C-6''	78.1	77.1#	78.6	—	—
6'-Me	28.4	26.9#	—	28.4	28.3
6''-Me	28.3	28.2	28.4	—	—
C-3'''	—	—	40.9	—	—
C-3a'''	—	—	26.9	—	—
C-4'''	—	—	149.6	—	—
C-5'''	—	—	113.7	—	—
OCH <sub>3</sub>	—	—	—	61.8	61.8

<sup>#</sup> Assigned by comparison with the cyclo derivative of rheediaxanthone B (DELLEMONACHE *et al.*, 1981)

# Assigned may be interchanged

UV and IR spectra of a xanthone. The chelated hydroxyl and C-6 hydroxyl, which were indicated by typical UV spectrum, underwent modifications with additives (MESQUITA *et al.*, 1968). The presence of the free hydroxyl was confirmed by methylation with diazomethane ( $M^+$  at  $m/z$  422). The  $\alpha,\alpha$ -dimethylallyl and 2,2-dimethylchromene groups were characterized by  $^1H$  NMR data. The signal at  $\delta$  13.50 confirmed the chelated hydroxyl. The localization of the chromene group at C-2 and C-3 and C-1 hydroxyl was established by comparison of their  $^1H$  and  $^{13}C$  NMR data (Table 1) with reported chemical shifts for the closely related compound (SEN *et al.*, 1980). The signal of the  $OCH_3$  group in the  $^{13}C$  NMR at  $\delta$  61.8 indicates that it is bonded to C-5, between two *ortho* groups (CHAUDHURI *et al.*, 1978). The localization of the  $\alpha,\alpha$ -dimethylallyl at C-7 was confirmed by the long range ( $J^3$ ) coupling, and NOE experiments.

## EXPERIMENTAL

### GENERAL EXPERIMENTAL PROCEDURE

All melting points were uncorrected. UV spectra were recorded on a Perkin\_elmer 402 and IR spectra on a Perkin-Elmer 298 spectrophotometer,  $^{13}C$  NMR spectra on a Bruker spectrometer operating at 50.0 MHz and  $^1H$  NMR spectra on a Bruker spectrometer operating at 100 and 200 MHz.  $^{13}C$  NMR and  $^1H$  NMR spectra were measured with tetramethylsilane (TMS) as

internal reference. Mass spectra were recorded on a HP-5987A instrument at 70 eV.

### PLANT MATERIAL

The root bark of *R. acuminata* was collected in Araguacema, Goiás State, Brazil, and identified by the botanist Dr. William A. Rodrigues, INPA, Manaus, Brasil.

### EXTRACTION AND ISOLATION OF CONSTITUENTS

The root bark (3.05 kg) was reduced to saw dust and extracted at room temperature with petrol ether. The solvent was evaporated giving 94.2g of the extract. Part of it (10.0g) was chromatographed on silica giving the following compounds: 1,2,3-friedelin and friedelanol, eluted with hexane/acetone 4%; 1 (13.0mg) was recrystallized from hexane-acetone (1:1); 2(8.0mg) was purified by CCCP; 3 (60.8mg) was recrystallized from hexane-diethyl ether (1:1).

### SPECTROSCOPY DATA OF CONSTITUENTS

1,5-dihydroxy-6',6'-dimethyl-2H-pyran(2',3':3,2)-6",6"-dimethyl-2H-pyran(2",3":6,7)xanthone (pyranojacareubin, 1),  $C_{23}H_{20}O_6$ , mp 261-263° (yellow needles). IR KBr max (cm-1): 3480, 2969, 1638, 1605, 1467, 1374, 1199, 1157.  $^1H$  NMR ( $CDCl_3$ :  $\delta$  13.80 (1H, 1-OH), 7.48 (1H, s, H-8), 6.73 (1H, d,  $J=10.0$  Hz, H-4'), 6.44 (1H, s, H-4), 6.45 (1H, d,  $J=10.0$  Hz, H-4"), 5.74 (1H, d,  $J=10.0$  Hz, H-5"), 5.60 (1H, d,  $J=10.0$  Hz, H-5'), 1.55 + 1.45 (6H + 6H, s, 4 x Me). MS  $m/z$  (rel. int.): 392 [M]<sup>+</sup> (18), 377

[M-Me]<sup>+</sup> (100), 323 (14), 267 (1), 181 (M-Me-Me)<sup>2+</sup> (24).

1,5-dihydroxy-6',6'-dimethyl-2H-pyran(2',3':6,7)-6'',6''-dimethyl-2H,4H-pyran(2'',3'':3,2) xanthone (dihydropyranojacareubin, 2) C<sub>23</sub>H<sub>22</sub>O<sub>6</sub>, mp 212-215° (yellow needles). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 13.8 (1H, s, 1-OH), 7.48 (1H, s, H-8), 6.45 (1H, d, J=10.0 Hz, H-4''), 6.44 (1H, s, H-4), 5.74 (1H, d, J=10.0 Hz, H-5''), 2.90 (2H, t, J=7.0 Hz, H-4'), 1.91 (2H, t, J=7.0 Hz, H-5'), 1.55 + 1.45 (6H + 6H, s, 4 x Me). MS m/z (rel. int.): 394 [M]<sup>+</sup> (12), 379 (44), 323 (15)

Hydrogenation of 2: Compound 2 (62.0 mg) was hydrogenated by the usual method giving as a pale yellow needles, mp 234-235° (methanol-hexane). IR max (KBr, cm<sup>-1</sup>): 3420, 2920, 1630, 1600, 1580, 1440, 1150, 870, 820. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.45 (1H, s, H-8), 6.34 (1H, s, H-4), 2.84 (2H, t, J=6.2 Hz, H-4'), 2.65 (2H, t, J=6.6 Hz, H-4''), 1.83 (2H, t, J=6.2 Hz, H-5'), 1.78 (2H, t, J=6.6 Hz, H-5''), 1.37 (3H, s, Me), 1.31 (3H, s, Me).

1,6-dihydroxy-5-methoxy-6',6'-dimethyl-2H-pyran(2',3':3,2)-7-(3,3-dimethylprop-2-enyl) xanthone (acuminatine, 3) C<sub>24</sub>H<sub>24</sub>O<sub>6</sub>, mp 152-154°. IR max (KBr, cm<sup>-1</sup>): 3350, 2930, 2890, 2840, 1650, 1580, 1460, 1350, 1190, 1150, 810, 770. UV ε max (EtOH, nm) (log ε): 244(4.79), 279 (4.83), 289 (4.83), 334, (4.60); ε max (AlCl<sub>3</sub>, nm) (after 30 min.): 243, 289, 358; ε max (HCl, nm): 246, 296, 358; ε max (NaOH, nm): 247, 280, 316 sh, 386; ε max (HCl, nm): 243, 280, 334; ε max (NaOAc, nm): 244, 279, 380; ε max

(H<sub>3</sub>BO<sub>3</sub>, nm): 244, 289, 334. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 13.50 (1H, s, 1-OH), 7.73 (1H, s, H-8), 6.72 (1H, d, J = 10.0 Hz, H-4'), 6.58 (1H, s, 6-OH), 6.37 (1H, s, H-4), 5.59 (1H, d, J=10.0 Hz, H-5'), 5.35 (1H, t, J=7.0 Hz, H-2''), 4.10 (3H, s, OMe), 3.40 (2H, d, J=7.0 Hz, H-1''), 1.76 (3H, s, Me), 1.74 (3H, s, Me), 1.50 (6H, s, 2 x Me). MS m/z (rel. int.): 408 [M]<sup>+</sup> (28), 393 [M-Me]<sup>+</sup> (100), 377 (5), 361 (5), 355 (3), 335 (10), 323 (5), 189 (6), 182 (2), 174 (3), 169 (5), 38 (69).

Mono-O-methyl ether of 3. Compound 3 (14.0 mg) was suspended in Et<sub>2</sub>O, treated with excess CH<sub>2</sub>N<sub>2</sub> and left overnight. The solvent was evaporated and the product was recrystallized from hexane-diethyl ether (14.4 mg) mp 135-138°. IR max (KBr, cm<sup>-1</sup>): 2980, 2920, 1650, 1600, 1570, 1420, 1160, 830. MS m/z (rel. int.): 422 [M]<sup>+</sup> (26), 407 [M-Me]<sup>+</sup> (100), 393 (13), 377 (3), 361 (2), 349 (3), 335 (2), 323 (7), 196 (9).

## RESULTS AND DISCUSSION

From the ether extract of the root bark of *Rheedia acuminata* were isolated, by chromatographic techniques pyranojacareubin;

1,5-dihydroxy-6',6'-dimethyl-2H-pyran(2',3':6,7)-6'',6''-dimethyl-2H,4H-pyran(2'',3'':2,3)xanthone, a new xanthone 1,6-dihydroxy-5-methoxy-6',6'-dimethyl-2H-pyran(2',3':3,2)-7-(3,3-dimethylprop-2-enyl) xanthone together with friedelin and friedelanol. The identification or structural elucidation of the compounds were based on spectroscopic techniques.

## Literature Cited

- BENNETT, G. J.; LEE, H. H. 1989. Xanthenes from Guttiferae. *Phytochemistry*, 28(4), 967.
- CHANG, C. H.; LIN, C. C.; HATTORI, M.; NAMBA, T. 1989. Four prenylated xanthenes from *Cidrania cochinchinensis*. *Phytochemistry*, 28(2):595.
- CHAUDHURI, R. K.; ZYMALKOWSKI, F.; FRHAM, A. W. 1978. <sup>13</sup>C NMR-Spectroscopy of Polymethoxyxanthenes. *Tetrahedron*, 34:1837.
- DELLE MONACHE, F.; BOTTA, B.; NICOLETTI, M.; BARROS COELHO, J. S. de; ANDRADE LYRA, J. D. de 1981. Three new xanthenes and macluraxanthenes from *Rheedia benthamiana* Pl. Triana (Guttiferae). *J. Chem. Soc. Perkin Trans.*, 1, 484.
- DELLE MONACHE, G.; DELLE MONACHE, F.; MARINI BETTOLO, G. B.; ALVES DE LIMA, R. de 1983. Chemical investigation of the genus *Rheedia*. I. Prenylated xanthenes from *Rheedia gardneriana*. *J. Nat. Prod.*, 46, 655.
- DELLE MONACHE, G.; BOTTA, B.; MELLO, J. F., GOELHO, J. S. de B.; MENICHINI, F. 1984. Chemical investigation of the genus *Rheedia*. IV. Three new xanthenes from *Rheedia brasiliensis*. *J. Nat. Prod.*, 47(4): 620.
- DELLE MONACHE, G.; DELLE MONACHE, F.; WATERMAN, P. G.; CRICHTON, E. G.; LIMA, R. A. de 1984. Minor xanthenes from *Rheedia gardneriana*. *Phytochemistry*, 23(8):1757.
- MESQUITA, A. A. L.; CORREA, D. e B.; GOTTLIEB, O. R.; MAGALHÃES, M. T. 1968. Métodos para investigação estrutural de xantonas. II. Espectroscopia no ultra-violeta e no visível. *An. Acad. Brasil. Ciênc.*, 40(2):167.
- SEN, A. K.; SARKAR, K. K.; MUZUMDER, P. C.; UUSVUORI, N. B. R.; HASE, T. A. 1980. A xanthone from *Garcinia mangostana*. *Phytochemistry*, 19:2223.

Aceito para publicação em 30/7/93