

Differential Diagnosis between Bartholin Cyst and Vulvar Leiomyoma: Case Report

Diagnóstico diferencial entre cisto de Bartholin e leiomioma vulvar: relato de caso

Kelly Alessandra da Silva Tavares¹ Thomas Moscovitz¹ Marcos Tcherniakovsky¹
 Luciano de Melo Pompei¹ César Eduardo Fernandes¹

¹ Department of Gynecology and Obstetrics, Faculdade de Medicina do ABC, Santo André, São Paulo, Brazil

Address for correspondence Kelly Alessandra da Silva Tavares, MD, Rua dos Democratas, 277, apartamento 41, 04305-000, São Paulo, Brazil (e-mail: kellyalessandradasilva@yahoo.com.br).

Rev Bras Ginecol Obstet 2017;39:433–435.

Abstract

Keywords

- ▶ vulvar neoplasms
- ▶ Bartholin gland
- ▶ leiomyoma
- ▶ histology
- ▶ case report

Genital leiomyomas are rare tumors that can often be misdiagnosed as Bartholin cyst. We report a case of a 32-year-old patient who had a cystic nodulation in the left labium majus that was suggestive of Bartholin cyst. A resection surgery was performed, and the definitive histopathology diagnosis was vulvar leiomyoma. The macroscopic features of cystic lesions difficult the differential diagnosis between leiomyoma and Bartholin cyst; therefore, a histopathologic examination is often recommended.

Resumo

Palavras-Chave

- ▶ neoplasias vulvares
- ▶ glândula de Bartholin
- ▶ leiomioma
- ▶ histologia
- ▶ relato de caso

Os leiomiomas genitais são tumores raros, e frequentemente são diagnosticados como cisto de Bartholin. Relatamos o caso de uma paciente de 32 anos em que ocorreu nodulação cística no grande lábio esquerdo sugestiva de cisto de Bartholin. Uma cirurgia de ressecção foi realizada, e o diagnóstico histopatológico definitivo foi leiomioma vulvar. As características macroscópicas das lesões císticas dificultam o diagnóstico diferencial entre o leiomioma e o cisto de Bartholin, de modo que o exame histopatológico é frequentemente recomendado.

Introduction

Uterine leiomyomas are benign monoclonal tumors, and the most frequent site for their occurrence is the muscle cells of the myometrium. However, they develop in any site where smooth muscle cells are present,¹ such as the vulva, the vagina, the ovaries, the urinary bladder, the urethra, the round ligaments, the uterosacral ligaments, the inguinal canal and the retroperitoneum.^{2,3}

Genital leiomyomas are rare tumors.⁴ Less than 160 cases of vulvar leiomyoma have been reported in the literature, and this condition can often be misdiagnosed as Bartholin cyst.^{4–6}

Some features that support the diagnosis of Bartholin cyst are everted labia minora and cystic consistency of the swelling. However, finding inverted labia minora and firm consistency of the swelling suggests vulvar leiomyoma. A histopathologic examination is often recommended for the final diagnosis.⁷

Case Report

Patient Information

A 32-year-old white woman presented to the gynecological outpatient service at our institution complaining of vaginal pain and a vulvar nodulation that had developed six months

received
 October 29, 2016
 accepted
 May 5, 2017

DOI <https://doi.org/10.1055/s-0037-1604178>.
 ISSN 0100-7203.

Copyright © 2017 by Thieme Revinter Publicações Ltda, Rio de Janeiro, Brazil

License terms



before. She stated feeling no abdominal pain and having no fever and urinary and intestinal disorders. She reported tubal ligation and prolonged menstrual bleeding.

Physical Examination

Her general physical and systemic examinations were unremarkable. Upon local examination, a 5 × 5 cm cystic lesion in the lower part of left labium majus was observed. It was clinically diagnosed as Bartholin cyst.

Diagnostic Assessment

Baseline investigations were performed with a transvaginal ultrasound that evidenced multiple intramural uterine leiomyomas, with the largest measuring 28 mm. The uterine volume was 239 cc. The vulvar region was not assessed with ultrasound because the first hypothesis was Bartholin cyst.

Interventions

Nine months after the initial consultation, the patient came back for surgery, and we observed a growth in the tumor during this period; we found a fibroelastic mass of ~ 10 cm. The tissue appeared to be a leiomyoma, which made us question the initial hypothesis of Bartholin cyst. A specimen was sent for a histopathologic examination.

Follow-Up and Outcomes

The patient had an uneventful postoperative recovery, and was discharged on the first postoperative day. A gross examination of the excised tumor of the left labium majus showed: a piece with irregular surface, mousey and firm tissue, measuring 8 × 6 × 3 cm. Microscopy revealed a mature tumor of mesenchymal nature characterized by proliferation of spindle cells similar to smooth muscle cells. There were no indications of malignancy. The final diagnosis was leiomyoma.

The patient was called for follow-up after two weeks and was asymptomatic; one year later, no recurrence was observed.

Discussion

Uterine leiomyomas are very common benign tumors; however, their occurrence in the vulva is rare. Less than 160 cases of vulvar leiomyoma have been reported in the literature.⁴⁻⁶

The mean age when these tumors tend to occur varies from 13 to 71 years, and they usually occur as solitary lesions, with an average tumor size of 0.5 to 15 cm.⁸ Corroborating with the cases described in the literature, we report a 8-cm single vulvar lesion in a 32-year-old woman who complained of pain in the vulvar region 6 months prior to the consultation. Nevertheless, vulvar leiomyomas are usually asymptomatic, but sometimes can cause swelling and local discomfort.

Ultrasound is the most reliable and widely used diagnostic tool for uterine and extra uterine leiomyomas, and magnetic resonance imaging is sparingly used in cases that are difficult to diagnose.⁴ The simultaneous development of vulvar and uterine leiomyomas is very common.⁹ In our case report, a transvaginal ultrasound evidenced an increase in uterine volume associated with the leiomyoma.

When the patient was first evaluated, the main hypothesis was a Bartholin cyst. Most vulvar leiomyomas are usually misdiagnosed as Bartholin cyst or abscess initially.^{4-6,10,11} After the surgery, a specimen was sent for a histopathologic examination that confirmed the vulvar leiomyoma diagnosis. Degenerative alterations are very common; some authors have reported myxoid degeneration in some cases of vulvar leiomyoma.^{12,13} However, no degenerative alteration was observed in our case.

These lesions need to be surgically excised, and the prognosis is good if they are completely removed.¹⁴ Close long-term follow-up is required because of the high risk of recurrence,⁸ but no recurrence was observed after one year of follow-up in this patient. Excision of the tumor with some of the surrounding normal tissue is the treatment of choice. It decreases the rate of recurrence and increases the five-year survival rate.⁸

In conclusion, vulvar leiomyomas are very rare. The macroscopic features of cystic lesions difficult the differential diagnosis between leiomyoma and Bartholin cyst. A histopathologic examination is often recommended for the final diagnosis.

Informed Consent

Consent was taken for the procedure and publication.

Conflicts of Interest

The authors of this manuscript have no conflicts of interest to disclose.

Contribution to Authorship

KAST was responsible for the conception and design of the study, the acquisition of data, and the draft of the article; TM and MT critically revised the article for important intellectual content; LMP critically revised the article for important intellectual content, and gave the final approval for the version that was submitted; CEF gave the final approval for the version that was submitted.

Note

This research received no specific grants from any public, private or non-profit funding agencies.

References

- Weston G, Healy DL. Uterine fibroids. In: Shaw RW, Luesley D, Monga A, editors. *Gynaecology*. 4th ed. Edinburgh: Elsevier Science; 2011. p. 473-85
- Hillard P. Benign diseases of the female reproductive tract. In: Berek J, Adashi E, Hillard P, editors. *Novak's gynecology*. 12th ed. Baltimore: Williams & Wilkins; 1996. p. 331-97.
- Kunhardt Urquiza E, de la Cruz SI, Fernández Martínez RL, Hernández Zúñiga VE. Myomatosis of rare localization. *Ginecol Obstet Mex* 1997;65(12):541-544
- Fasih N, Prasad Shanbhogue AK, Macdonald DB, et al. Leiomyomas beyond the uterus: unusual locations, rare manifestations. *RadioGraphics* 2008;28(07):1931-1948
- Reyad MM, Gazvani MR, Khine MM. A rare case of primary leiomyoma of the vulva. *J Obstet Gynaecol* 2006;26(01):73-74
- Zhao T, Liu X, Lu Y. Myxoid epithelial leiomyoma of the vulva: a case report and literature review. *Case Rep Obstet Gynecol* 2015; 2015:894830

- 7 Pandey D, Shetty J, Saxena A, Srilatha PS. Leiomyoma in vulva: a diagnostic dilemma. *Case Rep Obstet Gynecol* 2014; 2014:386432
- 8 Nielsen GP, Rosenberg AE, Koerner FC, Young RH, Scully RE. Smooth-muscle tumors of the vulva. A clinicopathological study of 25 cases and review of the literature. *Am J Surg Pathol* 1996; 20(07):779–793
- 9 Zlatkov V, Doganov N, Macaveeva V. is A case of simultaneous development of leiomyomas of the labia majora and the uterine corpus. *Rev Fr Gynecol Obstet* 1989;84(04):351–353
- 10 Youssef A, Neji K, M'barki M, Ben Amara F, Malek M, Reziga H. Leiomyoma of the vulva. *Tunis Med* 2013;91(01):78–80
- 11 Francis SA, Wilcox FL, Sissons M. Bartholin's gland leiomyoma: a diagnostic and management dilemma. *J Obstet Gynaecol Res* 2012;38(06):941–943
- 12 Aguilera Martínez V, Pérez Santana ME, Avila Contreras MdeL, Mendoza E. [Vulvar leiomyoma. Report of a case]. *Ginecol Obstet Mex* 2011;79(06):382–385Spanish.
- 13 Zhou J, Ha BK, Schubeck D, Chung-Park M. Myxoid epithelioid leiomyoma of the vulva: a case report. *Gynecol Oncol* 2006; 103(01):342–345
- 14 Spies JB, Bradley LD, Guido R, Maxwell GL, Levine BA, Coyne K. Outcomes from leiomyoma therapies: comparison with normal controls. *Obstet Gynecol* 2010;116(03):641–652