



Erratum for: “Participation of hepatic α/β -adrenoceptors and AT₁ receptors in glucose release and portal hypertensive response induced by adrenaline or angiotensin II” [Braz J Med Biol Res (2018) 51(12): e7526]

L.J.T. de Araújo¹, M.R. Nagaoka², D.R. Borges³ and M. Kouyoumdjian¹

¹Departamento de Bioquímica, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, SP, Brasil

²Departamento de Biociências, Universidade Federal de São Paulo, Baixada Santista, SP, Brasil

³Departamento de Medicina, Escola Paulista de Medicina, Universidade Federal de São Paulo, SP, Brasil

Erratum for: Braz J Med Biol Res | doi: 10.1590/1414-431X20187526

The Journal would like to correct the Author of Correspondence for the article “Participation of hepatic α/β -adrenoceptors and AT₁ receptors in glucose release and portal hypertensive response induced by adrenaline or angiotensin II” that was published incorrectly in volume 51 no. 12 (2018) in the Brazilian Journal of Medical and Biological Research < <http://dx.doi.org/10.1590/1414-431X20187526> >

The correct Author of Correspondence is: M. Kouyoumdjian: <mkouyoumdjian@gmail.com>



Participation of hepatic α/β -adrenoceptors and AT₁ receptors in glucose release and portal hypertensive response induced by adrenaline or angiotensin II

L.J.T. de Araújo¹, M.R. Nagaoka², D.R. Borges³ and M. Kouyoumdjian¹

¹Departamento de Bioquímica, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, SP, Brasil

²Departamento de Biociências, Universidade Federal de São Paulo, Baixada Santista, SP, Brasil

³Departamento de Medicina, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, SP, Brasil

Abstract

It has been previously demonstrated that the hemodynamic effect induced by angiotensin II (All) in the liver was completely abolished by losartan while glucose release was partially affected by losartan. Angiotensin II type 1 (AT₁) and adrenergic (α 1- and β -) receptors (AR) belong to the G-proteins superfamily, which signaling promote glycogen breakdown and glucose release. Interactive relationship between AR and AT₁-R was shown after blockade of these receptors with specific antagonists. The isolated perfused rat liver was used to study hemodynamic and metabolic responses induced by All and adrenaline (Adr) in the presence of AT₁ (losartan) and α 1-AR and β -AR antagonists (prazosin and propranolol). All antagonists diminished the hemodynamic response induced by Adr. Losartan abolished hemodynamic response induced by All, and AR antagonists had no effect when used alone. When combined, the antagonists caused a decrease in the hemodynamic response. The metabolic response induced by Adr was mainly mediated by α 1-AR. A significant decrease in the hemodynamic response induced by Adr caused by losartan confirmed the participation of AT₁-R. The metabolic response induced by All was impaired by propranolol, indicating the participation of β -AR. When both ARs were blocked, the hemodynamic and metabolic responses were impaired in a cumulative effect. These results suggested that both ARs might be responsible for All effects. This possible cross-talk between β -AR and AT₁-R signaling in the hepatocytes has yet to be investigated and should be considered in the design of specific drugs.

Key words: Liver perfusion; Angiotensin II; Adrenaline; AT₁R; Adrenoceptors

Introduction

The first observations of the renin-angiotensin system (RAS) and pressor effects in the kidney and its role in hypertension was made in 1898 by Tigerstedt and Bergman (1). In 1976, Borges and co-workers (2) described, for the first time, the hepatic conversion of angiotensin I to angiotensin II (All), followed by All inactivation. Further studies showed that All produces an increase in the hepatic portal pressure and metabolic responses. The hemodynamic effect was completely abolished by losartan (angiotensin II type 1 receptor (AT₁-R)-dependent mechanism) while metabolic responses such as glucose release and O₂ consumption were partially affected by losartan (AT₁-R-independent mechanism) (3). AT₁ and adrenergic (α 1- and β -) receptors (AR) belong to the G-proteins superfamily that promote, following signalization, an increase of intracellular calcium that culminates with the glycogen breakdown and glucose release. An interactive relationship between AR and AT₁-R was found after blockade of these receptors with specific antagonists (4,5). Exposure to elevated

catecholamines or All results in homologous desensitization of both adrenergic and AT₁-mediated vascular smooth contraction in the rat or rabbit aorta (6–8). This desensitization mediated by G-protein coupled receptors may result from changes in receptors, G proteins, carriers, or the interaction among these component systems (9,10). Therefore, selective antagonism of AR may clarify a possible alternative site for All interaction, leading to glucose release. The present work was designed to study the effects of AT₁R and AR blockade on All-induced hepatic glucose release and portal pressure.

Material and Methods

Animals

Adult male Wistar rats (*Rattus norvegicus albinus*) (270–320 g) obtained from Centro de Desenvolvimento de Modelos Animais para Medicina e Biologia (CEDEME) of the Universidade Federal de São Paulo (UNIFESP) were

Correspondence: M. Kouyoumdjian: <mkouyoumdjian@gmail.com>

Received June 13, 2018 | Accepted August 27, 2018

fed standard laboratory diet (Purina[®], Brasil) and water *ad libitum*. The animal experimental procedure was carried out in accordance to the guidelines of the Ethics in Research Committee of UNIFESP (CEP 1456/09).

Liver perfusion

Rat liver perfusion was performed as previously described (11). Briefly, the rat was anesthetized with 1.3 g/kg *ip* urethane (Sigma Chemical, USA). The abdominal and thoracic cavities were opened and the portal vein (entry via) and the inferior vena cava (exit via) were cannulated. The livers were exsanguinated and perfused (with no recirculation) with Krebs-Henseleit bicarbonate buffer (pH 7.5), containing 1 mg/mL BSA (Sigma Chemical), at 37°C, saturated with an oxygen/carbon dioxide mixture (95/5%) and at a constant flow (3–4 mL·min⁻¹·g⁻¹ liver). Liver viability was ensured by bile secretion and oxygen uptake monitored continuously by an oximeter (Delta OHM, Italy) connected to the efferent cannula during the experiment. The portal pressure was also continuously monitored with an open vertical column attached before the afferent cannula. After 20 min of stabilization previously determined (glucose release and portal pressure), 2 nmol All or 40 nmol Adr (Sigma Chemical) was injected *in bolus* through the portal vein cannula. Aliquots of perfusate were collected at 0 and every 30 s until 5, 6, 8, and 10 min for glucose determination.

The agonist-induced response was observed for 5 min in the absence or presence of 17.5 μM propranolol chlorhydrate (Pro; Medley, Brazil) (β-AR antagonist) administered by gavage 1 h before the experiment and/or 10 μM losartan potassium (Los; EMS, Brazil) (AT₁-R antagonist) and/or 25 μM prazosin chlorhydrate (Pra; Pfizer, USA) (α₁-AR antagonist) both added to Krebs solution 5 min before agonist injection.

Animals were divided into 12 experimental groups (6 for All and 6 for Adr): Control (absence of antagonists), Los, Pro, Pra, Los + Pro, and Pro + Pra. None of the antagonists *per se* altered the studied parameters.

Liver viability – bile production and oxygen consumption

Bile was collected over approximately two periods of 10 min (before and after agonist injection) and is reported as μL·min⁻¹·g⁻¹ liver. As bile production was similar (0.9 ± 0.04) in both periods, the average oxygen uptake was 2.0 ± 0.1 μmol/g in all protocols, to confirm liver viability.

Portal hypertensive response (PHR)

The mean values obtained for PHR 5 min before the agonist injection was considered baseline portal pressure. The difference between the pressure value observed after injection of the agonist at different times and baseline value was considered the portal pressure gain (cmH₂O). The graph of the portal pressure gain as a function of perfusion time (min) was used to calculate the area under

the curve (AUC) of portal pressure, which represents the PHR, reported as cmH₂O/min.

Glucose release (GluR)

The release of hepatic glucose was determined in the perfusate aliquots using the commercial kit Glucose PAP (Labtest[®], Brasil). Due to the absence of glucose in the perfusion fluid, we observed continuous and linear glucose release during the stabilization period, which was considered the baseline glucose release. The difference between baseline glucose release and following agonist injection was considered the gain of glucose release, mmol·min⁻¹·g⁻¹ liver. The graph of the gain as a function of perfusion time (min) was used to calculate the AUC of glucose release during the experiment, represented as GluR, reported as mmol/g liver.

Statistical analysis

Parameters were compared among groups by ANOVA and Newman-Keuls post-test with the level of significance set at P < 0.05. Data are reported as means ± SE. Analysis was performed using the GraphPad Prism software (version 6.0; Graph Pad Software, USA).

Results

Bile production (Adr: P=0.3686; All: P=0.0829) and oxygen consumption (Adr: P=0.6302; All: P=0.0648) were similar among groups, confirming liver viability. Both Adr and All induced an increase in the portal pressure and glucose release. The PHR of 40 nmol Adr and 2 nmol All were evaluated in the presence of AR and AT₁ antagonists. Figure 1A shows that all antagonists studied decreased the PHR induced by Adr compared to the control group. Los abolished the PHR induced by All (Figure 1B); on the other hand, AR antagonists had no effect when used alone. When antagonists were used together (Pro + Pra or Pro + Los) in the perfusion experiment, they caused a decrease in the PHR compared to the control group (P < 0.0001). Pra alone and the mixture of Pro + Pra or Pro + Los decreased the glucose release induced by Adr (Figure 2A) while Los or Pro alone and the mixture of Pro + Pra or Pro + Los decreased the metabolic effect induced by All (Figure 2B).

Discussion

Both β-AR and AT₁-R antagonists, as well as angiotensin-converting enzyme inhibitors are used as therapeutic drugs for several cardiac, renal, and vascular conditions, including hypertension. Although the liver is not a target organ and it might not be implicated directly in these diseases, AR and ATR are present in the cellular plasma membrane. Therefore, we studied the hepatic participation of AR and ATR in the portal hypertensive response and glucose release of Adr and All in the rat perfused liver.




Figure 1 Hepatic portal hypertensive response following *in bolus* injection of 40 nmol adrenaline (Adr) or 2 nmol angiotensin II (All). Portal hypertensive response (PHR) was calculated from the graphs “portal pressure gain × perfusion time”, after Adr or All injection, in the presence or absence of antagonists. Los: losartan; Pra: prazosin; Pro: propranolol. Data are reported as means ± SE. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001: compared with the control group; Adr: Pro vs Pra (**); Pro vs Pro + Pra (*); Pra vs Los (**); Pro + Pra vs Pro + Los (**). All: vs Pra (**); Pra vs Pro (*) and Pra vs Pro + Los (***) (ANOVA followed by Newman Keuls).




Figure 2. Glucose release from the perfused liver following *in bolus* injection of 40 nmol adrenaline (Adr) or 2 nmol angiotensin II (All). Glucose release was calculated from the graphs “glucose output x perfusion time”, after Adr or All injection, in the presence or absence of antagonists. Los: losartan; Pra: prazosin; Pro: propranolol. Data are reported as means ± SE. *P < 0.05, **P < 0.01, ***P < 0.001: compared with the control group; Adr: Pro vs Pra (*); All: Los vs Pra (**); Pro vs Pra (**); Pra vs Pro + Pra (*); Pra vs Pro + Los (***) (ANOVA followed by Newman Keuls).

Adr is a catecholamine that interacts with hepatic AR and signals by coupling to the stimulatory G-protein G α leading to activation of adenylyl cyclase and inducing

glycogen breakdown and glucose release through cAMP-dependent pathway (12,13). Furthermore, the stimulation of AR also increases the portal pressure, and this effect is

sensitive to α_1 - and β_2 -AR antagonists (14). Our results suggested that the GluR was mainly mediated by α_1 -AR, as reported recently by de Oliveira and co-workers (15). This high sensitivity to α_1 -adrenergic antagonists was also observed in other studies (14,15) and is strong evidence of predominant participation of α_1 -ARs in the liver (16). We also observed that the participation of β -ARs appears to have a secondary role, as it did not decrease the GluR. Los caused a significant decrease in the portal hypertensive response induced by Adr confirming the participation of AT₁R. Although there was not a significant decrease in the GluR in the presence of Los, a partial participation of AT₁R in this response might be important. Interestingly, when Los + Pro were added to the perfusion media, there was a sum of the effects, significantly decreasing the hemodynamic as well as the metabolic effect.

The hemodynamic effect (PHR) of All was abolished by Los while the GluR was only diminished, as reported previously (3). The GluR induced by All was also impaired by Pro indicating the participation of β -AR in this response. When both the ARs were blocked, the hemodynamic as well as the metabolic response was impaired showing a cumulative effect of the antagonists. Therefore, this result

showed that both ARs might also be responsible for All effects or there might be some sort of direct or indirect interaction impairing AT₁-R signaling. A study with mouse cardiomyocytes showed direct interaction between β -ARs and AT₁-Rs; this interaction would elicit a phenomenon by which selective β -AR antagonism inhibits signaling of AT₁-receptors, whereas selective AT₁-R antagonism inhibits downstream signaling of β -AR.

Moreover, the mechanism for this dual trans-inhibition of two independent receptors by a single antagonist might be via functional uncoupling of the signaling receptor from its cognate G protein (5). Therefore, whether this cross-talk between β -AR and AT₁-R signaling also occurs in the hepatocytes has not yet been investigated and should be considered in the design of specific drugs. Further experiments may explain the mechanisms of this interaction and might be important in the development of drugs highly specific for pathologies involving these vasoactive peptides.

Acknowledgments

The authors received financial support from FAPESP (grant #2011/13974-8) and CAPES (#33009015001P0).

References

- Phillips MI, Schmidt-Ott KM. The discovery of renin 100 years ago. *New Physiol Sci* 1999; 14: 271–274, doi: 10.1152/physiologyonline.1999.14.6.271.
- Borges DR, Limões EA, Prado JL, Camargo AC. Catabolism of vasoactive polypeptides by perfused rat liver. *Naunyn Schmiedeberg Arch Pharmacol* 1976; 295: 33–40.
- Nascimento EA, Gioli-Pereira L, Carvalho LT, Santos EL, Pesquero JB, Kouyoumdjian M, et al. Hemodynamic and metabolic effects of angiotensin II on the liver. *Peptides* 2005; 26: 315–322, doi: 10.1016/j.peptides.2004.09.017.
- Abdulla MH, Sattar MA, Khan MA, Abdullah NA, Johns EJ. Influence of sympathetic and AT-receptor blockade on angiotensin II and adrenergic agonist-induced renal vasoconstrictions in spontaneously hypertensive rats. *Acta Physiol (Oxf)* 2009; 195: 397–404, doi: 10.1111/j.1748-1716.2008.01895.x.
- Barki-Harrington L, Luttrell LM, Rockman HA. Dual inhibition of beta-adrenergic and angiotensin II receptors by a single antagonist: a functional role for receptor-receptor interaction in vivo. *Circulation* 2003; 108: 1611–1618, doi: 10.1161/01.CIR.0000092166.30360.78.
- Carrier O, Wedell EK, Barron KW. Specific alpha-adrenergic receptor desensitization in vascular smooth muscle. *Blood Vessels* 1978; 15: 247–258, doi: 10.1159/000158170.
- Silva EG, Ferreira AT, Paiva ACM. Angiotensin tachyphylaxis in normal and everted rings of rabbit aorta. *Eur J Pharmacol* 1988; 153: 185–190, doi: 10.1016/0014-2999(88)90605-X.
- Kuttan SC, Sim MK. Angiotensin II-induced tachyphylaxis in aortas of normo- and hypertensive rats: changes in receptor affinity. *Eur J Pharmacol* 1993; 232: 173–180, doi: 10.1016/0014-2999(93)90771-9.
- Suzuki E, Tsujimoto G, Hashimoto K. Different desensitization mechanisms of two alpha 1-adrenergic receptor subtypes in the contraction of rabbit aorta. *Br J Clin Pharmacol* 1990; 30: S121–S124, doi: 10.1111/j.1365-2125.1990.tb05481.x.
- Johnson MD, Wang HY, Ciechanowski D, Friedman E. Reduced G-protein function in desensitized rat aorta. *J Pharmacol Exp Ther* 1991; 259: 255–259.
- Borges DR, Kouyoumdjian M. The recognition factor for clearance of plasma kallikrein is located on its heavy chain and not exposed on prokallikrein. *J Hepatol* 1992; 16: 115–121, doi: 10.1016/S0168-8278(05)80103-5.
- Guimarães S, Moura D. Vascular adrenoceptors: an update. *Pharmacol Rev* 2001; 53: 319–356.
- Sharabi K, Tavares CD, Rines AK, Puigserver P. Molecular pathophysiology of hepatic glucose production. *Mol Aspects Med* 2015; 46: 21–33, doi: 10.1016/j.mam.2015.09.003.
- Peixoto JS, Comar JF, Moreira CT, Soares AA, Oliveira AL, Bracht A, et al. Effects of *Citrus aurantium* (bitter orange) fruit extracts and *p*-synephrine on metabolic fluxes in the rat liver. *Molecules* 2012; 17: 5854–5869, doi: 10.3390/molecules17055854.
- de Oliveira AL, de Paula MN, Comar JF, Vilela VR, Peralta RM, Bracht A. Adrenergic metabolic and hemodynamic effects of octopamine in the liver. *Int J Mol Sci* 2014; 14: 21858–21872, doi: 10.3390/ijms141121858.
- Reinhart PH, Taylor WM, Bygrave FL. Studies on alpha-adrenergic-induced respiration and glycogenolysis in perfused rat liver. *J Biol Chem* 1982; 257(4): 1906–1912, doi: 10.3390/molecules17055854.