

## Effects of extract, fractions and 2,3-dihydromyricetin-3-O- $\alpha$ -L-rhamnoside from *Pradosia huberi* (Ducke) Ducke on rat isolated mesenteric arteries

Alessandra A. N. Medeiros,<sup>1</sup> Fernando A. Medeiros,<sup>1</sup> Thyago M. Queiroz,<sup>2</sup>  
Josean F. Tavares, Marcelo S. Silva,<sup>2</sup> Isac A. Medeiros<sup>\*2</sup>

<sup>1</sup>Instituto de Pesquisas Científicas e Tecnológicas do Amapá, Av. Feliciano Coelho, Bairro do Trem, Caixa Postal 1509, 68900-260 Macapá-AP, Brasil,

<sup>2</sup>Laboratório de Tecnologia Farmacêutica "Delby Fernandes de Medeiros", Universidade Federal da Paraíba, Caixa Postal 5009, 58051-970 João Pessoa-PB, Brazil.

**RESUMO:** "Efeito do extrato, frações e da 2,3-diidromiricetina-3-O- $\alpha$ -L-raminosídeo obtidos da *Pradosia huberi* (Ducke) Ducke em artéria mesentérica isolada de rato". *Pradosia huberi* (Ducke) Ducke (Sapotaceae), espécie Amazônica popularmente conhecida como "casca-doce" é utilizada na medicina tradicional no tratamento de gastrite. O extrato etanólico de suas cascas é rico em polifenóis que podem apresentar um grande número de atividades, incluindo efeito vasorelaxante e cardioprotetor. O objetivo deste estudo foi avaliar as propriedades farmacológicas do extrato etanólico (EPH), de frações e da 2,3-diidromiricetina-3-O- $\alpha$ -L-raminosídeo isolados de *P. huberi*, em artéria mesentérica isolada de rato. O EPH foi fracionado resultando nas seguintes frações: CHCl<sub>3</sub>, CHCl<sub>3</sub>:AcOEt (1:1), AcOEt, AcOEt:MeOH (1:1) e MeOH. Da fração MeOH foi isolada a 2,3-diidromiricetina-3-O- $\alpha$ -L-raminosídeo e identificada através de espectro de RMN de <sup>1</sup>H e <sup>13</sup>C, além de comparações com os dados de literatura. EPH (1-100  $\mu$ g/mL) promoveu relaxamento dependente de concentração no tônus vascular induzido por 10  $\mu$ M de fenilefrina (CE<sub>50</sub>=17,1 $\pm$ 2,9  $\mu$ g/mL; E<sub>max</sub>=87,4 $\pm$ 2,9 %, n=8). A fração MeOH também relaxou os anéis mesentéricos (CE<sub>50</sub>=31 $\pm$ 2,0  $\mu$ g/mL; E<sub>max</sub>=54 $\pm$ 12,5%, n=6), porém com menor eficácia quando comparado ao efeito de EPH. Tanto o efeito de EPH com de MeOH foram completamente abolidos após a remoção do endotélio vascular. A fração AcOEt:MeOH (1:1) e o flavonoide isolado induziram vasorelaxamento. O estudo demonstrou que o EPH e a fração MeOH de *Pradosia huberi* apresentam propriedade vasorelaxante que pode ser completamente dependente da presença do endotélio. O flavonoide isolado não é o responsável por este efeito vasorelaxante.

**Unitermos:** *Pradosia huberi*, artéria mesentérica, vasodilatação, endotélio dependente, 2,3-diidromiricetina-3-O- $\alpha$ -L-raminosídeo, floresta amazônica.

**ABSTRACT:** *Pradosia huberi* (Ducke) Ducke (Sapotaceae), an Amazonian species, is popularly known as "casca-doce" and used in the folk medicine for the treatment of gastritis. The ethanol extract of the bark contains mainly polyphenolic compounds, which are known to show a large number of activities, including cardioprotective and vasorelaxant effects. The aim of this study was to evaluate the pharmacological properties induced by *P. huberi* ethanol extract (PHEE) and fractions and 2-3-dihydromyricetin-3-O- $\alpha$ -L-rhamnoside derived from this extract, in isolated rat mesenteric arteries. PHEE was separated and the following fractions were obtained: CHCl<sub>3</sub>, CHCl<sub>3</sub>:AcOEt (1:1), AcOEt, AcOEt:MeOH (1:1) and MeOH. We isolated 2-3-dihydromyricetin-3-O- $\alpha$ -L-rhamnoside from the MeOH fraction, which was identified by <sup>1</sup>H and <sup>13</sup>C NMR spectra and compared with data in the literature. PHEE (1-100  $\mu$ g/mL) induced concentration-dependent relaxations of 10  $\mu$ M phenylephrine-induced tone (EC<sub>50</sub>=17,1 $\pm$ 2,9  $\mu$ g/mL; E<sub>max</sub>=87,4 $\pm$ 2,9 %, n=8). The MeOH fraction also relaxed mesenteric rings (EC<sub>50</sub>=31 $\pm$ 2,0  $\mu$ g/mL; E<sub>max</sub>=54 $\pm$ 12,5%, n=6) but less effectively when compared to PHEE. Both effects were completely abolished after removal of the vascular endothelium. The AcOEt:MeOH (1:1) fraction and the isolated flavonoid were ineffective in eliciting vasorelaxation. The study demonstrates that PHEE and MeOH fraction of *Pradosia huberi* possess a vasorelaxant effect, which may be completely dependent upon endothelium. The isolated flavonoid is not responsible for this vasorelaxant effect.

**Keywords:** *Pradosia huberi*, mesenteric artery, vasodilatation, dependent-endothelium, 2,3-dihydromyricetin 3-O- $\alpha$ -L-rhamnoside, Amazon Rainforest.

## INTRODUCTION

*Pradosia huberi* (Ducke) Ducke, Sapotaceae is a medicinal plant that is common in the Amazon Rainforest, popularly known as casca-doce, pau-doce, amapá-doce or paracauba, and used in local folk medicine in the treatment of gastric and digestive problems. This species has the nomenclatural synonym *Glycoxylon huberi* Ducke (Corrêa, 1986). There are few phytochemical and biological studies of *P. huberi* reported in the literature; however, the hydroalcoholic extract from *P. huberi* bark has shown antisecretory and gastroprotective activity, besides no acute toxicity (5000 mg/kg; *p.o.*) (Kushima et al., 2005).

Phytochemical screening of the ethanolic extract was positive for the presence of compounds such as flavonoids, terpenoids, quinones, alkaloids, tannins and saponins (Ferreira et al., 2005). Flavonoids from stem bark were identified as 2,3-dihydromyricetin 3-O- $\alpha$ -L-rhamnoside, astilbin, engelitin and 2,3-dihydromyricetin (Jacquemin et al., 1985), all of which are also found in various plants. It has been reported that these flavonoids have pharmacological properties, including the following: anti-inflammatory (Kanbara et al., 1994; Yun et al., 2000), anti-oxidative effects (Yang et al., 2004), inhibition of lipid peroxidation (Yun et al., 2000), block of uterine contraction in rats (Carneiro et al., 1993).

Flavonoids are plant-derived polyphenolic substances commonly found in plants and consumed in the diet. Many of these compounds possess cardiovascular protective properties (Curin & Andriantsitohaina, 2005) which can be explained by the combination of the antioxidant, antiplatelet and anti-inflammatory effects along with their positive effects on restoration of endothelial function or modulation of vascular tone (Fitzpatrick et al., 1993; Woodman & Chan, 2004; Curin & Andriantsitohaina, 2005).

To date, this species has not been studied with regard to cardiovascular activity. Thus, the aim of this work was to evaluate the pharmacologic properties of *Pradosia huberi* ethanolic extract (PHEE), fractions and isolated substance for vasorelaxant activity in rat superior mesenteric artery.

## MATERIAL AND METHODS

### Plant material

The bark of *Pradosia huberi* (Ducke) Ducke, Sapotaceae, was collected in the city of Porto Grande, Amapá State, Brazil. The species was identified and a voucher specimen (N<sup>o</sup> 012519) was deposited in the Herbário Amapaense (HAMAB) of the Instituto de Pesquisas Científicas e Tecnológicas do Estado do Amapá (IEPA).

## Phytochemical study

After drying at 40 °C, the plant material was pulverized (3.2 kg) and extracted with 95% EtOH under maceration at room temperature for ten days. The solvent was removed by rotary evaporation under vacuum at 45 °C, yielding 200 g of *Pradosia huberi* ethanolic extract (PHEE). A sample of 20 g of PHEE was separated on a silica gel column under reduced pressure with the eluents CHCl<sub>3</sub>, CHCl<sub>3</sub>:AcOEt (1:1), AcOEt, AcOEt:MeOH (1:1) and MeOH, resulting in the following yields: 0.2, 0.4, 1.2, 3.0 and 4.0 g, respectively. Samples of each fraction obtained were used for pharmacological testing. The MeOH fraction was chosen for isolation since it presented the greatest amount of constituent and showed the highest vasorelaxant activity compared to the other fraction obtained. An aliquot of 2 g of the MeOH fraction was separately submitted to Sephadex LH-20 column chromatography using MeOH as eluent, from which 28 fractions of 30 mL were collected, which after analysis by TLC were grouped according to their R<sub>f</sub>. The fraction 14-22 was rechromatographed on Sephadex LH-20 with MeOH elution, isolating 2,3-dihydromyricetin-3-O- $\alpha$ -L-rhamnoside (45 mg, Figure 1), which corresponded to 22.5% of the yield in relation to the MeOH fraction. For chemical identification of the isolated compound, <sup>1</sup>H and <sup>13</sup>C NMR spectra were acquired using a Mercury Varian spectrometer operating at 200 MHz for <sup>1</sup>H and 50.3 MHz for <sup>13</sup>C NMR and recorded in CD<sub>3</sub>OD.

### Animals

Male Wistar rats (250-300 g) were used for the experiments. Animals were housed under conditions of controlled temperature (21±1 °C) and lighting (light-dark cycle of 12 h), with free access to water and pelleted feed (Purina-Brazil). The study was approved by the Animal Care and Use Committees of the Federal University of Paraíba (N<sup>o</sup> 0603/07).

### Drugs

The drugs used were L-phenylephrine chloride and acetylcholine chloride (both from Sigma, St. Louis, MO, USA). For the experiments, PHEE was dissolved in distilled water. All the stock solutions were prepared in distilled water and kept at 4 °C.

### Preparation of isolated rat superior mesenteric artery rings

The superior mesenteric arteries were removed and cleaned free of connective tissue and fat. Mesenteric rings (1-2 mm) were obtained and suspended by cotton threads in an organ bath containing 10 mL Tyrode's solution (pH 7.4), maintained at 37 °C and gassed with a 95% O<sub>2</sub>

+ 5% CO<sub>2</sub> mixture. Rings were stabilized under a resting tension of 0.75 g for 1 h. During this time the solution was changed every 15 min to prevent the accumulation of metabolites that could otherwise lead to misinterpretations (Altura & Altura, 1970). The isometric contraction was recorded by a force transducer (Miobath-4, WPI, Sarasota, FL, EUA) coupled to an amplifier-recorder (Miobath-4, WPI, Sarasota, FL, EUA) and to a personal computer equipped with an analog-to-digital converter board. In some experiments, the endothelium layer was removed by gently rubbing the intimal surface of the vessels with a cotton ball. The presence of functional endothelium was assessed by the ability of acetylcholine (10 μM) to induce more than 90% relaxation of vessels pre-contracted with 10 μM phenylephrine (PHE), and the absence of relaxation in response to acetylcholine was taken as evidence that the vessel segments were functionally denuded of endothelium (Furchgott; Zawadzki, 1980). PHEE was cumulatively applied after contractile responses induced by PHE (10 μM).

#### Effect of PHEE, MEF, MAF and 2,3-dihydromyricetin-3-O-α-L-rhamnoside on sustained contractions induced by phenylephrine (10 μM) in isolated preparations from rat superior mesenteric arteries


After an equilibration period, the rings with or without functional endothelium were pre-contracted with the agonist, and once the response to the second administration of PHE (10 μM) reached a plateau, increasing cumulative concentrations of PHEE (1-100 μg/mL), MEF (1-100 μg/mL), MAF (1-100 μg/mL) or 2,3-dihydromyricetin-3-O-α-L-rhamnoside (1-100 μg/mL) were added to the bath. The relaxations were measured by comparing the tension developed before and after addition of PHEE, MEF, MAF or 2,3-dihydromyricetin-3-O-α-L-rhamnoside.

#### Data analysis

Values are expressed as means ±S.E.M. When appropriate, statistical significance was examined with Student's t -test or one-way ANOVA followed by Bonferroni's post-hoc test, using Graph Pad Prism TM 4.0 software. The EC<sub>50</sub> values were calculated by nonlinear regression of individual concentration–response curves, and  $p < 0.05$  was considered significant.

## RESULTS

The analyses of the spectral data as well as the assignments of all carbons and hydrogens (Table 1) and comparison with literature values (Gellért et al., 1981; Jacquemin et al., 1985; Shen et al., 1993; Slimestad et al., 1994; Wu et al., 1998; Du et al., 2005) allowed the identification of 2,3-dihydromyricetin-3-O-α-L-rhamnoside (**1**).



#### Relaxant effect of *Pradosia huberi* extract, fractions and isolated constituent on the PHE-induced sustained contractions


PHEE inhibited PHE-induced sustained contraction in the rat mesenteric rings in a concentration-dependent manner, in the preparations with preserved functional endothelium. EC<sub>50</sub> of PHEE effect on contraction induced by PHE was 17.1±2.9 μg/mL, and the maximal value for the relaxant effect ( $E_{max}$ ) was 87.4±2.9%,  $n=8$ ,  $p < 0.001$ \*\*\*. The concentration-response curve of PHEE was completely abolished after removal of functional endothelium (Figure 1 and Figure 2A). MEF was also able to induce concentration-dependent relaxations of 10 μM PHE-induced tone with an EC<sub>50</sub> value of 31±2.0 μg/mL. This vasodilator effect was significantly smaller when compared to that induced by PHEE, with a strong reduction in the  $E_{max}$  of 54±12.5%,  $n=6$ , as shown in Figure 2B. MEF was not able to induce relaxation in the rings without functional endothelium.

MAF did not induce a concentration-dependent vasodilator effect in the preparations, either with intact endothelium or after endothelium removal (Figure 2C). On the contrary, there was a tendency toward contraction of the rat mesenteric rings with intact endothelium at the highest concentrations of this fraction. In addition, the isolated flavonoid (2,3-dihydromyricetin-3-O-α-L-rhamnoside) was not capable of inhibiting the contractions induced by phenylephrine in the isolated preparations. On the contrary, a tendency toward contraction was observed in preparations with intact endothelium (Figure 2D).

## DISCUSSION


The present study showed that PHEE exerted a vasorelaxant effect in phenylephrine-induced contractions of rat superior mesenteric rings. Removal of functional endothelium completely abolished this relaxant response to PHEE, suggesting that vasorelaxation caused by PHEE was endothelium-dependent. Furthermore, the isolated flavonoid was not responsible for this vasorelaxant effect.

The <sup>13</sup>C NMR (APT) spectrum showed a total of twenty-one signals, including nine to no hydrogenated



**Figure 1.** Relaxant effects of PHEE on isolated mesenteric rings pre-contracted with phenylephrine. Panel A shows a typical recording obtained in intact endothelium rings, and panel B in removed endothelium rings.

carbons, eleven to methynic carbons and a methyl carbon. The spectral region between  $\delta$ C 168.9 and  $\delta$ C 95.8, characteristic of aromatic carbon signals, and the signal at  $\delta$ C 197.5, characteristic of carbonyl carbon signals, as well as comparison with literature data (Shen et al., 1993; Wu et al., 1998; Almeida et al., 2005; Du et al., 2005; Sinkkonen et al., 2005) and chemotaxonomy of the genus *Pradosia*, suggest a flavonoid skeleton such as an aglycone. Signals at  $\delta$ C 83.7 C and  $\delta$ C 76.8 are compatible with a dihydroflavonol structure, which can be supported by the  $^1$ H NMR spectrum showing doublets at  $\delta$ H 4.86 and  $\delta$ H 4.61 (each  $J = 11.0$  Hz), characteristic of H-2 and H-3 of the dihydroflavonol structure (Jacquemin et al., 1985; Wu et al., 1998; Du et al., 2002). The determination of the stereochemical trans-axial relationship between the protons at C-2 and C-3 was evident from the 11.0 Hz coupling constant (Jacquemin et al., 1985; Slimestad et al., 1994; Wu et al., 1998; Du et al., 2005). The oxymethynic carbon signals between  $\delta$ C 69.2-71.9, together with the methyl carbon signal at  $\delta$ C 17.8 (H-6", at  $\delta$ H 0.92, d,  $J = 6.2$  Hz, 3H) indicated that the rhamnose sugar was attached. The  $^{13}$ C NMR spectrum showed a downfield shift of 5.2 ppm for C-3 when compared with the data of 2,3-dihydromyricetin (Shen et al., 1993), indicating the location of the rhamnose moieties to be the C-3 (Slimestad et al., 1994; Wu et al., 1998; Du et al., 2005). The  $\alpha$ -configuration of rhamnose was established



**Figure 2.** Line plot showing the effects of increasing concentrations of PHEE (A), MEF (B), MAF (C) or isolated compounds of *Pradosia huberi* (D) on phenylephrine (10 $\mu$ M)-induced contraction in mesenteric rings of rats with and without the functional endothelium. Results are means  $\pm$  S.E.M..



**Table 1.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data of 2,3-dihydromyricetin-3-*O*- $\alpha$ -L-rhamnoside ( $\delta$  (ppm),  $J$  (Hz), measured in  $\text{CD}_3\text{OD}$ ).

C	2,3-dihydromyricetin-3- <i>O</i> - $\alpha$ -L-rhamnoside		1		2	
	$\delta\text{H}$	$\delta\text{C}$	$\delta\text{H}$	$\delta\text{C}$	$\delta\text{H}$	$\delta\text{C}$
4		197.5		197.2		194.3
5		165.4		163.2		163.3
7		168.9		166.7		166.9
9		164.2		162.4		162.1
10		101.9		100.4		101.0
1'		129.0		127.1		126.8
3'		147.0		145.6		145.8
4'		135.2		133.4		145.1
5'		147.0		145.6		
CH						
2	4.86 (d, $J=11,0$ Hz)	83.7	4.90 (d, $J=10.5$ Hz)	83.2	5.24 (d, $J=9.8$ Hz)	81.5
3	4.61 (d, $J=11,0$ Hz)	76.8	4.38 (d, $J=10.5$ Hz)	71.6	4.63 (d, $J=9.8$ Hz)	75.6
6	5.90 (d, $J=2,2$ Hz)	96.3	5.89 (d, $J=1.6$ Hz)	95.9	5.90 (d, $J=2.1$ Hz)	96.0
8	5.87 (d, $J=2,2$ Hz)	95.8	5.85 (d, $J=1.6$ Hz)	94.9	5.88 (d, $J=2.1$ Hz)	95.0
2''	6.50 (s)	108.0	6.40 (s)	106.9	6.88 (s)	114.7
5''					6.74 (s)	115.3
6''	6.50 (s)	108.0	6.40 (s)	106.9	6.74 (s)	118.7
1'''	4.17 (d, $J=2,8$ Hz)	102.7			4.07 (s)	100.0
2'''	4.00 (dd, $J=3,2; 1,4$ Hz)	71.9			3.36 (br, s)	70.1
3'''	3.41 (dd, $J=9,8; 3,0$ Hz)	71.9			3.42 (dd, $J=9,4; 2,8$ Hz)	70.4
4'''	3.20 (dd, $J=9,2; 9,2$ Hz)	70.3			3.15 (dd, $J=9,4; 9,4$ Hz)	71.6
5'''	2.40 (dd, $J=9,4; 6,2$ Hz)	69.2			3.88 (qd, $J=9,4; 6,2$ Hz)	68.9
CH <sub>3</sub>						
6'''	0.92 (d, $J=6,2$ Hz)	17.8			1,05 (d, $J=6,2$ Hz)	17.6

(1) 2,3-dihydromyricetin (Shen, et al., 1993) and (2) 2,3-dihydroquercetin-3-*O*- $\alpha$ -L-rhamnoside (Du et al., 2005)

by the anomeric proton at  $\delta\text{H}$  4.17 and from the 2.8 Hz coupling constant (Slimestad et al., 1994; Wu et al., 1998).

The endothelium is formed by a monolayer of cells that covers the lumen of blood vessels and serves as a secretory gland able to produce contractant as well as relaxing factors that control vascular tone (Curin & Andriantsitohaina, 2005). Under physiological conditions, there is a balance between endothelial factors released, where the effect of relaxing agents prevails. These factors include nitric oxide (NO), endothelium-derived hyperpolarizing factor (EDHF) and prostacyclin ( $\text{PGI}_2$ ) (Moncada & Vane, 1979; Furchgott & Zawadzki, 1980; Feletou & Vanhoutte, 1988).

Many reports show that the effect of polyphenols on the endothelium is mainly due to NO production (Andriambeloson et al., 1997; Duarte et al., 2004; Zenebe et al., 2003), increase in intracellular concentration of  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ), activation of  $\text{K}^+$  channels in the endothelium, inhibition of  $\text{Ca}^{2+}$ -ATPases of the endoplasmic reticulum in endothelial cells (Li et al., 2000; McKenna et al., 1996), or modulation of NO levels by the action on the phosphodiesterases (PDE)

PDE-2 and PDE-4 in endothelial cells (Beretz et al., 1986a; Beretz et al., 1986b; Lugnier & Schini, 1990).

MEF also induced a concentration-dependent relaxation of the preparations pre-contracted with phenylephrine, only in intact endothelium rings ( $\text{EC}_{50}=31\pm 2.0$   $\mu\text{g}/\text{mL}$ ;  $E_{\text{max}}=54\pm 12.5\%$ ,  $n=6$ ). However, such effect was shown to be less potent and effective when compared to the effect produced by PHEE ( $\text{EC}_{50}=17,1\pm 2,9$   $\mu\text{g}/\text{mL}$ ;  $E_{\text{max}}=87.4\pm 2.9\%$ ,  $n=8$ ).

Both MAF, with the majority substances of the ethanol extract of *Pradosia huberi*, and 2,3-dihydromyricetin 3-*O*- $\alpha$ -L-rhamnoside were not effective in relaxing mesenteric rings.

A particular feature of phytomedicines is their complex composition, *i.e.*, the "phytochemical complex" which includes a variety of phytochemicals with different biological activities. Some of these phytochemicals are responsible for specific effects, while other components play an additional role. However, a wider array of effects and the healing properties are frequently guaranteed only by the phytocomplex (Pietta, 2000).

We can conclude that PHEE possesses a vasorelaxant

effect in isolated mesenteric rings and that this effect is totally dependent on the vascular endothelium. The loss of activity of the fractions and 2,3-dihydromyricetin-3-O- $\alpha$ -L-rhamnoside may be due to the action of the constituents present in PHEE (phytoextract). Since the extract consists primarily of flavonoids, these data are in line with the literature that show an endothelium-dependent vasodilator effect of flavonoids and other polyphenols (Fitzpatrick et al., 1993; Rice-Evans et al., 1996; Lemos et al., 1999).

## ACKNOWLEDGEMENTS

The authors thank Instituto de Pesquisas Científicas e Tecnológicas do Estado do Amapá (IEPA/GEA), CNPq and Laboratório de Tecnologia Farmacêutica (LTF/UFPB) for the financial or technical support of this work.

## REFERENCES

- Almeida SCX, Lemos TLG, Silveira ER, Pessoa ODL 2005. Constituintes químicos voláteis e não-voláteis de *Cochlospermum vitifolium* (Willdenow) Sprengel. *Quim Nova* 28: 57-60.
- Altura BM, Altura BT 1970. Differential effects of substrate depletion on drug induced contractions of rabbit aorta. *Am J Physiol* 219: 1698-1705.
- Andriambeloso E, Kleschyov AL, Muller B, Beretz A, Stoclet JC, Andriantsitohaina R 1997. Nitric oxide production and endothelium-dependent vasorelaxation induced by wine polyphenols in rat aorta. *Br J Pharmacol* 120: 1053-1058.
- Beretz A, Anton R, Cazenave JP 1986a. The effects of flavonoids on cyclic nucleotide phosphodiesterases. *Prog Clin Biol Res* 213: 281-296.
- Beretz A, Briancon-Scheid F, Stierle A, Corre G, Anton R, Cazenave JP 1986b. Inhibition of human platelet cyclic AMP phosphodiesterase and of platelet aggregation by a hemisynthetic flavonoid, amentoflavone hexaacetate. *Biochem Pharmacol* 35: 257-262.
- Carneiro E, Calixto J, Delle M, Franco Y, Rosendo A 1993. Isolation, chemical identification, and pharmacological evaluation of eucryphin, astilbin, and engelitin obtained from the bark of *Hymenaea martiana*. *Int J Pharmacol* 31: 38-46.
- Corrêa MP 1986. *Dicionário das plantas úteis do Brasil e das exóticas cultivadas*. Rio de Janeiro: Ministério da Agricultura. IBDF.
- Curin Y, Andriantsitohaina R 2005. Polyphenols as potential therapeutic agents against cardiovascular diseases. *Pharmacol Rep* 57: 97-107.
- Du Q, Cai W, Xia M, Ito YG 2002. Purification of (+)-dihydromyricetin from leaves of *Ampelopsis grossedentata* using high-speed countercurrent chromatography with scale-up triple columns. *J Chromatogr A* 973: 217-220.
- Du Q, Li L, Jerz G 2005. Purification of astilbin and isoastilbin in the extract of *Smilax glabra* rhizome by high-speed countercurrent chromatography. *J Chromatogr A* 1077: 98-101.
- Duarte J, Andriambeloso E, Diebolt M, Andriantsitohaina R 2004. Wine polyphenols stimulate superoxide anion production to promote calcium signaling and endothelium-dependent vasodilatation. *Physiol Res* 53: 595-602.
- Félétou M, Vanhoutte PM 1988. Endothelium-dependent hyperpolarization of canine coronary smooth muscle. *Br J Pharmacol* 93: 515-524.
- Ferreira ES, Medeiros FA, Medeiros AAN 2005. Estudo fitoquímico da espécie *Pradosia huberi* (Ducke): usada na produção de fitoterápicos do IEPA. *Rev Cien Soc* 2: 43-53.
- Fitzpatrick DF, Hirschfield SL, Coffey RG 1993. Endothelium-dependent vasorelaxing activity of wine and other grape products. *Am J Physiol* 265: H774-H778.
- Furchgott RF, Zawadzki JV 1980. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 288: 373-376.
- Gellért M, Szendrei K, Reisch J 1981. Dihydromyricetin-3-O-rhamnoside from leaves of *Catha edulis*. *Phytochemistry* 20: 1759-1760.
- Jacquemin H, Boissonnat A, Faugeras G, Tillequin F, Delaveau P 1985. Flavonoids of *Glycoxydon huberi* Ducke. *Ann Pharm Fr* 43: 521-522.
- Kanbara T, Mizutani K, Tamura K, Kataoka S 1994. Extraction of astilbins from *Engelhardtia chrysolepis* leaves and their anti-inflammatory activities. *Jpn Kokai Tokkyo Koho JP* 06,256,194.
- Kushima H, Hurima-Lima CA, Santos MA, Viana E, Coelho-Ferreira, Brito ARMS 2005. Gastroprotective activity of *Pradosia huberi* on experimentally induced gastric lesions in rodents: Role of endogenous sulphhydryls and nitric oxide. *J Ethnopharmacol* 101: 61-67.
- Lemos VS, Freitas MR, Muller B, Lino YD, Queiroga CE, Cortes SF 1999. Dioclein, a new nitric oxide- and endothelium-dependent vasodilator flavonoid. *Eur J Pharmacol* 386: 41-46.
- Li HF, Chen SA, Wu SN 2000. Evidence for the stimulatory effect of resveratrol on  $Ca^{2+}$ -activated  $K^+$  current in vascular endothelial cells. *Cardiovasc Res* 45: 1035-1045.
- Lugnier C, Schini VB 1990. Characterization of cyclic nucleotide phosphodiesterases from cultured bovine aortic endothelial cells. *Biochem Pharmacol* 39: 75-84.
- McKenna E, Smith JS, Coll KE, Mazack EK, Mayer EJ, Antanavage J, Wiedmann RT, Johnson RG Jr 1996. Dissociation of phospholamban regulation of cardiac sarcoplasmic reticulum  $Ca^{2+}$ -ATPase by quercetin. *J Biol Chem* 271: 24517-24525.
- Moncada S, Vane JR 1979. Pharmacology and endogenous roles of prostaglandins endoperoxides, thromboxane A2 and prostacyclin. *Pharmacol Rev* 30: 293-331.
- Pietta P 2000. Phytomedicines creating safer choices. In: Watson RR *Vegetables, fruits and herbs in health promotion*. 1.ed. Florida: CRC Press LLC, p.73-74.
- Rice-Evans CA, Miller NJ, Paganga G 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids.

*Free Radical Biol Med* 20: 933-956.

- Shen C, Chang Y, Ho LK 1993. Nuclear magnetic resonance studies of 5,7-dihydroflavonoids. *Phytochemistry* 34: 843-845.
- Sinkkonen J, Liimatainen J, Koronen M, Pihlaja K 2005. Spectral assignments and reference data: a new dihydroflavonol from *Pinus sylvestris* L. *Magn Reson Chem* 43: 348-349.
- Slimestad R, Andersen OM, Francis GW 1994. Ampelopsin-7-O-glucoside and other dihydroflavonol-7-O-glucosides from needles of *Picea abies*. *Phytochemistry* 35: 550-552.
- Woodman OL, Chan ECH 2004. Vascular and anti-oxidant of flavonols and flavones. *Clin Exp Pharmacol Physiol* 31: 786-790.
- Wu QL, Whang SP, Du LJ, Zhang SM, Yang JS, Xiao PG 1998. Cromone glycosides and flavonoids from *Hipericum japonicum*. *Phytochemistry* 49: 1417-1420.
- Yang S, Zhang Y, Peng L, Ning Z, Zhang J 2004. Antioxidant properties of dihydromyricetin. *Zhongguo Liangyou Xuebao* 19: 82-84, apud *Chemical Abstracts* 129: 289-360g.
- Yun BS, Lee IK, Kim JP, Chung SH, Shim GS, Yoo ID 2000. Lipid peroxidation inhibitory activity of some constituents isolated from the stem bark of *Eucalyptus globulus*. *Arch Pharmacol Res* 23: 147-150.
- Zenebe W, Pechanova O, Andriantsitohaina R 2003. Red wine polyphenols induce vasorelaxation by increased nitric oxide bioactivity. *Physiol Res* 52: 425-432.