



# The Brazilian Journal of INFECTIOUS DISEASES

[www.elsevier.com/locate/bjid](http://www.elsevier.com/locate/bjid)



## Original article

# JC virus/human immunodeficiency virus 1 co-infection in the Brazilian Amazonian region



Izaura Maria Vieira Cayres-Vallinoto\*, Antonio Carlos Rosário Vallinoto,  
Giselle Priscila dos Anjos Pena, Vânia Nakauth Azevedo,  
Luiz Fernando Almeida Machado, Marluísa de Oliveira Guimarães Ishak, Ricardo Ishak

Universidade Federal do Pará (UFPA), Instituto de Ciências Biológicas, Laboratório de Vírus, Belem, Para, Brazil

### ARTICLE INFO

#### Article history:

Received 14 December 2015

Accepted 10 May 2016

Available online 5 June 2016

#### Keywords:

JCV

HIV-1

Co-infection

Brazilian Amazon region

### ABSTRACT

JC virus (JCV) is a member of the *Polyomaviridae* family and is associated to a severe disease known as progressive multifocal leukoencephalopathy, PML, which is gradually increasing in incidence as an opportunistic infection among AIDS patients. The present study aimed to investigate the occurrence of JCV among HIV-1 carriers including their types and molecular subtypes and the possible association with disease. Urine samples from 66 HIV-1 infected subjects were investigated for the presence of the virus by amplifying VP1 (215 bp) and IG (610 bp) regions using the polymerase chain reaction. JCV was detected in 32% of the samples. The results confirmed the occurrence of type B (subtype Af2); in addition, another polyomavirus, BKV, was also detected in 1.5% of samples of the HIV-1 infected subjects. Apparently, there was no significant difference between mono- (HIV-1 only) and co-infected (HIV-1/JCV) subjects regarding their TCD4<sup>+</sup>/TCD8<sup>+</sup> lymphocyte counts or HIV-1 plasma viral load. Self admitted seizures, hearing and visual losses were not significantly different between the two groups.

© 2016 Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Polyomaviruses are small agents which infect a variety of species including man, non-human primates, cattle, rodents, rabbits, and birds. Two human polyomaviruses are important pathogens, JC virus (JCV) and BK virus (BKV), as they are able to cause a severe, usually fatal, neurological disease (progressive multifocal leukoencephalopathy, PML) and renal infection in immunosuppressed patients, respectively.<sup>1–3</sup> The viruses share

approximately 72% of nucleotide homology and a high degree of identity (80%) of amino acid sequence of a capsid protein, VP1.<sup>4,5</sup> JCV infects the human host and establishes a persistent replicative cycle (including latency) following the acute infection, which favors its dissemination within human populations and its carriage through generations.

Although JCV is found to infect more than 70% of the adult human population as a productively persistent infection, PML is a rare disease.<sup>5–9</sup> It is a manifestation that follows malignant conditions, as well as a consequence of

\* Corresponding author.

E-mail address: [ivallinoto@ufpa.br](mailto:ivallinoto@ufpa.br) (I.M. Cayres-Vallinoto).

<http://dx.doi.org/10.1016/j.bjid.2016.05.005>

1413-8670/© 2016 Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

primary or secondary immunodeficiencies, particularly in acquired immunodeficiency syndrome (AIDS). The emergence of human immunodeficiency virus 1 (HIV-1) infection elicited an additional interest in JCV with the frequent finding of HIV-1 carriers co-infected with JCV.<sup>10</sup> The incidence of PML is gradually increasing among AIDS patients, initially as a consequence of the increase of HIV-1 infections, and secondly, as an increase in life expectancy as a result of better antiretroviral drugs.<sup>10</sup> JCV infects HIV-1 carriers and is one of the major opportunistic infections of the central nervous system.<sup>11–15</sup> In our previous study we described JCV and BKV infections among Amerindian tribes, Afro-descendant communities, as well as in the urban area of Belém, state of Para, north of Brazil.<sup>16</sup> The results showed a large genetic variability of strains circulating in the region, infecting a large group of individuals. The presence of European, Asiatic, and African subtypes associated to the ethnic origin of the population samples investigated highlighted the idea that JCV is a good marker for studying the early migration of human populations, reflecting their early and late history.

The present study aimed to describe the occurrence of JCV, including their types and molecular subtypes among HIV-1 carriers and the possible association with neurological disease.

## Materials and methods

### Population examined

A cross-sectional study was performed, using a convenience sample which included 66 HIV-1 infected persons. A single urine sample was obtained from HIV-1 infected persons residing in the city of Belém, attending the University Hospital João de Barros Barreto (HUJBB), for their routine clinical care, and laboratory follow-up determinations of TCD4<sup>+</sup>/TCD8<sup>+</sup> lymphocyte counts and HIV-1 plasma viral load (HPVL). All patients were receiving antiretroviral therapy (ART) at the time of sample collection. The duration of ART ranged from one to seven years. Half of them were on ART for no more than five years, with mean and median values of four years. The project was approved by the Ethics Committee of the University Hospital Joao de Barros Barreto (Protocol Number 2092/05). The participants were briefed about the project and those who accepted to take part signed an informed consent. All patients were adults (18 years or older) and were capable to provide answers to a questionnaire which included questions regarding demographic, social and cultural aspects of their life style. The patients were also asked to inform, over common neurological events which are usually considered as baseline symptoms associated to JCV infection, at least once, since their initial diagnosis of HIV-1 infection.

### Quantification of HPVL and TCD4<sup>+</sup>/TCD8<sup>+</sup> lymphocyte counts

HPVL was determined by the bDNA method (VERSANT<sup>®</sup> HIV-1 RNA 3.0 Assay bDNA “Versant Assay”, Siemens, USA) following the manufacturer’s directions. Whole blood samples were

processed within four hours of collection for the determination of TCD4<sup>+</sup> counts by flow cytometry (FacsCount, Becton & Dickinson, San Jose, CA, USA) using the FacsCount<sup>™</sup> Reagents immunomonitoring kit, following the protocol recommended by the manufacturer (Becton Dickinson, San Jose, CA, USA). Both procedures were routinely conducted at the Virology Laboratory, which is one of the National Reference Laboratory for the two determinations.

### Urine sample collection and DNA extraction

Urine samples (50 mL) were collected in specific vial collectors and transported to the Virology Laboratory of the Institute of Biological Sciences of the Universidade Federal do Para (ICB/UFGPA) and stored at –20 °C until assaying for JCV infection. The urine samples were centrifuged at 5000 rpm for 15 min, the urinary sediment washed three times with sterile saline solution and the cell pellet used for DNA extraction according to the protocol using the EZ-DNA kit (Gentra Systems, Inc., Minneapolis, MN, USA), as previously described.<sup>16</sup>

### Molecular analysis

Polymerase chain reaction – PCR was performed for amplification of the VP1 gene (215 bp) and the genomic region IG (610 bp) of JCV, using a thermocycler (Mastercycler Personal, Eppendorf, Germany), and the amplified products were subjected to purification for subsequent sequencing analysis of the nucleotide bases, as previously described.<sup>16</sup>

### Sequencing and subtype analysis

The amplified products of the IG region were used for determining genetic relationships among the strains detected and those recorded in Genbank database. The amplified fragments were submitted to a direct sequencing assay (both forward and reverse) according to the protocol of the ABI Prism Dye Terminator Cycle Sequencing Ready Kit (Life Technologies, Foster City, CA, USA) and the products were loaded on the ABI Prism 310 DNA Sequencer (Life Technologies, Foster City, CA, USA). The nucleotide sequences were used together with other JCV sequences available in the Genbank as previously described.<sup>16</sup> Nucleotides sequences obtained in the present study are available in the Genbank databases under the accession numbers KR062063, KR062064 and KR062065 and were submitted to Basic Local Alignment Search Tool – Blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) aiming to identify the virus subtype.

### Statistical analysis

The frequency of HIV-1/JCV co-infection was estimated by direct counting. To determine whether the variables TCD4<sup>+</sup>, TCD8<sup>+</sup> and HIV-1 viral load followed a normal distribution, D’Agostino’s test was used, and the mean values of the laboratory markers were compared using ANOVA. A p-value <0.05 was considered statistically significant using the BioEstat 5.0 software.<sup>17</sup>

**Table 1 – Distribution of HIV-1 carriers investigated for JCV infection according to sex and age.**

Age range	Male (%)	Female (%)	Total (%)
11–20	0	1	1
21–30	10	6	16
31–40	18	9	27
41–50	10	4	14
>50	3	1	4
No information	3	1	4
Total	44 (66.7)	22 (33.3)	66 (100)

## Results

JCV infection was investigated in 66 urine samples of HIV-1 carriers, outpatients and inpatients, attending the University Hospital. Table 1 shows that most of the patients were male (66.7%), within the age range of 21–50 years old (86.4%).

Amplification of VP1 was obtained in 21 (32%) samples (18 males and 3 females) and 11 of them amplified the two regions, VP1 and IG. Sequencing of nine samples from IG region submitted to Basic Local Alignment Search Tool – Blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) were identified as belonging to type B, subtype Af2, with 100% identity, confirming the JCV/HIV-1 co-infection. The molecular analysis also showed the amplification of a nucleotide sequence of 541 bp, characteristic of BKV polyomavirus, showing 97.4% identity with KOM-3 line and a co-infection BKV/HIV-1 in 1.5%.

TCD4<sup>+</sup> and TCD8<sup>+</sup> lymphocyte counts and HPVL (Fig. 1) were compared between the mono-infected (HIV-1 only) and co-infected (JCV/HIV-1) patients. The average counts of TCD4<sup>+</sup> were 130.61 and 105.25 cells/mL ( $p=0.5649$ ), and the values for TCD8<sup>+</sup> were 589.97 and 679.66 cells/mL ( $p=0.2416$ ), respectively. The mean values for HPVL ( $\log_{10}$ ) were 4.563 and 4.867 ( $p=0.5605$ ), respectively (Fig. 1).

Neurological events were investigated regarding the presence of seizures, as well as, hearing and vision losses (Table 2). The presence of these baseline symptoms at least once after laboratory diagnosis of HIV-1 infection were admitted by 46.3% and 52.6%, of the mono and double infected individuals,

**Table 2 – Frequency of self-admitted neurological alterations among HIV-1 carriers and co-infected with JCV.**

Clinical symptoms	HIV-1 carriers (%)	JCV/HIV-1 coinfectd (%)	Total
With neurological symptoms	19 (46.3)	10 (52.6)	29 (48.3)
Without neurological symptoms	22 (53.7)	9 (47.4)	31 (51.7)
Total	41 (100)	19 (100)	60 (100)


$\chi^2 = 0.031; p = 0.860.$

respectively, but the difference between the frequencies was not significant ( $p=0.860$ ).

## Discussion

The information obtained from HIV-1 carriers were sufficient to establish the socio demographic variables and indicate possible trends of the infection. The age range observed (20–60 years) was in agreement with the epidemic in Brazil.<sup>18</sup> Similar results were seen in the USA with patients with non-Hodgkin's lymphoma, HIV-1 negative patients,<sup>19</sup> and healthy Koreans.<sup>20–23</sup> It is believed that JCV infection occurs early in life and therefore it is necessary to monitor the upsurge of disease among healthy individuals or with associated pathologies within different population groups as the virus prevalence is usually high.<sup>5,6,8,9</sup>

Infection was defined by the presence of two codifying genomic regions, VP1 and IG, which have been fairly correlated with seroepidemiological information among different populations.<sup>23–27</sup> The variation of prevalence rates are also commonly observed among HIV-1 carriers. The present work showed 32% prevalence of co-JCV/HIV-1 infection, which is similar to that reported by others among HIV-1 carriers with or without LPM.<sup>7,8,28–32</sup> The search for the virus in other biological



**Fig. 1 – Comparison of the mean values of (A) TCD4<sup>+</sup>, (B) TCD8<sup>+</sup> lymphocyte counts and (C) mean of HIV-1 viral load between HIV-1 and HIV-1/JCV infected patients.**

fluids including urine, plasma, liquor and cerebral tissue did not increase significantly the prevalence of JCV.<sup>30</sup>

Higher prevalence rates have been reported in Brazil in the CSF of HIV-1 carriers,<sup>33</sup> in Germany, in the blood,<sup>34</sup> in France, in brain tissue and blood<sup>35,36</sup> and in Italy, where very high and very low rates were described in the CSF of HIV-1 carriers.<sup>37,38</sup>

Co-infection with BKV is also a commonly reported finding among HIV-1 infected persons.<sup>8,39,40</sup> There was only one BKV/HIV-1 co-infected person in the examined group, in contrast to what has been reported by other. The presence of BKV has been previously reported in our geographical area in Afro-descendants living in the region of Rio Trombetas, Para,<sup>6</sup> among renal patients and the general population of urban centers.<sup>41</sup> The low prevalence of BKV indicates its low levels of dissemination.

HIV-1 infection was more frequent among males, which is consistent with the overall epidemic in Brazil.<sup>18</sup> The observation of sex and age distribution of HIV-1 carriers or not, were described by seven laboratories from England and Wales,<sup>7</sup> among aborigenes from Central Africa (pigmies and Bantus), in patients from general hospital in Australia, including immunocompromised persons,<sup>27</sup> and in Brazil<sup>40</sup> but there were no statistically significant differences of prevalence between sexes.

TCD4<sup>+</sup> and TCD8<sup>+</sup> lymphocyte counts, as well as, HIV-1 viral load were not different between mono- and double-infected subjects, which is a common finding,<sup>8,39,40,42</sup> suggesting that the frequency of JCV shedding may not be influenced or altered by the immunodeficiency process, but it remains a matter of controversy yet to be defined.

The investigation of possible disease among the examined group was approached by a self-admitted present or past common symptoms of major perception, including seizures, hearing and vision losses. There was no difference between the frequencies found among the mono- and double-infected groups. Although JCV is a major opportunistic infection of the CNS,<sup>11–15,43</sup> PML is not commonly reported in Brazil.<sup>40</sup> Either we are not performing the correct diagnosis of minor and major symptoms of PML, or reactivation of the virus, although a common feature, is without clinical consequences in HIV-1 infected persons in the North and Southeast regions of the country.

Until now, there have been few reports of JCV and BKV of either mono-infection or HIV-1 co-infection in Brazil. The results obtained, so far, show a large frequency of excretion of JCV and sometimes of BKV. The clinical consequences of the infections with both viruses should not be ignored and should deserve a deeper look into the actual clinical and epidemiological picture, including the improvement and dissemination of laboratory diagnosis of both infections.

## Funding

The present study was fully supported by grants from the STD/AIDS and Hepatitis Department, Ministry of Health/UNESCO, and the Brazilian National Council for Scientific and Technological Development (CNPq).

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgments

We thank the subjects and send them our best wishes.

## REFERENCES

1. Padgett BL, Walker DL, Zurhein GM, et al. Cultivation of Papova-like virus from human brain with progressive multifocal leucoencephalopathy. *Lancet*. 1971;1:1257–60.
2. Gardner SD, Field AM, Coleman DV, Hulme B. New human Papovavirus (BK) isolated from urine after renal transplantation. *Lancet*. 1971;1:1253–7.
3. White MK, Khalili K. Polyomaviruses and human cancer: molecular mechanisms underlying patterns of tumorigenesis. *Virology*. 2004;324:1–16.
4. Hale AD, Bartkeviute D, Dargeviute A, et al. Expression and antigenic characterization of the major capsid proteins of Human Polyomaviruses BK and JC in *Saccharomyces cerevisiae*. *J Virol Methods*. 2002;104:93–8.
5. Mcnees AL, White ZS, Zanwar P, et al. Specific and quantitative detection of Human Polyomaviruses BKV, JCV, and SV 40 by real time PCR. *J Clin Virol*. 2005;34:52–62.
6. Gee GV, Manley K, Atwood WJ. Derivation of a JC virus-resistant human glial cell line: implications for the identification of host cell factors that determine viral tropism. *Virology*. 2003;314:101–9.
7. Knowles WA, Sasnauskas K. Comparison of cell culture-grown JC virus (primary human fetal glial cells and the JCI cell line) and recombinant JCV VP1 as antigen for the detection of anti-JCV antibody by haemagglutination inhibition. *J Virol Methods*. 2003;109:47–54.
8. Behzad-Behbahani A, Klapper PE, Valley PJ, et al. Detection of BK virus and JC virus DNA in urine samples from immunocompromised (HIV-infected) and immunocompetent (HIV-non-infected) patients using polymerase chain reaction and microplate hybridization. *J Clin Virol*. 2004;29:224–9.
9. Ryschkewitsch C, Jensen P, Hou J, et al. Comparison of PCR-southern hybridization and quantitative real-time PCR for the detection of JC and BK viral nucleotide sequences in urine and cerebrospinal fluid. *J Virol Methods*. 2004;121:217–21.
10. Henmi C, Sawa H, Iwata H, et al. Establishment of an immunoscreening system using recombinant VP1 protein for the isolation of a monoclonal antibody that blocks JC virus infection. *Biochem Biophys Res Commun*. 2005;327:242–51.
11. Sacktor N. The epidemiology of human immunodeficiency virus-associated neurological disease in the era of highly active antiretroviral therapy. *J Neurovirol*. 2002;8:115–21.
12. Berger Jr. Progressive multifocal leucoencephalopathy in acquired immunodeficiency syndrome: explaining the high incidence and disproportionate immunosuppressive conditions. *J Neurovirol*. 2003;9:38–41.
13. D'Arminio-Monforte A, Cinque P, Mocroft A, et al. Changing incidence of central nervous system diseases in the EuroSIDA cohort. *Ann Neurol*. 2004;55:320–8.
14. Vidal JE, Penalva de Oliveira AC, Fink MC, et al. Aids-related progressive multifocal leucoencephalopathy: a retrospective study in a referral center in São Paulo, Brazil. *Rev Inst Med Trop*. 2008;50:209–12.



15. Pizaa F, Finkb MC, Nogueira GS, et al. JC virus-associated central nervous system diseases in HIV-infected patients in Brazil: clinical presentations, associated factors with mortality and outcome. *Braz J Infect Dis.* 2012;16:153-6.
16. Cayres-Vallinoto IMV, Vallinoto ACR, Azevedo VN, et al. Human JCV infections as a Bio-Anthropological marker of the formation of Brazilian Amazonian populations. *PLoS ONE.* 2012;7:E46523, <http://dx.doi.org/10.1371/journal.Pone.0046523>.
17. Ayres M, Ayres JR, Ayres DL, Santos AS. *BIOSTAT 5.0: Aplicações Estatísticas nas Áreas das Ciências Biológicas e Médicas.* Belém: Sociedade Civil Mamirauá/MCT-CNPq/Conservation International; 2007.
18. Ministério da Saúde do Brasil. Boletim Epidemiológico AIDS e DST. Ano III - no 1 - 01a à 26a semanas epidemiológicas - janeiro a junho de 2014, ISSN: 1517-1159. [www.aids.gov.br](http://www.aids.gov.br).
19. Engels EA, Rollison DE, Hartge P, et al. Antibodies to JC and BK viruses among persons with non-Hodgkin lymphoma. *Int J Cancer.* 2005;117:1013-9.
20. Agostini HT, Ryschkewitsch CF, Stoner GL. Genotype profile of human polyomavirus JC excreted in urine of immunocompetent individuals. *J Clin Microbiol.* 1996;34:159-64.
21. Chang H, Wang M, Tsai RT, et al. High incidence of JCV viremia in JC-seropositive older individuals. *J Neurovirol.* 2002;8:447-51.
22. Gu Z-Y, Li Q, Si Y-L, et al. Prevalence of BK virus and JC virus in peripheral blood leukocytes and normal arterial walls in healthy individuals in China. *J Med Virol.* 2003;70:600-5.
23. Jeong B-H, Lee K-H, Choi E-K, et al. Genotyping of the JC virus in urine samples of healthy Korean individuals. *J Med Virol.* 2004;72:281-9.
24. Stoner GL, Jobs DV, Cobo MF, et al. JC virus as a marker of human migration to the Americas. *Microbes Infect.* 2000;2:1905-11.
25. Chima SC, Ryschkewitsch CF, Stoner GL. Molecular epidemiology of human polyomavirus JC in the Biaka pygmies and Bantu of Central Africa. *Mem Inst Oswaldo Cruz.* 1998;93:615-23.
26. Sundsfjord A, Osci A, Rosenqvist H, et al. BK and JC viruses in patients with systemic lupus erythematosus: prevalent and persistent BK viremia, sequence stability of the viral regulatory regions, and nondetectable viremia. *J Infect Dis.* 1999;180:1-9.
27. Whitley DM, Arden KE, Mackay IM, et al. Simultaneous detection and differentiation of human polyomaviruses JC and BK by a rapid and sensitive PCR-Elaha assay and a survey of the JCV subtypes within an Australian population. *J Med Virol.* 2004;72:467-72.
28. Agostini HT, Yanagihara R, Davis V, et al. Asian genotypes of JC virus in native Americans and in a Pacific island population: markers of viral evolution and human migration. *Proc Natl Acad Sci USA.* 1997;94:14542-6.
29. Ferrante P, Caldarelli-Stefano R, Omodeo-Zorini E, et al. Comprehensive investigation of the presence of JC virus in AIDS patients with and without progressive multifocal leukoencephalopathy. *J Med Virol.* 1997;52: 235-42.
30. Pfister LA, Letvin NL, Koralknik IJ. JC virus regulatory region tandem repeats in plasma and central nervous system isolates correlate with poor clinical outcome in patients with progressive multifocal leukoencephalopathy. *J Virol.* 2001;75:5672-6.
31. Lednicky JA, Vilchez RA, Keitel WA, et al. Polyomavirus JCV excretion and genotype analysis in HIV-infected patients receiving highly active antiretroviral therapy. *AIDS.* 2003;17:801-7.
32. Mischitelli M, Fioriti D, Videtta M, et al. Investigation on the role of cell transcriptional factor SP1 and HIV-1 TAT protein in PML onset or development. *J Cell Physiol.* 2005;204:913-8.
33. Fink MCDS, Penalva de Oliveira AC, Milagres FAP, et al. JC vírus DNA in cerebrospinal fluid samples from Brazilian AIDS patients with focal brain lesions without mass effect. *J Infect.* 2006;52:30-6.
34. Weber F, Goldmann C, Krämer M, et al. Cellular and humoral immune response in progressive multifocal leukoencephalopathy. *Ann Neurol.* 2011;49:636-42.
35. Jochum W, Weber T, Frye S, et al. Detection of JC virus by anti-VP1 immunohistochemistry in brains with progressive multifocal leukoencephalopathy. *Acta Neuropathol.* 1997;94:226-31.
36. Dubois V, Moret H, Lafon M-E, et al. JC virus genotypes in France: molecular epidemiology and potential significance for progressive multifocal leukoencephalopathy. *J Infect Dis.* 2001;183:213-7.
37. De Luca A, Giancola ML, Ammassari A, et al. The effect of potent antiretroviral therapy and JC virus load in cerebrospinal fluid on clinical outcome of patients with AIDS-associated progressive multifocal leukoencephalopathy. *J Infect Dis.* 2000;182:1077-83.
38. Giri JA, Gregoresky J, Silguero P, et al. Polyoma virus JC DNA detection by polymerase chain reaction in CSF of HIV infected patients with suspected progressive multifocal leukoencephalopathy. *Am Clin Lab.* 2001;20:33-5.
39. Markowitz RB, Thompson HC, Mueller JF, et al. Incidence of BK virus and JC virus viremia in human immunodeficiency virus-infected and uninfected subjects. *J Infect Dis.* 1993;167:13-20.
40. Nali LH, Centrone CC, Urbano PR, et al. High prevalence of the simultaneous excretion of polyomaviruses JC and BK in the urine of HIV-infected patients without neurological symptoms in São Paulo, Brazil. *Rev Inst Med Trop Sao Paulo.* 2012;54:201-5.
41. Pires EP, Bernardino-Vallinoto CV, Alves DM, et al. Prevalence of infection by JC and BK Polyomaviruses in kidney transplant recipients and patients with chronic renal disease. *Transpl Infect Dis.* 2011;13:633-7.
42. Matos A, Duque V, Beato S, et al. Characterization of JC human polyomavirus infection in a Portuguese population. *J Med Virol.* 2010;82:494-504.
43. Holman RC, Török TJ, Belay ED, et al. Progressive multifocal leukoencephalopathy in the United States, 1979-1994: increased mortality associated with HIV infection. *Neuroepidemiology.* 1998;17:303-9.