

The Elusive Renal Cell Carcinoma: Reversal Imaging of Arterial Phase to Improve Acuity

Erich K. Lang, Karl Zhang, Quan Nguyen, Daniel Thorner, Ernest Rudman

Department Radiology and Urology, SUNY Downstate Medical School, Brooklyn, New York, NY, USA

On routine physical examination the patient's physician noted microscopic hematuria. The finding was reconfirmed by 2 Dipsticks over an interval of 4 months. The patient was a known diabetic, controlled by diet. Otherwise the patient was asymptomatic, without significant past medical history at the time of this work-up, the 47 year old Caucasian male appeared to be in good general health. Laboratory data showed Hb of 15.1gm/dL, HCT 45%, RBC 4.8 million/uL, WBC 6200, Neu 62%, BUN 18 mg/dL, Creatinine 1.1 mg/dl, GRF 94 mL/min, A/G ratio 1.4, Glu 128 mg/dl, K 4.2 mmol/L, Na 145 MMOL/L Cl 108 mmol/L Urine analysis, spec grav 1018, 3-5 RBC/hpf, no WBC or bacteria on hpf, no casts, urine culture negative x 2. A KUB (Flat plate of abdomen) showed no opaque calculi nor other abnormalities. Cystoscopy and blue light cystoscopy revealed no abnormalities.

An enhanced 4 phase MDCT was performed. The pre-enhancement phase was entirely unremarkable; no parenchymal lesions were detected. Following administration of 100 ml non-ionic contrast medium at a flow rate of 5 mL/sec, the 12 second delayed arterial phase Ct demonstrated a relatively poorly enhancing 1.6 cm mass at the cortico-medullary junction (Figure-1), the lesion is much better shown on reversal image. Both the parenchymal phase CT (50 second delay) and the excretory phase CT (4 minutes delay Figure-2) demonstrate a non-enhancing 16 mm mass at the cortico-medullary junction (Figure-3).

In the light of a clinical history of diabetes and microscopic hematuria, the non-enhancing hypovascular mass seen on parenchymal and ex-

cretory phase CTs in the medulla might have been written off as Medullary Necrosis (With characteristic CT findings of a negative pre-enhancement phase CT, but a non-enhancing lesion shown on parenchymal and excretory phase; an early avascular necrosis) (1). However, the reversal image of the arterial phase CT clearly shows an enhancing lesion, though somewhat hypovascular for a RCC. The hypo-density on parenchymal and excretory phase CT reflects the characteristic "wash-out" phenomenon of RCCs in these phases. The tumor having no tubules is less dense than adjacent normal parenchyma. The correct diagnosis was made, and a laparoscopic resection carried out.



Figure 1 - Massive gas in the bladder, dissecting in the submucosal layer and extending into the space of Retzius.



Figure 2 - lateral extension of the gas-dissection in the pre-vesical space.




Figure 3 - gas dissecting into the ureter.

REFERENCES

1. Lang EK, Macchia RJ, Thomas R, Davis R, Ruiz-Deya G, Watson RA, et al.: Multiphasic helical CT diagnosis of early medullary and papillary necrosis. J Endourol. 2004; 18: 49-56.
2. Israel GM, Bosniak MA: How I do it: evaluating renal masses. Radiology. 2005; 236: 441-50.

Correspondence address:

Dr. Erich K. Lang
Departments of Urology and Radiology
SUNY, Downstate Health Science Center
455 Lenox Road
Brooklyn, NY, 11203, USA
E-mail: erich.lang@downstate.edu