

Genetic gain prediction of the third recurrent selection cycle in a popcorn population

Fabrcio Santana Santos¹, Antonio Teixeira do Amaral Jnior^{1*}, Silvrio de Paiva Freitas Jnior¹, Ramon Macedo Rangel¹, Carlos Alberto Scapim² and Freddy Mora²

¹Laborat3rio de Melhoramento Gen3tico Vegetal, Centro de Ci4ncias e Tecnologias Agropecu3rias, Universidade Estadual do Norte Fluminense, Av. Alberto Lamego, 2000, 28013-602, Parque Calif3rnia, Campos dos Goytacazes, Rio de Janeiro, Brazil.

²Departamento de Agronomia, Universidade Estadual de Maring3, Maring3, Paran3, Brazil. *Author for correspondence. E-mail: amaraljr@uenf.br

ABSTRACT. Although Brazil has reduced its importation of popcorn grain in recent years, the availability of improved varieties is still rare. In order to improve a popcorn population and to estimate the prediction of genetic gains, the third cycle of intrapopulation recurrent selection was carried out. One hundred ninety two half-sib families were obtained from the second cycle of the UNB-2U population and tested in a randomized complete block design with two replications within sets in two locations. The presence of statistical significance for sets for grain yield, poorly husked ear and broken plants proved the efficiency and the necessity for the use of the hierarchical statistical model by sets. Except for the poorly husked ear, there was significance for families within sets for the other evaluated traits, showing that there was sufficient genetic variability for exploitation in future cycles. Although the heritability for popping expansion was higher than for grain yield, the predicted genetic gain for the latter was superior to popping expansion because of the higher magnitude of the genetic standard deviation for grain yield. It is expected that in the fifth recurrent cycle, the first popcorn variety with yellow grains may be released for Rio de Janeiro State.

Key words: *Zea mays*, popcorn breeding, genetic gains, yield, popping expansion.

RESUMO. Predi3o de ganhos gen3ticos para o terceiro ciclo de sele3o recorrente em uma popula3o de milho pipoca. Embora o Brasil tenha reduzido a importa3o de gr3os de milho pipoca, a disponibilidade de variedade melhorada ainda 3 rara no pa3s. Com o objetivo de melhorar a popula3o UNB-2U de milho pipoca e estimar a predi3o de ganhos gen3ticos, implementou-se o terceiro ciclo de sele3o recorrente intrapopulacional. Cento e noventa e duas fam3lias de meios-irm3os foram obtidas e avaliadas em delineamento de blocos casualizados, com duas repeti3es, dentro de "sets" em dois locais. A presen3a de signific3ncia para "sets", para produ3o de gr3os, espigas mal empalhadas e plantas quebradas, comprovou a efici4ncia do uso do modelo estat3stico hierarquizado por "sets". Com exce3o das espigas mal empalhadas, houve signific3ncia para fam3lias dentro de "sets" para as demais caracter3sticas, revelando que h3 suficiente variabilidade gen3tica a ser explorada em ciclos futuros. Ainda que a herdabilidade para capacidade de expans3o (PE) tenha sido superior 3 da produ3o de gr3os (GY), o ganho predito para GY foi superior a PE, em decorr4ncia da maior estimativa da magnitude do desvio-padr3o gen3tico para GY. Espera-se que, no quinto ciclo de sele3o recorrente, seja poss3vel a recomenda3o da primeira variedade de milho pipoca com gr3os amarelos para o Estado do Rio de Janeiro.

Palavras-chave: *Zea mays*, melhoramento de milho pipoca, ganhos gen3ticos, produ3o de gr3os, capacidade de expans3o.

Introduction

In the last five years, the Brazilian popcorn market has changed, especially regarding the partnership between producers and packers, which has been reflected in the importation of the product. Until the beginning of the 2003-2004 growing season, national popcorn consumption was

approximately 80,000 tons, of which 75% corresponded to imported grains, mainly from Argentina (Galv3o *et al.*, 2000). In the 2004-2005 growing season, according to information from the packing companies, there was a marked reduction in importation of the product, to 20,000 tons, as a result of the large-scale use of the IAC-112 modified

simple hybrid, developed by the Campinas Agronomic Institute (Sawazaki, 2001).

However, the hybrid popcorn market in Brazil is still incipient, in spite of the importance of the crop that is valued at approximately 1.15 billion dollars annually in the United States of America (Pacheco *et al.*, 1998; Galvão *et al.*, 2000). Yet, obtaining an improved variety from a composite is also rare in Brazil, although the release of varieties derived from composites has been a viable alternative (Vilela *et al.*, 2008). As an example, the variety BRS ANGELA was derived from recurrent selection cycles of the CMS-43 composite, from Embrapa/CNPMS (Pacheco *et al.*, 2000).

The main reason for the dependency on imported popcorn is that breeding programs in the country are rare. As examples can be cited the breeding programs developed in Viçosa and Sete Lagoas (Minas Gerais State), Maringá and Londrina (Paraná State), and Piracicaba and Campinas (São Paulo State). A popcorn breeding program at Campos dos Goytacazes has also been established, aiming at the development of varieties with adaptation for the state of Rio de Janeiro.

The initial program stages involved the assessment of the potential of the UNB-2U population, especially in terms of genetic variance, in order to identify more efficient breeding strategies to produce improved varieties (Pereira and Amaral Júnior, 2001; Vilela *et al.*, 2008). Thus, the first recurrent selection cycle based on full-sib family selection resulted in predicted genetic gains of 4.69% (yield) and 10.39% (popping expansion) (Daros *et al.*, 2002). In the second cycle, based on S_1 families, the predicted genetic gains were 17.8% for popping expansion and 26.95% for yield (Daros *et al.*, 2004).

Thus, the main objective of this work was to estimate the genetic gain prediction of a third cycle of recurrent selection, and to evaluate the potential of the improved population for future releasing as a new popcorn variety.

Material and methods

One hundred and ninety-two half-sib families were obtained from the C_2 cycle. The field trials were established in December 2004 in Campos dos Goytacazes (21°45'15" S, 41°19'28" W) and in January 2005 in Itaocara (21°40'09" S, 42°04'34" W), in Rio de Janeiro State, Brazil. These environments are contrasting in terms of regions (about 110 kilometers apart), soil types and altitudes.

A completely randomized block design was used,

with six sets, two replications within sets, and each set consisting of 32 families and eight controls. Each family was cultivated in a 5.0 m row with inter-row spacing of 1.0 m and 0.2 m spacing between plants. Initially, fifty seeds were sown per row. One month after sowing, the plants were thinned to 25 plants per row. Soil preparation, fertilizer application and cultural practices were in conformity with the recommendations by Fancelli and Dourado Neto (2000).

The following traits were evaluated: a) popping expansion (PE) – ratio of expanded volume in relation to weight in 30 grams, in mL g⁻¹; b) grain yield (GY) – weight of grain in kg ha⁻¹; c) poorly husked ears (PHE) – number of ears not completely covered by husk; and d) broken plants (BP) – number of plants with stalk broken below the upper ear at harvest.

Popping expansion was assessed in a popcorn machine developed by Embrapa/Instrumentação Agropecuária, incorporating an electronic resistor and thermostat. Two grain samples were obtained and evaluated from each plot.

An analysis of variance was carried out on the measured traits, based on the completely randomized block design with two replications within sets in two environments, according to the genetic-statistical model proposed by Hallauer and Miranda Filho (1981), ie:

$$Y_{ijkl} = \mu + E_i + S_j + ES_{ij} + R/ES_{ijk} + F/S_{jl} + EF/S_{jil} + \xi_{ijkl}$$

where:

μ is the general mean;

E_i is the fixed effect of the i^{th} environment;

S_j is the random effect of j^{th} set;

ES_{ij} is the random effect of environment-by-set interaction;

R/ES_{ijk} is the random effect of the k^{th} replication within the environment-by-set interaction;

F/S_{jl} is the random effect of the l^{th} family within the set;

EF/S_{jil} is the random effect of environment-by-family interaction within the set; and

ξ_{ijkl} is the experimental error.

Each set was analyzed separated, and then the results from the six sets were grouped for each environment. Finally, the results for all sets were combined for both environments. All sources of variation, with the exception of environment, were assumed to be random. The GENES Program (Cruz, 2006) was used for data analysis.

The genetic parameters, phenotypic ($\hat{\sigma}_p^2$) and genotypic ($\hat{\sigma}_g^2$) variance and heritability on the

family mean basis (\hat{h}^2) were estimated as follows:

$$\hat{\sigma}_f^2 = \frac{MSF}{re}; \hat{\sigma}_g^2 = \frac{MSF - MSE}{re}; \text{ and } \hat{h}^2 = \frac{\hat{\sigma}_g^2}{\hat{\sigma}_f^2};$$

where: MSF and MSE are, respectively, the family mean square and error mean square; r and e are, respectively, the numbers of replications and environments (Falconer, 1987).

The estimates of genetic gain predictions were obtained by the expression $GS = p \times DP_g \times h \times i$, where p is the parentage control (1 for recombination of remnant seeds of the half-sib families), DP_g is the additive standard deviation among the test units, h is the heritability square root, and i is the selection intensity. The value of i was obtained by the ratio between z (standard curve height at the truncation point) and c (percentage of selected individuals) (Cruz, 2005).

Results and discussion

The coefficients of variation (CV) were considered high for all traits, except popping expansion (Table 1). In the second cycle of recurrent selection of this population (UNB-2U), high values of CV were also observed for these traits (Daros *et al.*, 2004). In the CMS-42 and CMS-43 popcorn populations, high magnitudes of CV for the traits percentage of lodged and broken plants were reported by Pacheco *et al.* (1998). The high values of the coefficients of variation are also in accordance with Coimbra (2000) and Santos *et al.* (2007).

Table 1. Mean squares, general mean, maximum and minimum family mean values and coefficients of variation of the four evaluated traits in the half-sib families, Campos dos Goytacazes and Itaocara, 2005.

Source	DF	Mean Square			
		PE (mL g ⁻¹)	GY (kg ha ⁻¹)	PHE (number)	BP (number)
E	1	176.07**	4447927.81**	168.25**	42.23ns
S	5	15.34ns	786828.61**	5.29*	48.74**
ES	5	17.32ns	89453.59ns	6.94**	54.96**
R/ES	12	24.46**	692591.00**	2.36ns	51.00**
F/S	186	23.72**	253805.04**	2.03ns	16.16*
EF/S	186	9.23ns	165124.99**	2.18ns	13.09ns
Error	372	8.60	132697.00	2.22	13.09ns
Maximum		39.36	3120.00	12.00	12.66
Average		23.24	1135.74	1.76	5.49
Minimum		10.33	380.00	0.00	0.00
CV (%)		12.62	32.07	84.76	64.72

**,*Significant by F-test at 1 and 5%, respectively. E is the environment effect; S is the random effect as result of set; ES is the random effect of environment-by-set interaction; R/ES is the random effect of the replication within the environment-by-set interaction; F/S is the random effect of the family within the set; EF/S is the random effect of environment-by-family interaction within the set. DF: degrees of freedom. CV: coefficient of variation.

The analysis of variance results indicate environment effects for the evaluated traits, except

for broken plants, showing that the environments differed sufficiently for most of the evaluated traits (Table 1).

Although the effect of environment was significant for GY, there was no significance for the EF/S variation source, showing a tendency to maintain the relative performance of the families in the different environments (Table 1).

Only the trait PE did not show statistical difference for sets, indicating the efficiency and effectiveness for the use of block designs divided into sets. If set effect were not included, it would probably produce variations that could result in loss of experimental accuracy (Table 1).

The source of variation F/S was significant by the F test, for all the evaluated traits, except for PHE (Table 1). This shows there is genetic variability for exploitation in future cycles, enabling progress with selection. Furthermore, it indicated that the recurrent selection objective of improving the population mean maintaining the genetic variability was attained in the third selection cycle.

The selective potential of the UNB-2U population for the main traits, PE and GY, was indicated by the difference in the mean of the selected families compared to the mean of the original population (Table 2). Additionally, the great discrepancy in the mean of the lower and upper limit for the main traits, PE and GY (Table 2), is another indication of the genetic variability in the population UNB-2U that is an essential condition to obtaining superior segregates. Similar results were obtained by Pacheco *et al.* (1998), who also based on the upper and lower limits for PE and GY to predict genetic gains in the CMS-42 and CMS-43 popcorn populations in the second recurrent selection cycle.

Table 2. Estimates of the means of the original population (\bar{X}_o), the selected families (\bar{X}_s), the controls (\bar{X}_r), lower limit (LL) and upper limit (UL) of the evaluated population.

Estimates	Traits			
	PE (mL g ⁻¹)	GY (kg ha ⁻¹)	PHE (number)	BP (number)
\bar{X}_o	23.30	1135.58	1.76	5.51
\bar{X}_s	25.74	1361.92	1.45	4.79
\bar{X}_r	19.72	1885.40	1.54	6.17
LL	10.30	380.00	0.00	0.00
UL	39.67	3120.00	12.00	33.00

It is timely to recognize that the upper limit can rise throughout the selection cycle, which means that a maximum value detected in a given cycle can be lower than the maximum for the following cycle, and thus successively with the increase of the frequency of favorable alleles followed by

recombination and generation of advanced cycle populations.

Based on genetic component estimates (Table 3), the heritability estimates of the PE trait are in line with results obtained by Sawazaki (1996), which ranged from 60 to 90% in popcorn half-sib families. Similar estimates for this trait in CMS-43 popcorn population in the second selection cycle among and within half-sib families were obtained by Pacheco *et al.* (1998).

In the present investigation, PE heritability was superior to that for GY, with values of 68.40 and 50.17%, respectively. Such results are in agreement with previous study conducted by Pereira and Amaral Júnior (2001) using design I (Comstock and Robinson, 1948) in the same base population. These results are also in line with other investigations (Pacheco *et al.*, 1998; Freitas Júnior *et al.*, 2006; Scapim *et al.*, 2006a; Rangel *et al.*, 2007; Rangel *et al.*, 2008). Consequently, as mentioned by Pacheco *et al.* (1998), greater genetic gains were expected for PE than for GY. However, in the present study, the gain for GY was superior to that predicted for PE because of the estimate of the magnitude of the GY standard deviation, which was much greater than for PE.

Table 3. Phenotypic variance ($\hat{\sigma}_p^2$), genotypic variance ($\hat{\sigma}_g^2$), family mean heritability (\hat{h}_x^2 %) and genetic gain (GG%) estimated for the evaluated traits.

Genetic Parameters	Traits			
	PE	GY	PHE	BP
$\hat{\sigma}_p^2$	6.28	68875.00	0.54	4.15
$\hat{\sigma}_g^2$	4.29	34559.52	0.00	0.75
\hat{h}_x^2 %	68.42	50.18	0.00	18.12
GG%	11.40	17.97	0.00	10.39

The genetic gain for the PE trait was 11.40% for the third selection cycle. Based on the initial mean, such gain is equivalent to an increase of 1.84 units year⁻¹. In relation to GY, the predicted gain was of 17.97% in comparison to the original population. Such gain is equivalent to an increase of 134.28 kg ha⁻¹ year⁻¹ (Table 3).

When the means obtained for PE and GY in the three first cycles were compared with the predicted mean for the third cycle, it was observed that there were increases in PE and GY as a result of the favorable allele concentration (Table 4). Linear regression analysis showed the evolution of gains in the successive selection cycles (Figures 1 and 2). It was observed that the selection strategy to obtain C₃ using half-sib families resulted in satisfactory gains for popping expansion and grain yield, which increased to desirable magnitudes. The use of S₁

families in a previous cycle (Daros *et al.*, 2004) did not compromise the genetic gains once after selection; the selected families are recombined, overcoming the inbreeding depression and recovering the vigor of the population (Scapim *et al.*, 2006b).

Table 4. Obtained means in the C₀, C₁, C₂ and predicted mean in the C₃ cycles^{1/} of recurrent selection in UNB-2U population for the popping expansion (PE) and grain yield (GY) in Campos dos Goytacazes and Itaocara.

Cycle	PE (mL g ⁻¹)	GY (kg ha ⁻¹)
C0	20.28	1960.00
C1	21.68	1990.00
C2	23.44	2025.00
C3	25.11	2080.00

^{1/} C₀ = mass selection; C₁ = full-sib recurrent selection; C₂ = S₁ families recurrent selection; and C₃ = half-sib recurrent selection.




Figure 1. Obtained Means in the three first cycles (0, 1 and 2) and predicted mean in the third cycle (3) of recurrent selection in UNB-2U population for the trait PE (mL g⁻¹).




Figure 2. Obtained Means in the three first cycles (0, 1 and 2) and predicted mean in the third cycle (3) of recurrent selection in UNB-2U population for the trait GY (kg ha⁻¹).

Judging from the progress of the first three cycles, it is expected that with two more cycles of recurrent selection, the resulted population will present desirable popping expansion and grain yield magnitudes that will propitiate the future release of the first bred variety of popcorn with yellow grains in Rio de Janeiro State. Also, it is expected that, in addition to the magnitude of the desirable traits, the improved population should also retain enough

genotypic variability to justify the continuity of recurrent selection procedures with the correspondent genetic gains.

Conclusion

The significance of the source of variation F/S for most of the traits showed that there is genetic variability for exploitation in future cycles, enabling progress with selection.

Although the heritability for PE was superior to GY, the predicted gain for the latter was superior to the former, resulting from the greater estimates of the genetic standard deviation for GY.

It is expected that after the fifth recurrent selection cycle, it will be possible to release the first popcorn variety with yellow grains for Rio de Janeiro State.

Acknowledgement

We thank Faperj and CNPq for funding this study.

References

- COIMBRA, R.R. *Seleção entre famílias de meios-irmãos da população DFT1-Ribeirão de milho pipoca*. 2000. Dissertação (Mestrado em Genética e Melhoramento)-Universidade Federal de Viçosa, Viçosa, 2000.
- COMSTOCK, R.E.; ROBINSON, H.F. The components of genetic variance in populations of biparental progenies and their use in estimating the average degree of dominance. *Biometrics*, Washington, D.C., v. 4, n. 3, p. 254-266, 1948.
- CRUZ, C.D. *Programa genes: versão windows: aplicativo computacional em genética e estatística*. Viçosa: UFV, 2006.
- CRUZ, C.D. *Princípios de genética quantitativa*. Viçosa: UFV, 2005.
- DAROS, M. *et al.* Genetic gain for grain yield and popping expansion in full-sib recurrent selection in popcorn. *Crop Breed. Appl. Biotechnol.*, Londrina, v. 2, n. 3, p. 339-344, 2002.
- DAROS, M. *et al.* Recurrent selection in inbred popcorn families. *Sci. Agric.*, Piracicaba, v. 61, n. 6, p. 609-614, 2004.
- FALCONER, D.S. *Introdução à genética quantitativa*. 1. ed. Viçosa: UFV, 1987.
- FANCELLI, A.L.; DOURADO NETO, D. *Produção de milho*. Guaíba: Agropecuária, 2000.
- FREITAS JÚNIOR *et al.* Capacidade combinatória de milho pipoca por meio de dialelo circulante. *Pesq. Agropecu. Bras.*, Brasília, v. 41, n. 11, 1599-1607, 2006.
- GALVÃO, J.C.C. *et al.* Comportamento de híbridos de milho pipoca em Coimbra, Minas Gerais. *Rev. Ceres*, Viçosa, v. 47, n. 270, p. 201-218, 2000.
- HALLAUER, A.R.; MIRANDA FILHO, J.B. *Quantitative genetics in maize breeding*. Ames: Iowa State University Press, 1981.
- PACHECO, C.A.P. *et al.* Estimativas de parâmetros genéticos nas populações CMS-42 e CMS-43 de milho pipoca. *Pesq. Agropecu. Bras.*, Brasília, v. 33, n. 12, p. 1995-2001, 1998.
- PACHECO, C.A.P. *et al.* BRS ANGELA: variedade de milho pipoca. Sete Lagoas: Embrapa-CNPMS, 2000. (Comunicado técnico, 27).
- PEREIRA, M.G.; AMARAL JÚNIOR, A.T. Estimation of genetic components in popcorn based on the nested design. *Crop Breed. Appl. Biotechnol.*, Viçosa, v. 1, n. 1, p. 3-10, 2001.
- RANGEL, R.M. *et al.* Prediction of popcorn hybrid and composite means. *Crop Breed. Appl. Biotechnol.*, Viçosa, v. 7, n. 3, p. 287-295, 2007.
- RANGEL, R.M. *et al.* Genetic parameters in parents and hybrids of circulant diallel in popcorn. *Gen. Mol. Res.*, Ribeirão Preto, v. 7, n. 4, p. 1020-1030, 2008.
- SANTOS, F.S. *et al.* Predição de ganhos genéticos por índices de seleção na população de milho-pipoca UNB-2U sob seleção recorrente. *Bragantia*, Campinas, v. 66, n. 3, p. 391-398, 2007.
- SAWAZAKI, E. *Parâmetros genéticos em milho pipoca*. Tese (Doutorado em Genética e Melhoramento de Plantas)-Universidade de São Paulo, Piracicaba, 1996.
- SAWAZAKI, E. A cultura do milho pipoca no Brasil. *O Agrônômico*, Campinas, v. 53, n. 2, p. 11-13, 2001.
- SCAPIM, C.A. *et al.* Combining ability of white grain popcorn populations. *Crop Breed. Appl. Biotechnol.*, Viçosa, v. 6, n. 1, p. 136-143, 2006a.
- SCAPIM, C.A. *et al.* Componentes genéticos de médias e depressão por endogamia em populações de milho pipoca. *Cienc. Rural*, Santa Maria, v. 36, n. 1, p. 36-41, 2006b.
- VILELA, F.O. *et al.* Effect of recurrent selection on the genetic variability of the UNB-2U popcorn population using RAPD markers. *Acta Sci. Agron.*, Maringá, v. 30, n. 1, p. 25-30, 2008.

Received on April 29, 2008.

Accepted on May 28, 2008.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.