



Brazilian Journal of
OTORHINOLARYNGOLOGY

www.bjorl.org



ORIGINAL ARTICLE

Polymorphism in GRHL2 gene may contribute to noise-induced hearing loss susceptibility: a meta-analysis[☆]

Xin Li ^a, Zhengping Zhu ^a, Wei Li ^b, Li Wei ^c, Baocheng Zhao ^d, Zheng Hao ^{d,*}



^a Nanjing Municipal Center for Disease Control and Prevention, Department of HIV/AIDS/STI Prevention and Control, Jiangsu, China

^b Southeast University, School of Public Health, Department of Epidemiology and Health Statistics, Key Laboratory of Environmental Medicine Engineering, Jiangsu, China

^c Nanjing Municipal Center for Disease Control and Prevention, Environmental Health Division, Jiangsu, China

^d Nanjing Zhongyangmen Community Health Service Center, Kang'ai Hospital, Center of Diagnosis and Treatment for Developmental Dysplasia of the Hip, Jiangsu, China

Received 9 July 2018; accepted 2 January 2019

Available online 23 February 2019

KEYWORDS

Noise-induced hearing loss;
Grainyhead like 2;
Molecular epidemiology;
Polymorphism;
Meta-analysis

Abstract

Instruction: Noise-induced hearing loss is a leading occupational disease caused by gene-environment interaction. The Grainy Like 2, GRHL2, is a candidate gene. In this regard, many studies have evaluated the association between GRHL2 and noise-induced hearing loss, although the results are ambiguous and conflicting.

Objective: The purpose of this study was to identify a precise estimation of the association between rs3735715 polymorphism in GRHL2 gene and susceptibility of noise-induced hearing loss.

Methods: A comprehensive search was performed to collect data up to July 8, 2018. Finally, 4 eligible articles were included in this meta-analysis comprising 2410 subjects. The pooled odds ratios with 95% confidence intervals were used to evaluate the strength of the association.

Results: Significant association was found in the overall population in the dominant model (GA/AA vs. GG, odds ratio = 0.707, 95% confidence interval = 0.594–0.841) and allele model (G allele vs. A allele, odds ratio = 1.189, 95% confidence interval = 1.062–1.333). When stratified by source of the subjects, we also found association between rs3735715 and noise-induced hearing loss risk in the dominant model (GA/AA vs. GG, odds ratio = 0.634, 95% confidence interval = 0.514–0.783) and allele model (G allele vs. A allele, odds ratio = 1.206, 95% confidence interval = 1.054–1.379).

[☆] Please cite this article as: Li X, Zhu Z, Li W, Wei L, Zhao B, Hao Z. Polymorphism in GRHL2 gene may contribute to noise-induced hearing loss susceptibility: a meta-analysis. Braz J Otorhinolaryngol. 2020;86:370–5.

* Corresponding author.

E-mail: charliehao@msn.com (Z. Hao).

Peer Review under the responsibility of Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial.

Conclusion: Rs3735715 polymorphism in GRHL2 gene may influence the susceptibility of noise-induced hearing loss. Additional large, well-designed and functional studies are needed to confirm this association in different populations.

© 2019 Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial. Published by Elsevier Editora Ltda. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

PALAVRAS-CHAVE

Perda auditiva
induzida por ruído;
Grainyhead like 2;
Epidemiologia
molecular;
Polimorfismo;
Metanálise

Polimorfismo no gene GRHL2 pode contribuir para a suscetibilidade à perda auditiva induzida por ruído: uma metanálise

Resumo

Introdução: Perda auditiva induzida por ruído é uma das principais doenças ocupacionais causadas pela interação gene-ambiente. O *Grainy Like 2*, ou *GRHL2* é um gene que tem sido considerado como candidato. Nesse sentido, muitos estudos avaliaram a associação entre o *GRHL2* e perda auditiva induzida por ruído, embora os resultados sejam ambíguos e conflitantes. **Objetivo:** Identificar uma estimativa precisa da associação entre o polimorfismo rs3735715 no gene *GRHL2* e a suscetibilidade à perda auditiva induzida por ruído.

Método: Uma pesquisa abrangente foi feita para coletar dados até 8 de julho de 2018. No fim, quatro artigos elegíveis foram incluídos nesta metanálise, abrangeram 2.410 indivíduos. As *odds ratios* agrupadas com intervalos de confiança de 95% foram usadas para avaliar a força da associação.

Resultados: Uma associação significante foi encontrada na população geral no modelo de dominância (GA/AA vs. GG, *odds ratio* = 0,707, intervalo de confiança 95% = 0,594-0,841) e modelo de alelo (alelo G vs. alelo A; *odds ratio* = 1,189, intervalo de confiança 95% = 1,062 a 1,333). Quando estratificados pelo local de trabalho dos indivíduos, também encontramos associação entre rs3735715 e risco de perda auditiva induzida por ruído no modelo de dominância (GA/AA vs. GG, *odds ratio* = 0,634, intervalo de confiança 95% = 0,514 ± 0,783) e modelo de alelo (alelo G vs. alelo A; *odds ratio* = 1,206, intervalo de confiança 95% = 1,054- 1,379).

Conclusão: O polimorfismo Rs3735715 no gene *GRHL2* pode influenciar a suscetibilidade à perda auditiva induzida por ruído. Estudos adicionais, amplos, bem desenhados e funcionais são necessários para confirmar essa associação em diferentes populações.

© 2019 Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Noise-induced hearing loss (NIHL) is one of the worldwide leading occupational disease especially in the developing countries and the second most frequent form of sensorineural hearing deficit after Age-Related Hearing Impairment (ARHI).¹ It is a complex hearing impairment induced by a combination of genetic and environment factors.² Recently, accumulating epidemiological evidence indicated that noise, organic solvents, heat, heavy metals, vibrations, smoking, drinking, high blood pressure and cholesterol levels are responsible environmental factors.²⁻⁸ Additionally, it was demonstrated that genetics contribute to the incidence of NIHL, deduced from animal experiments.^{9,10} Some of these studies used heterozygote or homozygote knockout mice model and confirmed that otocadherin 23 (cdh23) gene,¹¹ glutamate peroxidase 1 (gpx1) gene,¹² plasma membrane Ca²⁺-ATPase isoform 2 (pmca2) gene,¹³ and heat shock factor (hsf1) gene¹⁴ might associate with NIHL risk. In human, several association studies have demonstrated that CDH23 gene,^{15,16} human 8-oxOG DNA glycosylase 1 (hOGG1) gene,¹⁷ catalase (CAT) gene,¹⁸ heat shock protein 70 (HSP70) gene,¹⁹ potassium voltage-gated channel, Isk-related family,

member 1 (KCNE1) gene and potassium voltage-gated channel, KQT-like subfamily, member 4 (KCNQ4) gene²⁰ might be involved in the susceptibility to NIHL.

The *Grainy Like 2* (GRHL2) is a transcription factor which is associated with the composition of the organ of Corti.²¹ Grainyhead-like transcription factor family includes three members (GRHL1-GRHL3) regulating epithelial adhesion.²² GRHL2 is highly expressed in cochlear duct lining cells and plays an important role in the epithelial cell maintenance and embryonic development.²³ GRHL2 gene knockout mice were embryonically lethal.²² Recently, two research teams confirmed that GRHL2 gene might influence the susceptibility to ARHI and progressive autosomal dominant hearing loss (DFNA28).^{22,24} Up to now, promising but contradictory data showed that the GRHL2 gene might be responsible for the development of NIHL. GRHL2 gene is located on chromosome 8q22.3, including 15 introns and 16 exons. Currently, several candidate gene studies have focused on whether the GRHL2 gene is associated with NIHL risk while the results remain conflicting rather than conclusion. Yang et al. genotyped the potentially functional polymorphism rs3735715 and got a significant association.²⁵ Xu et al. confirmed this finding in another population.²⁶ But Li

Table 1 Basic information of the 4 studies in this meta-analysis.

Year	First author	Country	Ethnicity	Source of the subjects	Cases	Controls
2018	Yang	China	Asian	Steel factory	340	339
2016	Yang	China	Asian	Steel factory	283	281
2013	Li	China	Asian	Chemical fiber company	340	356
2016	Xu	China	Asian	Steel factory	236	235

et al. did not find any association between rs3735715 and NIHL susceptibility.²⁷ In this study, we performed a meta-analysis to estimate the overall association.

Methods


Search strategy and date extraction

We searched all literatures from PubMed, CNKI, Wang Fang, Web of Science and Springer databases, using the keyword (*GRHL2* or "Grainyhead Like 2" or rs3735715) and (NIHL or "Noise-Induced Hearing Loss"). The search time was not limited. The last search update was done on July 8th, 2018. Articles investigating GRHL2 and NIHL before July 8th, 2018 were all included in this meta-analysis. The first study was in 2013. Four literatures were included in our meta-analysis according to the following criteria: (1) was a case-control study; (2) was a study about the GRHL2 polymorphism and NIHL susceptibility; (3) with usable data for allele frequency; (4) was a paper written in English or Chinese. Two of the authors extracted the available data independently according to the criteria mentioned above. We extracted the information including publication year, the name of the first author, country, ethnicity, source of the subjects, and genotype distributions of the GRHL2 rs3735715 polymorphisms among cases and controls. The controversies were discussed within our research team, and we reached a consensus eventually.

Statistical methods

Summary Odds Ratios (ORs) with 95% Confidence Intervals (CIs) determined by Z-test were used to assess the strength of the association. If the *p*-value was less than 0.05, the association was considered significant. Stratified analysis was performed by source of the subjects. We used *Q* test to assess the between-study heterogeneity. The heterogeneity was considered to be significant if the *p*-value was less than 0.10. χ^2 statistic ($\chi^2 = 100\% \times (Q - df)/Q$) was also used to quantify heterogeneity. χ^2 greater than 50% indicated heterogeneity among studies. The fixed-effects model and random-effects model were used to pool the data appropriately.²⁸ The fixed-effects model was used when there was no heterogeneity existed. It assumes that all the studies are sampled from the populations with same effect size. The fixed-effects model makes an adjustment to study weights according to the in-study variance. The random-effects model based on the Dersimonian and Laird method was more suitable when the heterogeneity existed; otherwise the two methods provided the same results.

To test for publication bias in this meta-analysis, we performed both Egger's and Begg's test.²⁹ The publication bias was assessed by funnel plot and the linear regression asymmetry test.

**Figure 1** Flow diagram of study inclusion/exclusion.**Table 2** Genotype distributions of the GRHL2 rs3735715 polymorphisms among cases and controls.

Year	First author	Cases			Controls		
		GG	GA	AA	GG	GA	AA
2018	Yang	115	157	68	85	181	73
2016	Yang	94	126	63	67	154	60
2013	Li	122	161	57	119	162	75
2016	Xu	80	104	52	56	130	49

All analyses were performed using Stata software version 8.2 (Stata Corporation, College Station, TX, USA).

Results

The characteristics of the included studies

A total of 25 relevant studies were identified through database screening. Seven were excluded for duplicated records. After a detailed evaluation of the full-text of the 18 studies, 14 were excluded: two concerned about other polymorphisms, 4 reviews, 1 not for human, 6 not about NIHL and one lack of genetic distribution data. 4 articles were eventually included in the analysis. The flow of studies through this meta-analysis was shown in Fig. 1. Characteristics of the 4 studies and details of the genotype distributions were shown in Tables 1 and 2.




Figure 2 Begg's funnel plot for publication bias test using the dominant model (GA/AA vs GG).

The demographics of the subjects of the 4 studies in this meta-analysis

The average of age in cases was 40.7 ± 8.4 , 40.5 ± 8.1 , 39.3 ± 5.8 , and 40.4 ± 8.3 for Yang's (2018), Yang's (2016), Li's (2013), and Xu's (2016) study, respectively. The average of age in controls was 40.0 ± 8.4 , 39.8 ± 8.1 , 39.8 ± 5.8 , and 39.5 ± 8.2 for Yang's (2018), Yang's (2016), Li's (2013), and Xu's (2016) study, respectively. The hearing threshold level in cases was 51.0 ± 9.0 , 37.6 ± 11.7 , and 51.4 ± 8.8 for Yang's (2016), Li's (2013), and Xu's (2016) study, respectively. The hearing threshold level in controls was 11.7 ± 10.7 , 14.2 ± 3.9 , and 9.3 ± 9.1 for Yang's (2016), Li's (2013), and Xu's (2016) study, respectively. The demographics of the subjects of the 4 studies in this meta-analysis were detailed in Table 3.

Overall population

Overall, our meta-analysis showed a significant association between rs3735715 and NIHL risk in both dominant model and allele model. For dominant model (GA/AA vs. GG) OR = 0.707, 95% CI = 0.594–0.841. For allele model (G allele vs. A allele) OR = 1.189, 95% CI = 1.062–1.333 (Table 4).

Subgroup analysis by source of the subjects

Stratification by source of the subjects identified a significant association in steel factory population between GRHL2 rs3735715 polymorphism and NIHL risk. Among the subjects chosen from steel factory, significant association was found in the dominant model (GA/AA vs. GG) OR = 0.634, 95% CI = 0.514–0.783, and allele model (G allele vs. A allele) OR = 1.206, 95% CI = 1.054–1.379 (Table 4). We did not find any significant association between the GRHL2 rs3735715 polymorphism and NIHL risk in the chemical fiber company workers.

Heterogeneity and publication bias

The meta-analysis was performed using a fixed-effects model based on the Mantel-Haenszel method because the between-study heterogeneity was not found (Table 4).

Publication bias is always a concern in a meta-analysis. As shown in Fig. 2, there was no obvious asymmetry in the shape of the funnel. We also performed the Egger's test to evaluate the funnel plot symmetry. The results showed

no evidence of the publication bias ($t = -1.92$, $p = 0.194$ for GRHL2 rs3735715 GA/GG vs. GG).

Discussion

In the present study, the GRHL2 rs3735715G > A polymorphism was found to be associated with NIHL risk in the dominant model (GA/AA vs. GG) and allele model (G allele vs. A allele) including 1199 NIHL cases and 1211 controls. Comparing with the GG genotype, GA/AA genotype showed a decrease risk of NIHL. The G allele showed increasing risk comparing with A allele. In the subgroup analysis, the association was also found in steel factor workers, but not among chemical fiber company members. These results suggest that the potentially functional polymorphism rs3735715 may affect susceptibility to NIHL. As far as we know, this is the first comprehensive meta-analysis to estimate the association between the GRHL2 polymorphism and NIHL risk.

GRHL2, also known as Brother of Mammalian grainyhead (BOM) and Transcription Factor Cellular Promoter 2-Like 3 (TFCP2L3) is a member of GRHL transcription factor family which controls the development of multicellular epithelia by regulating cell junction formation and proliferation genes.^{30,31} The junction proteins and ion channels play a critical role in the otic epithelial cells in inner-ear development and homeostasis maintenance. In zebrafish model, the mutant of GRHL2 shows inner-ear defects. Injecting wide-type human GRHL2 mRNA could rescue the defects.³² GRHL2 might influence the susceptibility to ARHI and DFNA28.^{22,24} Just like ARHI and DFNA28, NIHL is one type of sensory impairment. Although they are not completely the same type hearing loss, some features are totally correspond, such as the hearing threshold in high frequency are most affected, and the sensorineural and progressive nature.²⁴

In 2009, Konings et al.³³ performed a large-scale association study in two independent noise-exposed populations to identify susceptibility genes for NIHL which did not find an association between GRHL2 and NIHL risk. The difference may be multifactorial, such as ethnic difference, different inclusion criteria, lifestyle, genetic and environmental factors. Our results about the rs3735715 polymorphism agreed with Yang's,^{25,34} and Xu's²⁶ studies while differed from Li's²⁷ analysis. The reasons may be in Li's study,²⁷ they chose workers from the chemical fiber company while others using steel factory subjects.^{25,26,34} Additionally, the definition of NIHL cases differed too. In Li's study,²⁷ the workers with hearing threshold worse than 25 dB in high frequency were defined as NIHL. Otherwise in the other researches, the subjects with hearing threshold worse than 40 dB in high frequency were defined as NIHL.^{25,26,34} The negative result of Li's study about rs3735715 may also due to another reason. The effect of this polymorphism on the risk of NIHL might be too small to be detectable with the small sample size. The reason remains unclear, and studies using the same ethnic population are still needed to confirm these findings in the future.

An advantage of this meta-analysis was that numbers of subjects were pooled together from each independent study, which increased the statistical power significantly. Second, the quality of articles included in this analysis was satisfactory according to a consistent selection criterion. Third, based on this meta-analysis, functional study of rs3735715 in GRHL2 might be conducted to replicate these observations. There were some limitations of our meta-analysis. First, in the present study potential confounding factors (such as age, gender, exposure level, exposure time etc.) were

Table 3 The demographics of the subjects of the 4 studies in this meta-analysis.

First author (year)	Age (years) (mean \pm SD)		Gender		Hearing threshold level (dB) (mean \pm SD)		Time of exposition to the noise (years) (mean \pm SD)	
	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls
Yang (2018)	40.7 \pm 8.4	40.0 \pm 8.4	326 male 17 female	326 male 17 female	NA	NA	NA	NA
Yang (2016)	40.5 \pm 8.1	39.8 \pm 8.1	274 male 12 female	274 male 12 female	51.0 \pm 9.0	11.7 \pm 10.7	18.9 \pm 9.1	18.3 \pm 8.8
Li (2013)	39.3 \pm 5.8	39.8 \pm 5.8	306 male 34 female	317 male 39 female	37.6 \pm 11.7	14.2 \pm 3.9	17.0 \pm 6.9	17.0 \pm 7.0
Xu (2016)	40.4 \pm 8.3	39.5 \pm 8.2	239 male 0 female	239 male 0 female	51.4 \pm 8.8	9.3 \pm 9.1	18.7 \pm 9.2	18.7 \pm 9.2

NA, data not available.

Table 4 Meta-analysis of the GRHL2 rs3735715 polymorphism and NIHL risk.

Population	Comparison	Test of association	<i>p</i> ^a	Test of heterogeneity	
				<i>p</i> ^b	<i>I</i> ² (%)
Overall	GA/AA vs. GG	0.707 (0.594–0.841)	<0.001	0.343	10
	AA vs. GA/GG	0.928 (0.761–1.130)	0.455	0.584	0
	G allele vs. A allele	1.189 (1.062–1.333)	0.003	0.977	0
Source of the subjects steel factory	GA/AA vs. GG	0.634 (0.514–0.783)	<0.001	0.964	0
	AA vs. GA/GG	1.001 (0.794–1.261)	0.993	0.815	0
	G allele vs. A allele	1.206 (1.054–1.379)	0.006	0.971	0
Chemical fiber company	GA/AA vs. GG	0.897 (0.656–1.226)	0.496	–	–
	AA vs. GA/GG	0.755 (0.515–1.106)	0.149	–	–
	G allele vs. A allele	1.149 (0.928–1.421)	0.202	–	–

^a *p*-value determined by Z test.^b *p*-value determined by *Q*-test.

not adjusted. Second, we only chose four papers written in English or Chinese. Although we estimated the publication bias using Egger's and Begg's test, we cannot ignore the possibility of bias. Third, we did not perform further evaluation of the potential interactions. The gene-gene or gene-environment interactions may modulate NIHL risk.³⁵

Conclusion

In conclusion, this meta-analysis found an association between the GRHL2 rs3735715 polymorphism and NIHL risk, suggesting GRHL2 might influence the susceptibility of NIHL. In the future, more extensive studies are still required to confirm these findings in different ethnic populations.

Funding

This work was supported by the Humanities and Social Sciences of Ministry of Education Planning Fund of China (grant no. 16YJA840014).

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

This work was supported by the Humanities and Social Sciences of Ministry of Education Planning Fund of China (grant no. 16YJA840014).

References

- Konings A, Van Laer L, Van Camp G. Genetic studies on noise-induced hearing loss: a review. *Ear Hear*. 2009;30:151–9.
- Sliwinska-Kowalska M. Organic solvent exposure and hearing loss. *Occup Environ Med*. 2008;65:222–3.
- Upile T, Sipaul F, Jerjes W, Singh S, Nouraei SA, El MM, et al. The acute effects of alcohol on auditory thresholds. *BMC Ear Nose Throat Disord*. 2007;7:4.
- Taylor W, Pearson J, Mair A, Burns W. Study of noise and hearing in jute weaving. *J Acoust Soc Am*. 1965;38:113–20.
- Campo P, Lataye R. Noise and solvent, alcohol and solvent: two dangerous interactions on auditory function. *Noise Health*. 2000;3:49–57.
- Ward WD. Endogenous factors related to susceptibility to damage from noise. *Occup Med*. 1995;10:561–75.
- Mehrparvar AH, Mirmohammadi SJ, Hashemi SH, Davari MH, Mostaghaci M, Mollasadeghi A, et al. Concurrent effect of noise exposure and smoking on extended high-frequency pure-tone thresholds. *Int J Audiol*. 2015;54:301–7.
- Fechter LD, Chen GD, Rao D. Chemical asphyxiants and noise. *Noise Health*. 2002;4:49–61.

9. White CH, Ohmen JD, Sheth S, Zebboudj AF, Mchugh RK, Hoffman LF, et al. Genome-wide screening for genetic loci associated with noise-induced hearing loss. *Mamm Genome.* 2009;20:207–13.
10. Davis RR, Newlander JK, Ling X, Cortopassi GA, Krieg EF, Erway LC. Genetic basis for susceptibility to noise-induced hearing loss in mice. *Hear Res.* 2001;155:82–90.
11. Holme RH, Steel KP. Progressive hearing loss and increased susceptibility to noise-induced hearing loss in mice carrying a Cdh23 but not a Myo7a mutation. *J Assoc Res Otolaryngol.* 2004;5:66–79.
12. Ohlemiller KK, McFadden SL, Ding DL, Lear PM, Ho YS. Targeted mutation of the gene for cellular glutathione peroxidase (Gpx1) increases noise-induced hearing loss in mice. *J Assoc Res Otolaryngol.* 2000;1:243–54.
13. Kozel PJ, Davis RR, Krieg EF, Shull GE, Erway LC. Deficiency in plasma membrane calcium ATPase isoform 2 increases susceptibility to noise-induced hearing loss in mice. *Hear Res.* 2002;164:231–9.
14. Fairfield DA, Lomax MI, Dootz GA, Chen S, Galecki AT, Benjamin IJ, et al. Heat shock factor 1-deficient mice exhibit decreased recovery of hearing following noise overstimulation. *J Neurosci Res.* 2005;81:589–96.
15. Sliwinska-Kowalska M, Noben-Trauth K, Pawelczyk M, Kowalski TJ. Single nucleotide polymorphisms in the cadherin 23 (Cdh23) gene in polish workers exposed to industrial noise. *Am J Hum Biol.* 2008;20:481–3.
16. Kowalski TJ, Pawelczyk M, Rajkowska E, Dudarewicz A, Sliwinska-Kowalska M. Genetic variants of Cdh23 associated with noise-induced hearing loss. *Otol Neurotol.* 2014;35:358–65.
17. Shen H, Cao J, Hong Z, Liu K, Shi J, Ding L, et al. A functional Ser326Cys polymorphism in Hogg1 is associated with noise-induced hearing loss in a Chinese population. *PLOS ONE.* 2014;9:e89662.
18. Konings A, Van Laer L, Pawelczyk M, Carlsson PI, Bondeson ML, Rajkowska E, et al. Association between variations in cat and noise-induced hearing loss in two independent noise-exposed populations. *Hum Mol Genet.* 2007;16:1872–83.
19. Konings A, Van Laer L, Michel S, Pawelczyk M, Carlsson PI, Bondeson ML, et al. Variations in Hsp70 genes associated with noise-induced hearing loss in two independent populations. *Eur J Hum Genet.* 2009;17:329–35.
20. Pawelczyk M, Van Laer L, Fransen E, Rajkowska E, Konings A, Carlsson PI, et al. Analysis of gene polymorphisms associated with K⁺ ion circulation in the inner ear of patients susceptible and resistant to noise-induced hearing loss. *Ann Hum Genet.* 2009;73:411–21.
21. Lu J, Cheng X, Li Y, Zeng L, Zhao Y. Evaluation of individual susceptibility to noise-induced hearing loss in textile workers in China. *Arch Environ Occup Health.* 2005;60:287–94.
22. Han Y, Mu Y, Li X, Xu P, Tong J, Liu Z, et al. Grhl2 deficiency impairs otic development and hearing ability in a zebrafish model of the progressive dominant hearing loss Dfna28. *Hum Mol Genet.* 2011;20:3213–26.
23. Peters LM, Anderson DW, Griffith AJ, Grundfast KM, San AT, Madeo AC, et al. Mutation of a transcription factor, Tfcp2L3, causes progressive autosomal dominant hearing loss, Dfna28. *Hum Mol Genet.* 2002;11:2877–85.
24. Van Laer L, Van Eyken E, Fransen E, Huyghe JR, Topsakal V, Hendrickx JJ, et al. The Grainyhead Like 2 Gene (Grhl2), Alias Tfcp2L3, is associated with age-related hearing impairment. *Hum Mol Genet.* 2008;17:159–69.
25. Yang QY, Xu XR, Jiao J, He LH, Yu SF, Gu GZ, et al. Association between grainyhead-like 2 gene polymorphisms and noise-induced hearing loss. *Beijing Da Xue Xue Bao Yi Xue Ban.* 2016;48:409–13.
26. Xu X, Yang Q, Jiao J, He L, Yu S, Wang J, et al. Genetic variation in Pou4F3 and Grhl2 associated with noise-induced hearing loss in Chinese population: a case-control study. *Int J Environ Res Public Health.* 2016;13.
27. Li X, Huo X, Liu K, Li X, Wang M, Chu H, et al. Association between genetic variations in Grhl2 and noise-induced hearing loss in Chinese high intensity noise exposed workers: a case-control analysis. *Ind Health.* 2013;51:612–21.
28. Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst.* 1959;22:719–48.
29. Egger M, Davey SG, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ.* 1997;315:629–34.
30. Chen W, Xiao LZ, Oh JE, Shin KH, Kim RH, Jiang M, et al. Grainyhead-Like 2 (Grhl2) inhibits keratinocyte differentiation through epigenetic mechanism. *Cell Death Dis.* 2012;3:e450.
31. Petrof G, Nanda A, Howden J, Takeichi T, McMillan JR, Aristodemou S, et al. Mutations in Grhl2 result in an autosomal-recessive ectodermal dysplasia syndrome. *Am J Hum Genet.* 2014;95:308–14.
32. Rifat Y, Parekh V, Wilanowski T, Hislop NR, Auden A, Ting SB, et al. Regional neural tube closure defined by the grainy head-like transcription factors. *Dev Biol.* 2010;345:237–45.
33. Konings A, Van Laer L, Wiktorek-Smagur A, Rajkowska E, Pawelczyk M, Carlsson PI, et al. Candidate gene association study for noise-induced hearing loss in two independent noise-exposed populations. *Ann Hum Genet.* 2009;73:215–24.
34. Yang Q, Wang J, Xu X, He L, Yu S, Chen G, et al. Relationship between interaction of three kinds of hereditary deafness gene and susceptibility of high-frequency hearing loss. *Chinese J Ind Med.* 2018;83–6.
35. Marchini J, Donnelly P, Cardon LR. Genome-wide strategies for detecting multiple loci that influence complex diseases. *Nat Genet.* 2005;37:413–7.