

SHORT COMMUNICATION

Notes on the hematology of free-living *Phrynops geoffroanus*  
(Testudines: Chelidae) in polluted rivers of Southeastern Brazil

Bruno O. Ferronato <sup>1, 2, 5</sup>; Alexander Genoy-Puerto <sup>2</sup>; Carlos I. Piña <sup>3</sup>; Franco L. Souza <sup>4</sup>;  
Luciano M. Verdade <sup>1</sup> & Eliana R. Matushima <sup>2</sup>

<sup>1</sup> Laboratório de Ecologia Animal, Departamento de Ciências Biológicas, ESALQ, Universidade de São Paulo, Caixa Postal 09, Piracicaba, SP 13418-900, Brasil. E-mail: brunoferronato@hotmail.com; lmv@esalq.usp.br

<sup>2</sup> Laboratório de Patologia Comparada de Animais Silvestres, Departamento de Patologia, Universidade de São Paulo, São Paulo, SP 05508-270, Brasil. E-mail: alexandergenoy@usp.br; ermatush@usp.br

<sup>3</sup> CICYTTP-CONICET. Proyecto Yacaré. PO Box 3105, Diamante, Entre Ríos, Argentina. E-mail: cidcarlos@infoaire.com.ar

<sup>4</sup> Centro de Ciências Biológicas e da Saúde, Departamento de Ciências Biológicas, Universidade Federal do Mato Grosso do Sul, Campo Grande, MS 79070-900, Brasil. E-mail: flsouza@nin.ufms.br

<sup>5</sup> Corresponding author.

---

ABSTRACT. *Phrynops geoffroanus* (Schweigger, 1812) is the freshwater turtle with the widest geographical distribution in South America. During 2006, physical examination and hematological evaluation were performed on free-ranging turtles from two polluted rivers, the Piracicaba River (n = 51) and its tributary Piracicamirim (n = 42), in southeastern Brazil. Red blood cell and thrombocyte counts, mean corpuscular volume and mean corpuscular hemoglobin levels differed in turtles from the two water courses. Although free-ranging turtles showed ectoparasites and boat propeller lesions, animals apparently had no signs of clinical disease. In spite of our results, further monitoring of the demography and health status of *Phrynops geoffroanus* in anthropogenically altered environments is recommended.

KEY WORDS. Hemogram; lesion; pollution; turtle.

---

Geoffroy's side-necked turtle, *Phrynops geoffroanus* (Schweigger, 1812) is the freshwater turtle with the widest geographical distribution in South America, ranging from the Colombian Amazon to southern Brazil and northern Argentina, living in rivers, streams and lakes (PRITCHARD & TREBBAU 1984, ERNST & BARBOUR 1989). Recently, the species has been found inhabiting polluted urban rivers in Brazil (SOUZA & ABE 2000, BRITES & RANTIN 2004), exhibiting values of population density and biomass among the highest recorded for neotropical freshwater turtles (SOUZA & ABE 2000).

Hematologic values are important for monitoring health of free-ranging turtle populations (BOLTEN & BJORNDALE 1992, METIN *et al.* 2006). Most of hematological studies in free-ranging and captive turtles are reported as baseline values in endangered species (MARKS & CITINO 1990, BOLTEN & BJORNDALE 1992, CHRISTOPHER *et al.* 1999, BRENNER *et al.* 2002, DEEM *et al.* 2006, METIN *et al.* 2006, KARIZOE *et al.* 2007) or in diseases related to anthropogenic causes (JACOBSON *et al.* 1991, WORK *et al.* 2001, CHRISTOPHER *et al.* 2003). Investigation on animals living in disturbed habitats is crucial, since man-made alterations in the environment, such as habitat loss, degradation or pollution, are the main reasons for population declines of reptiles (GIB-

BONS *et al.* 2000, RODRIGUES 2005) and can negatively influence animal health (CHRISTOPHER *et al.* 1999). Anemia is one of the disorders that can be detected in species living in polluted habitats (KELLER *et al.* 2004, BELSKII *et al.* 2005). Our goal in this investigation was to describe hematological range values of *P. geoffroanus* from two disturbed sites in Southeastern Brazil.

From May through October 2006, free-living turtles were captured in the Piracicaba River (PIRA) and its tributary, Piracicamirim (PISCA), located in Piracicaba River basin (PRB), a highly developed basin (12.400 km<sup>2</sup>), with industries and a high human population density (KRUSCHE *et al.* 1997, MARTINELLI *et al.* 1999). PRB is located in the State of São Paulo, Southeastern Brazil. The main industrial activities include textile, paper and pulp, sugar and ethanol, metallurgic, food crops, tanning, chemical and fuel refineries (CETESB 2001). Sugar cane and citrus are the main agricultural products (KRUSCHE *et al.* 1997). The densest human population and industries are concentrated in the central part of the basin. PISCA is considered the most polluted tributary of PIRA. The main sources of pollution are non-treated domestic sewage and fertilizers used in sugar cane crops (OMETO *et al.* 2004). Its micro basin (133 km<sup>2</sup>) encompasses portions of three counties: Piracicaba, Rio das Pedras, and Saltinho (OMETO *et al.* 2000).

Turtles were captured in PIRA by active search using a motor boat and hand dip nets, in a region known as Monte Alegre (22°41'75"S, 47°33'58"W), located in the central area of the basin, in the suburban area of the city of Piracicaba. We captured turtles in PISCA close to its largest urbanized area on the University of São Paulo campus (22°42'51"S, 47°37'36"W). We used four gill nets (nylon; mesh size 3-5 cm; 1.5-2 m deep; 15 m long) extended perpendicularly between the river banks (SOUZA & ABE 2000) and checked every three hours. The animals were captured under Ibama license (Proc. 02010.000005/05-61).

Immediately after capture, turtles were transported in plastic containers to the laboratory in Piracicaba. Three ml of blood were collected from the external jugular vein (ROGERS & BOOTH 2004). Immediately after blood collection, two blood smears were prepared for each turtle for estimation of thrombocytes and white cell differential counts. The blood smears were air-dried and placed in cardboard holders. The remaining blood was placed in lithium-heparin tubes, chilled (4°C), and processed at the Veterinary School of Medicine of the University of São Paulo (FMVZ/USP) within 5-10 h of collection.

Packed cell volume was determined with microhematocrit capillary tubes filled from the lithium-heparin tubes and centrifuged at 10,000 rpm for five minutes in a microcentrifuge (MADER 2000). Plasma protein concentration was measured with a refractometer (ANDERSON *et al.* 1997). Hemoglobin concentration was determined by the cyanmethemoglobin method (CAMPBELL 1996). White and red blood cell counts were determined using the standardized recount system in a modified Neubauer chamber, with Natt-Herrick solution as diluent (FRYE 1991). White cell differential and thrombocytes were manually determined from modified May-Grunwald-Giemsa stained blood smears (ROSENFELD 1947). A hundred leukocytes were examined on each slide and were categorized as lymphocytes, eosinophils, heterophils, basophils and monocytes (SYPEK & BORYSENKO 1988, BRENNER *et al.* 2002, DEEM *et al.* 2006). Thrombocytes were counted every 1,000 red blood cells on the blood smear. Mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) were calculated from the packed cell volume, hemoglobin concentration, and red blood cell count (CAMPBELL 1996).

After blood sampling, animals were sexed based on their external morphological characteristics (ERNST & BARBOUR 1989, SOUZA & ABE 2000), measured at the straight-line carapace length, weighed and had a physical examination performed. The health status of each turtle was based on general body condition (DEEM *et al.* 2006), searching for clinical signs as sunken eyes, cachexia, lethargy and emaciation.

Descriptive statistics (mean, standard deviation, minimum and maximum) were performed. Data were tested for normality. Analysis of variance (General linear model) and analysis of mean (Anom) (Minitab for Windows 15) were performed for comparison between PIRA and PISCA turtles' blood values, where sex, sampling site and its interaction were considered as grouping factors.

A total of 51 adult turtles (29 males and 22 females) from PIRA and 42 adults (24 males and 18 females) from PISCA were captured. Mean straight-line carapace lengths were  $288 \pm 27$  mm (range: 236-348 mm) and  $291 \pm 32$  mm (range: 221-365 mm) from PIRA and PISCA respectively. Body mass was  $2280 \pm 675$  g (range: 1340-4100 g) and  $2444 \pm 814$  g (range: 950-5200 g) from PIRA and PISCA respectively. Fourteen animals (six males and eight females) from PIRA and eight (two males and six females) from PISCA had carapace injuries, probably caused by boat propeller strikes as we have noticed during the field work in PIRA. The traumatic lesions were completely healed in 12 of these turtles from PIRA and seven from PISCA. Ectoparasites (leeches) were observed on the skin and carapace in most of the animals from PIRA and PISCA, but this information was not registered systematically. Wild turtles were not cachectic or lethargic. They were in good body condition and apparently had no clinical signs of disease.

Results of *P. geoffroanus* hematology are listed in table I. Sex and "site by sex" interaction did not influence turtle blood values from both PIRA and PISCA ( $p > 0.05$ , Tab. I). Statistical differences occurred in 4 of 13 parameters analysed, due to location ( $p < 0.05$ , Tab. I). PIRA turtles had higher red blood cell and thrombocytes counts than PISCA turtles. However, PISCA turtles had higher mean corpuscular volume (MCV) and mean corpuscular hemoglobin (MCH). There were no differences in white blood cell and white cell differential counts between sites. Lymphocytes and heterophils were the most common white cells in PISCA turtles. For PIRA animals, lymphocytes were the most common in white cell differential counts (Tab. I).

The data presented herein provide hematological values for free-ranging *P. geoffroanus*. Even though free-ranging turtles were captured in disturbed and polluted sites, animals showed no detectable evidence of clinical disease. The shape of the lesions on the cracked and fractured carapaces is consistent with damage caused by motorboat strikes and show no signs of being caused by pathological agents, as recorded for other freshwater turtles (GARNER *et al.* 1997, KNOTKOVA *et al.* 2005).

Comparing the hematological values from PIRA turtles and its tributary PISCA, there was no variation due to sex or "site by sex" interaction. Rather, the site where animals were captured was responsible for the variation. Turtles from PIRA had a higher red blood cell count when compared with turtles from PISCA, what could be related to the river hydrodynamics. PIRA is larger and has a stronger water flow (MARTINELLI *et al.* 1999). Consequently, turtles in this site may have to swim more, thus consuming more oxygen, what could result in a higher number of erythrocytes. Although red blood cell count was higher in PIRA turtles, animals did not show differences in hemoglobin concentration and packed cell volume. PISCA turtles had a higher MCH and MCV than PIRA turtles. However, these variables are calculated values (CAMPBELL 1996), and probably are being influenced by the red blood cell count.

Table I. Hematology of *P. geoffroanus* from anthropogenically affected environments, southeastern Brazil. Piracicaba River (PIRA) and Piracicamirim stream (PISCA). Mean, standard deviation and range values in parentheses.

Measure	Females PIRA (n = 22)	Males PIRA (n = 29)	Females PISCA (n = 18)	Males PISCA (n = 24)
Packed cell volume%	21.1 ± 5.4 (13-32)	22 ± 4.5 (14-30)	21.1 ± 4.6 (15-32)	22.6 ± 5.6 (12-32)
Plasma protein concentration g dl <sup>-1</sup>	5.91 ± 1.29 (3.8-8.4)	6.2 ± 1.47 (3.4-9.2)	5.96 ± 1.18 (3.9-7.8)	6.81 ± 2.15 (3.6-11.2)
Hemoglobin g/dl	5.1 ± 1.38 (3.2-7.8)	5.68 ± 1.61 (2.8-9.5)	4.86 ± 1.06 (3.3-7.1)	5.38 ± 1.95 (2.9-11.2)
Red blood cell (x 10 <sup>3</sup> /μl) *	484 ± 178 (210-870) <sup>a</sup>	482 ± 140 (250-860) <sup>a</sup>	364 ± 146 (100-660) <sup>b</sup>	384 ± 99.7 (240-660) <sup>b</sup>
White blood cell (x 10 <sup>3</sup> /μl)	5.9 ± 3.5 (1.5-15)	5.7 ± 3.1 (1.5-12)	6.7 ± 3.0 (1.5-15)	6.6 ± 3.3 (3-16)
Absolute heterophils (x 10 <sup>3</sup> /μl)	1.9 ± 2.0 (0.2-7.5)	1.9 ± 1.5 (0.3-7.2)	2.5 ± 2.1 (0.5-8.7)	2.0 ± 1.8 (0.4-7)
Absolute eosinophils (x 10 <sup>3</sup> /μl)	0.89 ± 0.47 (0.2-1.8)	0.99 ± 0.78 (0.1-2.9)	0.93 ± 0.64 (0.2-2.5)	1.1 ± 0.8 (0.1-3.3)
Absolute basophils (x 10 <sup>3</sup> /μl)	0.43 ± 0.3 (0.07-1.0)	0.37 ± 0.3 (0.05-1.2)	0.48 ± 0.41 (0.05-1.8)	0.47 ± 0.34 (0-1.1)
Absolute lymphocytes (x 10 <sup>3</sup> /μl)	2.4 ± 1.4 (0.8-6.2)	2.3 ± 1.7 (0.7-7)	2.43 ± 1.5 (0.5-5.6)	2.8 ± 1.6 (0.6-6.4)
Absolute monocytes (x 10 <sup>3</sup> /μl)	0.17 ± 0.22 (0-0.7)	0.14 ± 0.11 (0-0.4)	0.26 ± 0.3 (0-1.2)	0.20 ± 0.29 (0-1.4)
Thrombocytes (x 10 <sup>3</sup> /μl) *	25 ± 28 (7.2-148) <sup>a</sup>	19 ± 11 (7.8-59) <sup>a</sup>	12 ± 7.8 (1.7-28) <sup>b</sup>	12.6 ± 7.4 (2.1-29) <sup>b</sup>
MCV fl *	469 ± 122(241.3-628.5) <sup>a</sup>	487 ± 141 (215.1-756.7) <sup>a</sup>	650 ± 246 (418-1500) <sup>b</sup>	601.2 ± 140.7 (400-1000) <sup>b</sup>
MCH pg *	112.9 ± 31.3(74.6-214) <sup>a</sup>	122.4 ± 37.6(68.2-243.5) <sup>a</sup>	149.9 ± 55.7 (90.9-330) <sup>b</sup>	144.7 ± 58.3 (93.9-329.4) <sup>b</sup>
MCHC g/dl	24.7 ± 6.1 (19.6-39)	26.7 ± 9.6 (13.3-51.1)	23 ± 1.9 (19.5-26.6)	23.6 ± 4.7 (17.8-41.4)

\*Values statistically different (p &lt; 0.05) between turtles from Piracicaba River and Piracicamirim stream.

Published data of freshwater turtles suffering from necrotic shell diseases (GARNER *et al.* 1997, KNOTKOVA *et al.* 2005) showed higher values of white blood cells, lymphocytes, heterophils and basophils than those obtained for the free-ranging *P. geoffroanus* in this study. However it is difficult to decide whether the hematological values obtained for *P. geoffroanus*, in this investigation, are low or normal, since there are no reference range values for free-ranging *P. geoffroanus* inhabiting unpolluted rivers. Future studies should focus on the hematology of *P. geoffroanus* from unpolluted environments for comparison with the data presented here.

Despite the fact that the turtles from PISCA and PIRA were apparently healthy, subclinical infections can occur. We recommend monitoring the health status of these turtles, with the addition of other techniques such as biochemistry analysis, necropsies, histological examination, and population size monitoring on a long term basis.

## ACKNOWLEDGMENTS

We thank S. Rossi and T.C. Sanches from Laboratório de Patologia Comparada de Animais Silvestres (FMVZ/USP), for their assistance in teaching the hematological procedures and techniques. We also thank T.S. Marques, I. Guardia and A.L.B. Longo from Laboratório de Ecologia Animal (ESALQ/USP) for helping in turtles' capture. We thank Valentine Lance, from San Diego State University, for a critical reading and an English review of this manuscript. The *P. geoffroanus*' ecology investigation was sponsored by FAPESP Research Grant (Proc. 2005/00210-9) and CNPq (Proc. 300087/2005-5). BOF was sponsored by CAPES scholarship.

## LITERATURE CITED

- ANDERSON, N.L.; R.F. WACK & R. HATCHER. 1997. Hematology and clinical chemistry reference ranges for clinically normal, captive new guinea snapping turtle (*Elseya novaeguineae*) and the effects of temperature, sex, and sample type. **Journal of Zoo and Wildlife Medicine** 28: 394-403.
- BELSKII, E.A.; N.V. LUGAS'KOVA & A.A. KARIDOVA. 2005. Reproductive parameters of adult birds and morphophysiological characteristics of chicks in the pied flycatcher (*Ficedula hypoleuca* Pall.) in technogenically polluted habitats. **Russian Journal of Ecology** 36: 329-335.
- BOLTEN, A.B. & K.A. BJØRNDAL. 1992. Blood profiles for a wild population of green turtles (*Chelonia mydas*) in the southern Bahamas: size-specific and sex-specific relationships. **Journal of Wildlife Diseases** 28: 407-413.
- BRENNER, D.; G. LEWBART; M. STEBBINS & D.W. HERMAN. 2002. Health survey of wild and captive bog turtles (*Clemmys muhlenbergii*) in North Carolina and Virginia. **Journal of Zoo and Wildlife Medicine** 33: 311-316.
- BRITES, V.L.C. & F.T. RANTIN. 2004. The influence of agricultural and urban contamination on leech infestation of freshwater turtles, *Phrynops geoffroanus*, taken from two areas of the Uberabinha river. **Environmental Monitoring and Assessment** 96: 273-281.

- CAMPBELL, T.W. 1996. Clinical Pathology, p. 248-257. *In*: D.R. MADER (Ed.). **Reptile medicine and surgery**. Philadelphia, W.B. Saunders, 1242p.
- CETESB. 2001. **Relatório de qualidade das águas interiores do estado de São Paulo**. São Paulo, Companhia de Tecnologia de Saneamento Ambiental, Série Relatórios Ano 2000, 227p.
- CHRISTOPHER, M.M.; K.H. BERRY; I.R. WALLIS; K.A. NAGY; B.T. HENEN & C.C. PETERSON. 1999. Reference intervals and physiologic alterations in hematologic and biochemical values of free-ranging desert tortoises in the Mojave Desert. **Journal of Wildlife Diseases** 35: 212-238.
- CHRISTOPHER, M.M.; K.H. BERRY; B.T. HENEN & K.A. NAGY. 2003. Clinical disease and laboratory abnormalities in free-ranging desert tortoises in California (1990-1995). **Journal of Wildlife Diseases** 39: 35-56.
- DEEM, S.L.; E.S. DIERENFELD; G.P. SOUNGUET; A.R. ALLEMAN; C. CRAY; R.H. POPPENG; T.M. NORTON & W.B. KARESH. 2006. Blood values in free-ranging nesting leatherback sea turtles (*Dermochelys coriacea*) on the coast of the Republic of Gabon. **Journal of Zoo and Wildlife Medicine** 37: 464-471.
- ERNST, C.H. & R.W. BARBOUR. 1989. **Turtles of the world**. Washington, Smithsonian Institution Press, 313p.
- FRYE, F.L. 1991. **Biomedical and Surgical Aspects of Captive Reptile Husbandry**. Malabar, Krieger Publishing Company, 712p.
- GARNER, M.M.; R. HERRINGTON; E.W. HOWERTH; B.L. HOMER; V.F. NETTLES; R. ISAZA; E.B. SHOTTS & E.R. JACOBSON. 1997. Shell disease in river cotters (*Pseudemys concinna*) and yellow-bellied turtles (*Trachemys scripta*) in a Georgia (USA) lake. **Journal of Wildlife Diseases** 33: 78-86.
- GIBBONS, J.W.; D.E. SCOTT; T.J. RYAN; K.A. BUHLMANN; T.D. TUBERVILLE; B.S. METTS; J.L. GREENE; T. MILLS; Y. LEIDEN; S. POPPY & C.T. WINNE. 2000. The global decline of reptiles, déjà vu amphibians. **Bioscience** 50: 653-666.
- JACOBSON, E.R.; J.M. GASKIN; M.B. BROWN; R.K. HARRIS; C.H. GARDINER; J.L. LAPOINTE; H.P. ADAMS & C. REGGIARDO. 1991. Chronic upper respiratory tract disease of free-ranging desert tortoises (*Xerobates agassizii*). **Journal of Wildlife Diseases** 27: 296-316.
- KAKIZOE, Y.; K. SAKAOKA; F. KAKIZOE; M. YOSHII; H. NAKAMURA; Y. KANOU & I. UCHIDA. 2007. Successive changes of hematologic characteristics and plasma chemistry values of juvenile loggerhead turtles (*Caretta caretta*). **Journal of Zoo and Wildlife Medicine** 38: 77-84.
- KELLER, J.M.; J.R. KUCKLICK; M.A. STAMPER; C.A. HARMS & P.D. MCCLELLAN-GREEN. 2004. Associations between organochlorine contaminant concentrations and clinical health parameters in loggerhead sea turtles from North Carolina, USA. **Environmental Health Perspectives** 112: 1074-1079.
- KNOTKOVA, Z.; S. MAZANEK; M. HOVORKA; M. SLOBODA & Z. KNOTEK. 2005. Haematology and plasma chemistry of bornean river turtles suffering from shell necrosis and haemogregarine parasites. **Veterinární Medicina** 50: 421-426.
- KRUSCHE, A.V.; F.P. CARVALHO; J.M. MORAES; P.B. CAMARGO; M.V.R. BALLESTER; S. HORNINK; L.A. MARTINELLI & R.L. VICTORIA. 1997. Spatial and temporal water quality variability in the Piracicaba river basin, Brazil. **Journal of the American Water Resources Association** 33: 1117-1123.
- MADER, D.R. 2000. Normal Hematology of Reptiles, p. 1126-1132. *In*: B.V. FELDMAN; N.C. JAIN & J.G. ZINKL (Eds). **Schalm's veterinary hematology: veterinary hematology**. Philadelphia, Blackwell Publishing, 1344p.
- MARKS, S.K. & S.B. CITINO. 1990. Hematology and serum chemistry of the radiated tortoise (*Testudo radiata*). **Journal of Zoo and Wildlife Medicine** 21: 342-344.
- MARTINELLI, L.A.; A.V. KRUSCHE; R.L. VICTORIA; P.B. CAMARGO; M. BERNARDES; E.S. FERRAZ; J.M. MORAES & M.C. BALLESTER. 1999. Effects of sewage on the chemical composition of Piracicaba river, Brazil. **Water, Air and Soil Pollution** 110: 67-79.
- METIN, K.; O. TÜRKÖZAN; F. KARGIN; Y. BASIMOĞLU KOCA; E. TASKAVAK & S. KOCA. 2006. Blood cell morphology and plasma biochemistry of the captive European Pond Turtle *Emys orbicularis*. **Acta Veterinaria Brunensis** 75: 49-55.
- OMETO, J.P.; L.A. MARTINELLI; M.V. BALLESTER; A. GESSNER; A.V. KRUSCHE; R.L. VICTORIA & M. WILLIAMS. 2000. Effects of land use on water chemistry and macroinvertebrates in two streams of the Piracicaba river basin, Southeast Brazil. **Freshwater Biology** 44: 327-337.
- OMETO, J.P.; A. GESSNER; L.A. MARTINELLI; M.C. BERNARDES; A.V. KRUSCHE & P.B. CAMARGO. 2004. Macroinvertebrates community as indicator of land-use changes in tropical watersheds, southern Brazil. **Ecology & Hydrobiology** 4: 35-47.
- PRITCHARD, P.C.H. & P. TREBBAU. 1984. *Phrynops geoffroanus* (Schweigger, 1812), p. 111-117. *In*: P.C.H. PRITCHARD & P. TREBBAU (Eds). **The turtles of Venezuela**. Athens, Society for Study of Amphibians and Reptiles, 414p.
- RODRIGUES, M.T. 2005. The conservation of Brazilian reptiles: challenges for a megadiverse country. **Conservation Biology** 19: 659-664.
- ROGERS, K.D. & D.T. BOOTH. 2004. A method of sampling blood from Australian freshwater turtles. **Wildlife Research** 31: 93-95.
- ROSENFELD, G. 1947. Método rápido de coloração de esfregaços de sangue: noções práticas sobre corantes pancromáticos e estudo de diversos fatores. **Memórias do Instituto Butantan** 20: 315-328.
- SYPEK, J. & M. BORYSENKO. 1988. Reptiles, p. 211-248. *In*: A.F. ROWLEY & N.A. RATCLIFFE (Eds). **Vertebrate blood cells**. Cambridge, University Press, 456p.
- SOUZA, F.L. & A.S. ABE. 2000. Feeding ecology, density and biomass of the freshwater turtle, *Phrynops geoffroanus*, inhabiting a polluted urban river in south-eastern Brazil. **Journal of the Zoological Society of London** 252: 437-446.
- WORK, T.M.; R.A. RAMEYER; G.A. BALAZS; C. CRAY & S.P. CHANG. 2001. Immune status of free-ranging green turtles with fibropapillomatosis from Hawaii. **Journal of Wildlife Diseases** 37: 574-581.

Submitted: 09.VI.2009; Accepted: 06.XII.2009.

Editorial responsibility: Lucélia Donatti