

# Acute static muscle stretching improves manual dexterity in young men

César Rafael M Costa<sup>1||</sup>, Raoni C Dos-Santos<sup>1||</sup>, Welington V de Paula<sup>1</sup>, Wallace MV Ribeiro<sup>1</sup>, Anderson Luiz B Silveira<sup>1||</sup>

<sup>1</sup> Universidade Federal Rural do Rio de Janeiro, Physical Education and Sports, Seropédica, RJ, Brazil

<sup>||</sup> Universidade Federal Rural do Rio de Janeiro, Physiological Science, Seropédica, RJ, Brazil

**OBJECTIVE:** This study aimed to evaluate the influence of acute muscle stretching on manual function.

**METHODS:** The sample consisted of 10 untrained men in a randomized, four test session cross-over experimental design. Each session was composed of only one of two protocols: a) control, or b) single series of passive static stretching; followed by either Minnesota Hand dexterity test or hand grip strength test with eletromyographical recording of reaction time. For data comparison, the Student T-test with significance level of  $p \leq 0.05$  was used.

**RESULTS:** Manual dexterity increased after stretching for both placing and turning tests, with no changes in hand grip strength or reaction time.

**CONCLUSION:** The results show that a 30 second static stretch of the hand decreases time to complete the Minnesota Hand Dexterity test without affecting handgrip strength or hand reaction time; thus it improves manual dexterity of young untrained men.

**KEYWORDS:** Muscle stretching, motor skills, reaction time.

Costa CRM, Dos-Santos RC, Paula WV, Ribeiro WMV, Silveira ALB. Acute static muscle stretching improves manual dexterity in young men. MedicalExpress (São Paulo, online). 2017;4(3):M170306

Received for Publication on May 1, 2017; First review on May 20, 2017; Accepted for publication on June 9, 2017; Online on June 26, 2017

E-mail: cesarufrrj@hotmail.com

## INTRODUCTION

Stretching exercises are often prescribed for persons with reduced motor skills or range of motion. Several reports describe this exercise technique as a necessary activity for better performance in daily life activities.<sup>1-4</sup> However, muscle stretching seems to evoke neural and mechanical adaptations that reduce strength, reaction time and body balance,<sup>3,5-7</sup> it also affects the responsiveness of relevant afferent structures, such as muscle spindles<sup>8</sup> and Golgi tendon organs,<sup>9</sup> which are important structures in the regulation of motor commands.

Manual dexterity, handgrip strength and reaction time are important components to assess hand function, which relate to better quality of daily life<sup>1,10</sup> and recreational activities;<sup>2,11</sup> they also act as indicators of cognitive and exercise performance<sup>12,13</sup> and are determinant factors in the ability to manipulate objects.<sup>14</sup> Additionally, manual dexterity and handgrip strength are closely interdependent,

in an interaction known as strength-dexterity trade-off.<sup>15</sup> Manual dexterity is also related to the central nervous system, since the cortex receives information from several proprioceptors<sup>8,16</sup> to modulate manual motor tasks increasing movement efficiency.<sup>14,17-19</sup> Hence, if muscle stretching does interfere with these factors it could alter the ability to manipulate objects. Because several sports, such as boxing,<sup>20</sup> handball,<sup>21</sup> basketball,<sup>22</sup> and many non-sporting activities such as musical performance<sup>23</sup> require agility and manual dexterity, the assessment of hand function is essential for the development of training techniques and functional rehabilitation of hand movements. However, there is a lack of knowledge about the effect of static stretching on manual dexterity. We hypothesized that muscle stretching would interfere with these factors and consequently alter hand dexterity.

The purpose of this study was to understand the effect of static stretching on manual dexterity, providing evidence for health professionals seeking to prescribe stretching exercises safely and efficiently for athletes and patients who need to improve their manual skills.

DOI: 10.5935/MedicalExpress.2017.03.06

## ■ MATERIALS AND METHODS

The present study was approved by the Research Ethics Committee of the Federal Rural University of Rio de Janeiro, under case # 23083.008201/2014-91. The study was carried out with 10 healthy untrained male subjects (age:  $22 \pm 2$  years; weight  $72.58 \pm 3.5$  kg; height:  $170 \pm 6$  cm; body fat percentile:  $12.69 \pm 3.39$  %; BMI:  $23.91 \pm 1.71$ ) with no present or pre-existent lesions or signs of pain in the arms or hands, no restriction to normal hand movements and no use of medications that could intervene with hand skills such as sedatives, psychotropics or beta blockers. All volunteers read and signed a consent form and were instructed not to participate in physical activities during the week preceding the performance of the experiment. In a previous visit to the laboratory, anthropometric measurements were obtained; subsequently, a familiarization session with the experimental protocols was carried out in order to decrease the influence of learning on results. The experiment had a cross-over randomized design, i.e. the same subjects were used in all experimental groups and compared with themselves; the order in which they performed the experiments was randomized, thus avoiding possible adaptations. The experiments were executed in a week with four testing sessions. Each session consisted of one of two protocols: a) control, in which participants performed no activity; or b) one instance of passive static stretch. After the protocol the participants performed either the Minnesota Hand Dexterity Test or the Hand Grip Test with eletromyographical analysis of hand reaction time. All procedures were carried out in a well-lit and silent environment.

### Muscle stretching

Muscle stretching consisted of a single instance of 30 seconds of passive static stretch. The participant sat with the shoulder in orthostatic position, with elbows at an angle of  $90^\circ$  and forearms supported by the table. The wrist was in supine position with fingertips pointing upward. The participant's wrist was passively stretched by the researcher to the limit of pain and held in position for 30s.

### Hand Dexterity Test

The subject sits on a chair adequate to his height in front of a table so that he has a complete view of the area and equipment.<sup>4</sup> To ensure the consistency of the procedures and test accuracy, general instructions are given before each session to avoid doubts. The volunteer is instructed to start the test immediately after the command and to proceed as fast as possible. To avoid distractions, the presence of observers during the test was not allowed and access to the areas surrounding the lab restricted. The Minnesota Manual Dexterity Test is performed using a board with holes (matrix) and a set of 60 discs (black/red) that fit into the matrix. Two

tests are performed: (a) the placing test, in which all discs must be fitted into the board as fast as possible, using only one hand; (b) the turning test, in which the discs start in the matrix, are removed, turned and refitted on the matrix so that all discs start with the black side up and end with the red side up. In both tests, the time to completion is measured in seconds, and a lower time indicates an increased performance.<sup>11</sup> The performance in this test indicates the capacity to execute a task that requires hand dexterity.<sup>1,4,11</sup>

### Hand Grip Strength

Hand grip strength was measured with a hydraulic dynamometer (Jamar, Sammons Preston Rolyan, 4, Bolingbrook, IL) as described elsewhere.<sup>10,24</sup> The participants sat in a comfortable armless chair, with the arm alongside the body, the elbow flexed at  $90^\circ$  and dominant hand parallel to the body holding the dynamometer. The opposite hand rested over the thigh. The participants were then instructed to perform a maximal contraction with the dominant hand for 3s in each of three tries; a rest period of 30s was allowed between tries and the mean values of the three attempts was used for analysis.

### Hand Reaction Time

For electromyographical analysis, electric signals were recorded on the skin above the forearm with bipolar electrodes and filtered to EMG Spikerbox (Backyard Brains), as described elsewhere.<sup>25,26</sup> To record the signals and measure the reaction time, we used the open-source audio processing program Audacity (Audacity®, Version 2.1.2). The recording was carried out during the above-described hand grip test and the assessment used the mean of values found in the 3 tries.


### Statistical Analysis

Data are displayed as mean  $\pm$  standard deviation. Data analysis was carried out with Prism v6.0 (GraphPad, EUA). Data normality was assessed by the Shapiro-Wilk test. The groups were compared with Student T-test and the adopted level of significance was  $p \leq 0.05$ .


## ■ RESULTS

Muscle stretching significantly improved hand dexterity. Figure 1A shows that the time to complete the placing test was significantly reduced after stretching, from  $55.53 \pm 1.78$  to  $53.63 \pm 3.23$  seconds ( $p=0.0139$ ). Figure 1B shows that the time to complete the turning test was also significantly reduced after stretching, from  $54.20 \pm 2.39$  vs.  $50.35 \pm 5.39$  seconds ( $p = 0.0200$ ). Figure 2 shows that static stretch did not affect hand grip strength ( $33.3 \pm 8.84$  vs.  $33.2 \pm 9.61$ ;  $p = 0.9238$ , Fig. 2a) or reaction time ( $200.8 \pm 30.08$  vs.  $199.2 \pm 41.56$ ;  $p = 0.8981$ , fig 2B)

**A: TIME TO COMPLETE PLACING TEST (SECONDS)**




**B: TIME TO COMPLETE TURNING TEST (SECONDS)**




**Figure 1:** Time to complete the Minnesota Dexterity Test (mean and std dev). Stretching decreased time to complete the placing (A) and turning (B) components of the Minnesota Dexterity Test, showing improved hand dexterity. \*  $p < 0.05$  vs. CTR.

**A: HAND GRIP STRENGTH (Kgf)**



**B: REACTION TIME (mseconds)**



**Figure 2:** Strength and reaction time (mean and std. dev.). Static muscle stretching did not affect hand grip strength (A) or reaction time (B).

## DISCUSSION

This study showed that 30 seconds of static stretching improved manual dexterity, with no change of handgrip strength or hand reaction time. The absence of changes of handgrip strength and hand reaction time after muscle stretching may be due to the moderate volume (1 instance) and intensity (30s) used in our study, because deleterious effects on strength and reaction time are related to greater volume (>2 sets) or intensities ( $\geq 45$ s) of muscle stretching,<sup>5,9,27-29</sup> as a result of a decreased motor unit activation.<sup>9,30</sup> The impact of muscle stretching on these variables relates to the stretching protocol used;<sup>31</sup> in this sense, conflicting results have been reported, in

which muscle stretching improved,<sup>32</sup> decreased<sup>5,6,9,27,28</sup> or had no effect on these variables,<sup>33-36</sup> in different stretching protocols. In a study with a similar methodological design Silva *et al.*<sup>6</sup> reported that acute static stretching did not affect strength in men; similar results are found in another study.<sup>37</sup> Decreases in strength after stretching were observed in other groups, such as women<sup>6</sup> or participants experienced in strength training,<sup>38</sup> both of which differ from our sample of untrained male subjects. However, Knudson<sup>29</sup> showed, in a sample similar to ours, an exponential decline of grip strength after stretching that was only significant after 4 sets of 10 seconds of static stretching, which demonstrates that lower intensities and volumes of stretching are not sufficient to induce changes on strength.

Interestingly, our results showed that passive static stretch, performed at an intensity range that does not affect strength, decreased placing and turning test time, which means an improvement of manual dexterity. It is conceivable that stretching induced an increase in cortical excitability, in order to compensate the inhibition of moto-neurons at spinal level, resulting in enhanced descending excitatory drive.<sup>39</sup> Likewise, muscle stretching increases the proprioceptors activity before the onset of movement enabling a basis for motor command programming and organization at central and peripheral level, and thus improved performance.<sup>8,39-41</sup> Additionally, central nervous system and proprioceptors are related to hand dexterity<sup>14,15,17,42</sup> through both spinal reflexes and central connections,<sup>41,43</sup> which ensure the generation of the correct pattern of muscle activity or automatic adjustments of hand movement.<sup>8,39,40</sup> Thus, muscle stretching possibly increased the quality of the information and the capacity of response of the subject, promoting a sequence of muscular activation and modulation of motor command better suited to the task. This increased cortical drive may have influenced the increase in dexterity found in this study. This study has limitations: participants were not controlled for intervenient factors such as hormonal levels, sleep time and feeding patterns, which may have influenced the results found on this study. Only one population was assessed in this study. Future studies should include different populations, such as participants trained in sports or with limitations to hand function.

## ■ CONCLUSION

Our results indicate that moderate volume and intensity of static stretching of the hand/forearm muscles may increase manual dexterity with no change in strength and hand reaction time. Therefore, stretching may increase hand dexterity if it is of an intensity that does not induce changes in other components of hand function. Greater volumes or intensities should be recommended with care, to avoid potential disturbances in manual dexterity. Further studies are pivotal to assess and establish a safe muscle stretching volume and intensity.

## ■ CONFLICTS OF INTEREST

The authors declare that they have no conflict of interest.

## ■ AUTHOR CONTRIBUTION

Study concept and design: CRMC, ALBS. Analysis and interpretation of data: CRMC, RCS, ALBS. Drafting/revision of the manuscript: CRMC, RCS, WVP, WMV, ALBS. Study supervision: ALB. All authors read and approved the final version of the manuscript.

## EFEITO AGUDO DO ALONGAMENTO ESTÁTICO SOBRE A DESTREZA MANUAL DE JOVENS DESTREINADOS

**OBJETIVO:** Este estudo objetivou avaliar a influência do alongamento sobre a função manual.

**MÉTODOS:** A amostra foi composta por 10 homens destreinados em um delineamento experimental cross-over randomizado, com quatro sessões de testes. Cada sessão foi constituída apenas por um dos protocolos: a) controle ou b) série única de alongamento estático passivo seguidos pelo minnesota Hand Dexterity Test ou pelo Teste de Preensão Manual com análise eletromiográfica do Tempo de Reação Manual. Para as comparações dos dados, o teste T de Student foi usado, com o índice de significância adotado de  $p \leq 0,05$ .

**RESULTADOS:** O alongamento aumentou a destreza manual em ambos *Placing e Turning Tests*, sem alterar a força de preensão manual ou o tempo de reação.

**CONCLUSÃO:** Os resultados indicam que o alongamento melhorou a destreza manual de jovens destreinados.

**PALAVRAS CHAVE:** Alongamento muscular; destreza motora; tempo de reação.

## ■ REFERENCES

1. Botelho MF, Azevedo A. Manual reaction speed and manual dexterity in elderly people: a comparative study between elderly practitioners and non-practitioners of physical activity. *Sport Sci.* 2009;2(1):35-43.
2. Carmeli E, Patish H, Coleman R. The aging hand. *J Gerontol A Biol Med Sci.* 2003 Feb;58(2):146-52. DOI: 10.1093/gerona/58.2.M146
3. Guissard N, Duchateau J. Neural aspects of muscle stretching. *Exerc Sport Sci Rev.* 2006;34(4):154-8. DOI: 10.1249/01.jes.0000240023.30373.eb
4. Lourenção MIP, Tsukimoto GR, Battistella LR. O teste de destreza manual Minnesota adaptado utilizado como avaliação do potencial de uso de membros superiores de pacientes hemiplégicos. *Acta fisiátrica.* 2007;14(1):56-61.
5. Behm DG, Bambury A, Cahill F, Power K. Effect of acute static stretching on force, balance, reaction time, and movement time. *Med Sci Sports Exerc.* 2004;36:1397-402. DOI: 10.1249/01.MS-S.0000135788.23012.5F
6. Silva GV de LC, Silveira ALB, Di Masi F, Bentes CM, Miranda HL, Novaes J da S. Efeito agudo do alongamento estático sobre a força muscular isométrica. *ConScientiae Saúde.* 2012;11(2):274-80. DOI:10.5585/ConsSaude.v11n2.3124
7. Marek SM, Cramer JT, Fincher AL, Massey LL, Dangelmaier SM, Purkayastha S, et al. Acute effects of static and proprioceptive neuromuscular facilitation stretching on muscle strength and power output. *J Athl Train.* 2005;40(2):94-103.
8. Park S, Toole T, Lee S. Functional roles of the proprioceptive system in the control of goal-directed movement. *Percept Mot Ski.* 1999;88(2):631-47. DOI: 10.2466/PMS.88.2.631-647
9. Fowles JR, Sale DG, MacDougall JD. Reduced strength after passive stretch of the human plantarflexors. *J Appl Physiol.* 2000;89(3):1179-88.
10. Reis MM, Arantes PMM. Medida da força de preensão manual- validade e confiabilidade do dinamômetro saehan. *Fisioter e Pesqui.* 2011;18:176-81. DOI: 10.1590/S1809-29502011000200013
11. Pinto MJC. Aptidão Física, Destreza Manual e Sensibilidade Proprioceptiva Manual no Idoso: Estudo em praticantes e não praticantes de actividade física. 2003. <http://hdl.handle.net/10216/10410>

12. Lebrun CM. Effect of the different phases of the menstrual cycle and oral contraceptives on athletic performance. *Sport Med.* 1993;16(6):400-30.
13. Seeman TE, Berkman LF, Charpentier PA, Blazer DG, Albert MS, Tinetti ME. Behavioral and psychosocial predictors of physical performance: MacArthur studies of successful aging. *J Gerontol A Biol Sci Med Sci.* 1995;50(4):M177-83. DOI: 10.1093/gerona/50A.4.M177
14. Olafsdottir HB, Zatsiorsky VM, Latash ML. The effects of strength training on finger strength and hand dexterity in healthy elderly individuals. *J Appl Physiol.* 2008;105(4):1166-78. DOI: 10.1152/jappphysiol.00054.2008
15. Shinohara M, Li S, Kang N, Zatsiorsky VM, Latash ML. Effects of age and gender on finger coordination in MVC and submaximal force-matching tasks. *J Appl Physiol.* 2003;94(1):259-70. DOI: 10.1152/jappphysiol.00643.2002
16. Laufer Y, Hocherman S, Dickstein R. Accuracy of reproducing hand position when using active compared with passive movement. *Physiother Res Int.* 2001;6(2):65-75. DOI: 10.1002/pri.215
17. Higo N. Training-induced Recovery of Manual Dexterity after a Lesion in the Motor Cortex. *Keio J Med.* 2010;59(1):4-9. DOI: 10.2302/kjm.59.4
18. Latash ML, Kang N, Patterson D. Finger coordination in persons with Down syndrome: atypical patterns of coordination and the effects of practice. *Exp Brain Res.* 2002;146(3):345-55. DOI: 10.1007/s00221-002-1189-3
19. Shinohara M, Latash ML, Zatsiorsky VM. Age effects on force produced by intrinsic and extrinsic hand muscles and finger interaction during MVC tasks. *J Appl Physiol.* 2003;95(4):1361-9. DOI: 10.1152/jappphysiol.00070.2003
20. Darby D, Moriarity J, Pietrzak R, Kutcher J, McAward K, McCrory P. Prediction of winning amateur boxers using pretournament reaction times. *J Sports Med Phys Fitness.* 2014;54(3):340-6.
21. Baştüre E, Stan Z, Acsinte A. The importance of coordination in the technical training specific to handball players. In: *Annals of "Dunarea de Jos" University of Galati Fascicle XV.* 2013. p. 7-12.
22. Kioumourtzoglou E, Derri V, Tzetzis G, Theodorakis Y. Cognitive, Perceptual, and Motor Abilities in Skilled Basketball Performance. *Percept Mot Skills.* 1998;86(3):771-86.
23. Metcalf CD, Irvine TA, Sims JL, Wang YL, Su AWY, Norris DO. Complex hand dexterity: A review of biomechanical methods for measuring musical performance. Vol. 5, *Frontiers in Psychology.* 2014. p. 414.
24. Bohannon RW, Peolsson A, Massy-Westropp N, Desrosiers J, Bear-Lehman J. Reference values for adult grip strength measured with a Jamar dynamometer: a descriptive meta-analysis. *Physiotherapy.* 2006;92(1):11-5. DOI: 10.1016/j.physio.2005.05.003
25. Marzullo TC, Gage GJ. The SpikerBox: a low cost, open-source bioamplifier for increasing public participation in neuroscience inquiry. *PLoS One.* 2012;7(3):e30837. DOI: 10.1371/journal.pone.0030837
26. Shannon KM, Gage GJ, Jankovic A, Wilson WJ, Marzullo TC. Portable conduction velocity experiments using earthworms for the college and high school neuroscience teaching laboratory. *Adv Physiol Educ.* 2014;38(1):62-70. DOI: 10.1152/advan.00088.2013
27. Behm DG, Button DC, Butt JC. Factors affecting force loss with prolonged stretching. *Can J Appl Physiol.* 2001;26(3):262-72. DOI: 10.1139/h01-017
28. Costa PB, Ryan ED, Herda TJ, Walter AA, Hoge KM, Cramer JT. Acute effects of passive stretching on the electromechanical delay and evoked twitch properties. *Eur J Appl Physiol.* 2010;108(2):301-10. DOI: 10.1007/s00421-009-1214-3
29. Knudson D, Noffal G. Time course of stretch-induced isometric strength deficits. *Eur J Appl Physiol.* 2005;94(3):348-51. DOI: 10.1007/s00421-004-1309-9
30. Winchester JB, Nelson AG, Kokkonen J. A Single 30-s Stretch Is Sufficient to Inhibit Maximal Voluntary Strength. *Res Q Exerc Sport.* 2009;80(2):257-61. DOI: 10.1080/02701367.2009.10599560
31. Behm DG, Chaouachi A. A review of the acute effects of static and dynamic stretching on performance. *Eur J Appl Physiol.* 2011;111(11):2633-51. DOI: 10.1007/s00421-011-1879-2
32. Costa PB, Graves BS, Whitehurst M, Jacobs PL. The acute effects of different durations of static stretching on dynamic balance performance. *J Strength Cond Res.* 2009;23(1):141-7. DOI: 10.1519/JSC.0b013e31818eb052
33. Alpkaya U, Koceja D. The effects of acute static stretching on reaction time and force. *J Sports Med Phys Fitness.* 2007;47(2):147-50.
34. Molacek ZD, Conley DS, Evetovich TK, Hinnerichs KR. Effects of low- and high-volume stretching on bench press performance in collegiate football players. *J Strength Cond Res.* 2010;24(3):711-6. DOI: 10.1519/JSC.0b013e3181c7c242
35. Perrier ET, Pavol MJ, Hoffman MA. The acute effects of a warm-up including static or dynamic stretching on countermovement jump height, reaction time, and flexibility. *J Strength Cond Res.* 2011;25(7):1925-31. DOI: 10.1519/JSC.0b013e3181e73959
36. Winke MR, Jones NB, Berger CG, Yates JW. Moderate static stretching and torque production of the knee flexors. *J Strength Cond Res.* 2010;24(3):706-10. DOI: 10.1519/JSC.0b013e3181c7c557
37. Cardozo G, Torres JB, Dantas EHM, Simão R. Comportamento da força muscular após o alongamento estático. *Rev Treinamento Desport.* 2006;7(1):73-6.
38. Costa e Silva GVL, Silveira ALB, Di Masi F, Bentes CM, de Sousa MSC, Novaes JS. Acute effect of different stretching methods on isometric muscle strength. *Acta Sci Heal Sci.* 2014;36(1):51-7. DOI: 10.4025/actascihealthsci.v36i1.15581
39. Gruber M, Linnamo V, Strojnik V, Rantalainen T, Avela J. Excitability at the Motoneuron Pool and Motor Cortex Is Specifically Modulated in Lengthening Compared to Isometric Contractions. *J Neurophysiol.* 2009;101(4):2030-40. DOI: 10.1152/jn.91104.2008
40. Duchateau J, Enoka RM. Neural control of shortening and lengthening contractions: influence of task constraints. *J Physiol.* 2008;586(24):5853-64. DOI: 10.1113/jphysiol.2008.160747
41. Kandel E, Schwartz J, Jessell T. *Principles of Neural Science, Fourth Edition.* McGraw-Hill Companies, Incorporated; 2000.
42. Latash ML, Turvey MT. *Dexterity and Its Development.* L. Erlbaum Associates; 1996.
43. Magill RA. *Motor Learning and Control: Concepts and Applications.* McGraw-Hill Companies; 2011.