

ORIGINAL ARTICLE

Effect of feeding frequency and water salinization on early development of *Pyrrhulina brevis*, an Amazonian ornamental fish

Arlindo dos S. PINHEIRO-JUNIOR¹, Leonnan C. C. de OLIVEIRA¹, Bruno José C. F. EIRAS¹, Wesley dos S. LIMA¹, Galileu C. VERAS², Lorena B. de MOURA¹, Ana L. SALARO³, Daniel A. V. CAMPELO^{1*} 

¹ Universidade Federal do Pará, Instituto de Estudos Costeiros, Alameda Leandro Ribeiro, s/n, 68600-000, Bragança, Pará, Brazil

² Universidade Federal de Minas Gerais, Escola de Medicina Veterinária, Av. Presidente Antônio Carlos 6627, 31270-901, Belo Horizonte, Minas Gerais, Brazil

³ Universidade Federal de Viçosa, Departamento de Biologia Animal, Av. Peter Henry Rolfs, s/n, 36570-900, Viçosa, Minas Gerais, Brazil

* Corresponding author: danielvc@ufpa.br;  <https://orcid.org/0000-0001-9204-3566>

ABSTRACT

Pyrrhulina brevis is an endemic fish from the Amazon basin that is valued in the ornamental fish market. In larviculture, fish are very sensitive to stressors and have a high mortality rate. Salinized water may provide more energy to overcome stress and feeding frequency is an important strategy to improve productive performance. Therefore, the aim of this study was to evaluate the best feeding frequency and the best level of water salinization for larval development of *P. brevis*. A total of 360 post-larvae of *P. brevis* (5.26 ± 1.65 mg, 5.57 ± 0.68 mm) were randomly distributed in 36 aquariums (1 L) in a 3x4 factorial scheme, to evaluate three concentrations of salt in water (0, 1 and 2 g L⁻¹) and four feeding frequencies (once, twice, three and four times a day). We offered 150 *Artemia* nauplii per day and per post-larvae during 15 days. The fish kept at 1 g L⁻¹ salinized water, and those fed 3 and 4 times a day presented significantly greater final length and weight, weight gain, and specific growth rate for length and weight. An interaction between salinized water and feeding frequency was observed for survival rate. We conclude that, during the initial rearing phase of *P. brevis*, feeding frequency of three times a day and salinized water at 1 g L⁻¹ is recommended.

KEYWORDS: productive performance, nutritional management, live foods, ornamental fish farming

Efeito da frequência alimentar e da salinização da água sobre o desenvolvimento inicial de *Pyrrhulina brevis*, um peixe ornamental da Amazônia

RESUMO

Pyrrhulina brevis é um peixe endêmico da bacia amazônica que apresenta bons valores no mercado de peixes ornamentais. Na larvicultura, os peixes são muito sensíveis aos estressores e apresentam alta taxa de mortalidade. A água salinizada pode fornecer mais energia para superar o estresse e a frequência alimentar é uma estratégia importante para melhorar o desempenho produtivo. Portanto, o objetivo deste trabalho foi avaliar a melhor frequência alimentar e o melhor nível de salinização da água para o desenvolvimento larval de *P. brevis*. Um total de 360 pós-larvas de *P. brevis* ($5,26 \pm 1,65$ mg, $5,57 \pm 0,68$ mm) foram distribuídos aleatoriamente em 36 aquários (1 L) em esquema fatorial 3x4, para avaliar três concentrações de sal na água (0, 1 e 2 g L⁻¹) e quatro frequências de alimentação (uma, duas, três e quatro vezes ao dia). Foram oferecidas 150 náuplios de *Artemia* por dia e por pós-larva, durante 15 dias. Os peixes mantidos com 1 g L⁻¹ de água salinizada e os alimentados 3 e 4 vezes ao dia apresentaram comprimento e peso finais, ganho de peso e taxa de crescimento específico para comprimento e peso significativamente maiores. Uma interação entre água salinizada e frequência alimentar foi observada para a taxa de sobrevivência. Concluímos que, durante a fase inicial de criação de *P. brevis*, recomenda-se uma frequência alimentar de três vezes ao dia e água salinizada a 1 g L⁻¹.

PALAVRAS-CHAVE: desempenho produtivo, manejo nutricional, alimento vivo, piscicultura ornamental

CITE AS: Pinheiro-Junior, A.S.; Oliveira, L.C.C.; Eiras, B.J.C.F.; Lima, W.S.; Veras, G.C.; Moura, L.B.; Salaro, A.L.; Campelo, D.A.V. 2023. Effect of feeding frequency and water salinization on early development of *Pyrrhulina brevis*, an Amazonian ornamental fish. *Acta Amazonica* 53: 107-113.

INTRODUCTION

During fish production, larviculture is one of the most critical phases (Abe *et al.* 2015). Post-larvae are more sensitive to stressors such as changes in water quality and nutritional management (Zuanon *et al.* 2011). The initial feeding directly influences the growth performance of the post-larvae and the supply of live food during larviculture has been the most used strategy to ensure the well-being and development of animals in the first days of life (Fosse *et al.* 2013).

Among live foods, *Artemia sp.* stands out for its ease of production and storage, in addition to its nutritional quality (Silva and Mendes 2006; Kestemont *et al.* 2007). Fish fed *Artemia* nauplii in the first days of life have good growth rates and low mortality (Campelo *et al.* 2020). However, it is important to control the daily food supply so that it can be used more efficiently by the post-larvae. Lack of food is harmful to the development of fish, and excess results in an unfavorable cost-benefit relation to the producer, as manpower for *Artemia* nauplii processing and fish feeding is a considerable part of larviculture expenses, and is harmful for the post-larvae, since excess food can decrease water quality (Oliveira *et al.* 2020).

Feeding frequency represents the number of meals offered to fish per day and is directly related to the improvement in growth performance variables (Xie *et al.* 2011). However, the ideal feeding frequency depends on the species, life stage and water quality (Thomassen and Fjaera 1996). Since post-larvae have a short gastrointestinal tract, the time spent by the food in the tract is also short (Lee *et al.* 2000). Providing food at an inappropriate frequency can promote intestinal overload, which results in lower nutrient absorption (Riche *et al.* 2004). Thus, determining the optimal feeding frequency is important to optimize food utilization and production costs (Abe *et al.* 2016).

Another relevant point to be considered during the larviculture of freshwater fish is the salinization of breeding water, through the addition of common salt (NaCl). The salinized water can help the initial development of the post-larvae, reducing ion losses and excessive water absorption due to the osmotic differences between the fish plasma and the external environment (Varsamos *et al.* 2005; Fabregat *et al.* 2015). This makes it possible to redirect energy to other physiological processes such as growth (Baldisserotto *et al.* 2007).

Salinized water also benefits *Artemia* nauplii, making this microcrustacean live longer and increasing its availability time as fresh food for the post-larvae (Beux and Zaniboni-Filho 2018). Adding salt to the water is also a good alternative to reduce stress in the larviculture environment (Jomori *et al.* 2012), since the increase in salt concentration in the water allows fish to reduce energy expenditure with the maintenance of osmotic balance (Sampaio and Bianchini 2002; Salaro *et*

al. 2012). However, the use of inadequate concentrations of salt in the water can compromise animal welfare and generate physiological and behavioral changes (Luz 2007; Luz and Santos 2008; Dias *et al.* 2016), that can result in reduced growth and survival of freshwater fish (Luz *et al.* 2013).

Pyrhulina brevis Steindachner, 1876 (Characiformes: Lebiasinidae) is an endemic species of the Amazon basin that has a cylindrical, elongated body and an upturned mouth, with gregarious habit, normally occurring in shoals, which makes it attractive to the ornamental market, and showing territorial behavior during the reproductive period (Weitzman and Weitzman 2003; Arias and Rossi 2005; Silva *et al.* 2016). These characteristics, as well as its rapid adaptation to rearing conditions, place *P. brevis* in the spotlight of the international ornamental fish trade (Abe *et al.* 2021).

Since *P. brevis* is an ornamental Amazonian fish that reproduces easily under captive conditions and has a promising consumer market, it is important to optimize protocols for the production of young forms, such as the best way to offer live food and to salinize the water to reduce stress and increase the viability of *Artemia* nauplii used as live food. Other studies have already evaluated the use of salinized water (Abe *et al.* 2015), prey concentrations (Oliveira *et al.* 2020) and feed deprivation cycles (Abe *et al.* 2021) for *P. brevis*. However, no study so far has tested the interaction of the water salinization level with the feeding frequency for this species. Thus, the present study aimed to evaluate the best feeding frequency and water salinization level during larviculture of *P. brevis* both in terms of fish growth and labor production costs.

MATERIAL AND METHODS

The experiment was approved by the ethics committee on the use of animals of Universidade Federal do Pará – UFPA (Process # 7656100517 CEUA/UFPA) and carried out at the Pisciculture Laboratory (LAPIS) of UFPA, Bragança campus, Pará state, Brazil.

Fish and experimental conditions

Post-larvae of *P. brevis* were obtained from four breeding pairs held at the Pisciculture Laboratory under controlled conditions. After hatching, the larvae were observed daily to monitor the consumption of the yolk sac. On the seventh day after hatching, the yolk sac was completely consumed and the mouth opening had reached the ideal size for the acceptance of exogenous food, characterizing the beginning of the post-larvae phase. For the initial biometrics, due to the small size and fragility of the fish, a sample of 40 individuals was used to estimate the initial mean weight and total length (5.26 ± 1.65 mg and 5.57 ± 0.68 mm, respectively). A total of 360 post-larvae were randomly selected and distributed in 36 aquariums (1L volume), in a 3x4 factorial scheme with three replications per treatment. The factor water salinization consisted of three

concentrations of sodium chloride in the water (0, 1 and 2 g L⁻¹) and the factor feeding frequency consisted of four levels: feeding live *Artemia* nauplii once a day (08:00), twice a day (08:00 and 14:00), three times a day (08:00, 10:00 and 14:00) and four times a day (08:00, 10:00, 14:00 and 16:00). The experiment duration was of 15 days, corresponding to the duration of larval development of *P. brevis*.

Artemia hatching

Overall, each post-larva was fed 150 nauplii of *Artemia* per day (1500 nauplii per replicate per day), as indicated by Abe *et al.* (2015). This amount was distributed among the number of daily feeding bouts corresponding to each treatment level. A total of 5 g L⁻¹ of *Artemia* cysts were hatched daily in a transparent container containing 2 L of salinized water at a concentration of 35 g L⁻¹. The containers for the hatching of the *Artemia* were kept at room temperature under artificial lighting of fluorescent lamps (15 W) and constant aeration for 24 hours, the period necessary for the hatching of the nauplii. After hatching, the nauplii were removed by siphoning to separate the cyst residues, then washed in running water to remove the saline water, so as not to influence the experimental treatments, and transferred to a 200-ml beaker. The density of nauplii was estimated by collecting 0.5 ml from the beaker and counting the number of nauplii in this sample under a stereomicroscope with 40x magnification (QUIMIS 14 Q714Z-2, Diadema, Brazil). From this sample, the volume to be supplied to each aquarium was calculated.

Water quality parameters

The water quality parameters were measured every three days. Temperature (26.28 ± 0.04 °C), pH (6.53 ± 0.05), dissolved oxygen (6.28 ± 0.06 mg L⁻¹) and electrical conductivity (1.83 ± 0.03 µS cm⁻¹) were measured with the aid of a multiparameter device (Horiba Advanced Techno Co. Ltd., Kyoto, Japan). Total ammonia (0.70 ± 0.03 mg L⁻¹) was measured using a Labcon Test Kit (Alcon Ltda., Brazil). All parameters remained in appropriate conditions for Amazonian fish farming (Silveira *et al.* 2009). Two hours after the last feeding of the day, 50% of the water of each aquarium was siphoned out and replaced to remove feces and possible food residues, to ensure water quality and the well-being of the post-larvae.

Growth performance

At the end of the experimental period, all fish were counted, measured and weighed with the aid of an analytical balance (Gehaka AG200, Real Parque, São Paulo, Brazil, precision 0.0001 g) and a digital caliper (Vonder, 0.01 mm). We determined the following growth performance parameters:

- Final length (FL) (mm);
- Final weight (FW) (mg);
- Weight gain (WG) (mg) = final weight – initial weight;

- Length gain (LG) (mm) = final length – initial length;
- Specific growth rate in weight (SGR_w) (% day⁻¹) = [(ln FW – ln initial weight)/15] *100, where 15 is the duration of the experiment in days;
- Specific growth rate in length (SGR_L) (% day⁻¹) = [(ln FL – ln initial length)/15] *100;
- Batch uniformity for weight (UW) (%) (Furuya *et al.* 1998) = (number of fish with weight ± 20% of the mean per experimental unit)/total number of fish per experimental unit *100;
- Batch uniformity for length (UL) (%) = (number of fish with length ± 20% of the mean per experimental unit)/total number of fish per experimental unit *100;
- Survival rate (SR) (%) = final number of fish/initial number of fish*100.

Labor cost

The labor cost was estimated based on the Brazilian minimum monthly wage in April 2021 (USD 200.20, with an exchange rate of 1 USD = BRL 5.49). Considering a workload of eight hours per day and 24 days per month, the labor cost per hour was USD 1.15. The total feeding time of the post-larvae in hours was calculated by multiplying the feeding frequency by a feeding time of 30 minutes for a standard 1,000 post-larvae, adjusted according to the survival rate among treatments. The labor cost (USD month⁻¹) was calculated as the ratio between the total feeding time for a 15-day trial period and the hourly labor value (Mankiw 2020).

Statistical analysis

The normality and homoscedasticity of the response variables were confirmed by the Lilliefors and Bartlett tests, respectively. Each growth performance variable was analyzed with two-way analysis of variance. When no interaction between salinity and feeding frequency was detected, the variable was analyzed with single-factor ANOVA. The labor cost was analyzed only as a single-factor ANOVA, since the salinization of water was not considered for the calculation. For significant ANOVAs, the post-hoc Tukey test was used for pairwise comparison of means. The significance level was 5% (P < 0.05) for all analysis. The data were analyzed using the SAEG 9.1 software.

RESULTS

The growth performance variables of FL, LG, FW, WG, SGR_w and SGR_L varied significantly with the feeding frequency and with the water salinization level, but there was no interaction between the factors (Table 1). No significant differences were observed for UW and UL (Table 1). For the fish survival rate, an interaction between feeding frequency and salinized water was observed (Table 2).

Table 1. Growth performance parameters of *Pyrhulina brevis* post-larvae submitted to different feeding frequencies and water salinization levels. FL = final length; LG = length gain; FW = final weight; WG = weight gain; SGR_w = specific growth rate for weight; SGR_L = specific growth rate of length; WU = weight uniformity; LU = length uniformity; SR = survival rate.

Treatment	Growth performance parameters								
	FL (mm)	LG (mm)	FW (mg)	WG (mg)	SGR _w (% dia ⁻¹)	SGR _L (% dia ⁻¹)	WU (%)	LU (%)	SR (%)
Feeding frequency (N feedings per day)									
1	9.9 ± 0.5C	5.7 ± 0.5C	7.5 ± 1.0B	6.0 ± 1.0B	10.73 ± 0.9B	5.7 ± 0.3B	46.4 ± 18.0	96.5 ± 4.6	97.8 ± 3.5
2	10.4 ± 0.3B	6.1 ± 0.3B	8.9 ± 0.5A	7.4 ± 0.5A	11.93 ± 0.4A	6.0 ± 0.2B	51.1 ± 14.6	95.6 ± 4.9	100.0 ± 0.0
3	10.8 ± 0.2A	6.5 ± 0.2A	9.1 ± 0.5A	7.6 ± 0.5A	12.04 ± 0.4A	6.2 ± 0.1A	53.3 ± 12.1	95.6 ± 5.9	100.0 ± 0.0
4	10.9 ± 0.2A	6.7 ± 0.2A	9.2 ± 0.5A	7.7 ± 0.5A	12.14 ± 0.4A	6.3 ± 0.2A	54.4 ± 12.1	98.8 ± 2.2	97.8 ± 3.5
Water salinization (g L⁻¹)									
0	10.1 ± 0.5C	5.9 ± 0.5C	7.8 ± 0.9B	6.3 ± 0.9B	11.0 ± 0.8B	5.8 ± 0.4C	96.6 ± 4.6	96.6 ± 0.0	98.3 ± 2.8
1	19.9 ± 0.3A	6.7 ± 0.3A	9.4 ± 0.5A	7.9 ± 0.5A	12.2 ± 0.4A	6.3 ± 0.2A	91.1 ± 5.9	100.0 ± 0.0	100.0 ± 0.0
2	10.5 ± 0.5B	6.3 ± 0.3B	7.4 ± 0.5C	7.4 ± 0.5C	11.9 ± 0.4C	6.0 ± 0.2B	95.7 ± 5.0	95.7 ± 5.0	98.3 ± 2.8
Feeding frequency	***	***	***	***	***	***	NS	NS	NS
Water salinization	***	***	***	***	***	***	NS	NS	NS
Interaction	NS	NS	NS	NS	NS	NS	NS	NS	**

Mean values within each variable and factor with different letters differ significantly according to a Tukey test at 5% probability. ANOVA: NS: not significant (P > 0.05); *P < 0.05; **P < 0.01; ***P < 0.001.

Table 2. Survival rate of *Pyrhulina brevis* post-larvae in different levels of water salinization and feeding frequency. Values are the mean ± standard deviation of three replicates.

Feeding frequency (N feedings per day)	Water salinization (g L ⁻¹)		
	0	1	2
1	93.3 ± 4.4 bB	100 ± 0.0 aA	100 ± 0.0 aA
2	100 ± 0.0 aA	100 ± 0.0 aA	100 ± 0.0 aA
3	100 ± 0.0 aA	100 ± 0.0 aA	100 ± 0.0 aA
4	100 ± 0.0 aA	100 ± 0.0 aA	93.3 ± 4.4 bB

Means with different lowercase letters in the same line indicate significant pairwise difference (P < 0.05) according to a Tukey test (n=3). Means with different capital letters in the same column indicate significant pairwise difference (P < 0.05) according to a Tukey test (n=3).

Post-larvae subjected to salinized water of 1 g L⁻¹ showed significantly higher values for FL, LG, FW, WG, SGR_w and SGR_L compared to the other salinization levels. FL, LG and SGR_L were significantly higher for feeding frequencies of three and four times a day, while FW, WG and SGR_w were significantly higher for feeding frequencies of two, three and four times a day (Table 1). The survival rate was 100% in all treatment combinations, except for fish fed once a day at 0 g L⁻¹ salinity and fish fed four times a day at 2 g L⁻¹ salinity, with two dead larvae each. The post-larvae were found dead with no signs of injury or any trace of disease.

As expected, the labor cost increased directly and significantly with the increase of the feeding frequency. The labor cost was of USD 8.0 ± 0.3, 16.3 ± 0.0, 24.5 ± 0.0 and 31.9 ± 1.1 for the feeding frequencies of one, two, three and four times a day, respectively.

DISCUSSION

In the present study, *P. brevis* post-larvae submitted to feeding frequencies of three and four times a day presented significantly higher values of final length, length gain, final weight and weight gain. Our results confirm that providing food at the appropriate feeding frequency can improve the development of *P. brevis* post-larvae. Similar results were observed for post-larvae of *Heros severus* Heckel, 1840, which had higher final weight, weight gain, specific growth rate, final length and length gain when fed three, four and five times per day (Paixão *et al.* 2019). Likewise, post-larvae of *Pterophyllum scalare* Schultzze, 1823, had higher values for weight gain, specific growth rate and survival rate when fed four times a day (Eiras *et al.* 2019). On the other hand, a feeding frequency of twice times a day is recommended in *Betta splendens* Regan, 1910, larviculture in order to obtain better growth, uniformity and survival (Sales *et al.* 2016).

Generally, fish in the early stages show better development when subjected to a higher feeding frequency, due to the higher metabolic rate of the animals at this stage of development (Pouey *et al.* 2012). A low feeding frequency implies a larger amount of food being supplied per meal, resulting in a suboptimal food consumption and nutrient absorption, due to the fast passage of the food through the gastrointestinal tract of the post-larvae (Luz and Portella 2005; Salaro *et al.* 2012; Abe *et al.* 2016). Post-larvae usually have more opportunities to obtain sufficient food at higher feeding frequencies (Xie *et al.* 2011). However, very high feeding frequencies can also cause hierarchy problems, generating food dispute and favoring the consumption of food by dominant fish, due to the competition for the smaller amount of food

per feeding bout. In addition, higher feeding frequencies can significantly increase the production cost and labor cost (Luz and Portella 2015; Okomoda *et al.* 2019).

In the present study, post-larvae subjected to salinized water of 1 g L⁻¹ had the best values for final length and weight, length and weight gain, and specific growth rate for length and weight, when compared to fish kept in non-salinized water or water salinized at 2 g L⁻¹. Similarly, other studies that tested the effect of water salinization on post-larvae of *P. brevis* obtained best production parameters at 1 g L⁻¹ (Oliveira *et al.* 2020) and at 2 g L⁻¹ (Abe *et al.* 2015), although the latter did not test concentrations lower than 2 g L⁻¹ and neither of the two studies related the salinized water with the feeding frequency of the fish. The use of salinized water at concentrations higher than 2 g L⁻¹ also resulted in negative effects on growth and survival of post-larvae of *Colossoma macropomum* (Cuvier, 1816), *Brycon amazonicus* (Spix and Agassiz, 1829), *Astronotus ocellatus* (Agassiz, 1831) (Jamori *et al.* 2013), *Hypsolebias radiseriatus* Costa, 2012 (Araújo *et al.* 2021), *Oreochromis niloticus* (Linnaeus, 1758) (Luz *et al.* 2013) and *Brycon vonoi* Lima, 2017 (Coraspe-Amaral *et al.* 2017). Therefore, our results confirm that freshwater fish species perform better when raised in slightly higher salinity levels compared to their natural conditions (Boeuf and Payan 2001), but generally lower than 2 g L⁻¹. The use of inadequate concentrations of salt in the water can lead to reduced development and survival of freshwater fish raised in captivity (Luz *et al.* 2012). High concentrations of salt in the water compromise animal welfare and generate physiological and behavioral changes (Luz 2007; Luz and Santos 2008; Dias *et al.* 2016). These effects may be related to osmoregulatory imbalance when the concentration of salts in water exceeds the limits of homeostasis (Barton 2002).

In this study, an interaction between feeding frequency and salinized water was observed for the survival rate of post-larvae, with the lowest survival being for fish submitted to a 2 g l⁻¹ treatment and feeding four times a day. The fish in this treatment could be suffering stress caused by the high salinized water and very fractional feeding (Hayashi *et al.* 2004; Luz *et al.* 2012), which may have benefited the dominant fish, so the combination of the two stressors harmed the fish's survival. In the same way, fish submitted to the treatment of one feeding per day in water without the addition of salt also had impaired survival, probably because the *Artemia* nauplii supplied in this treatment showed faster mortality and ended up being less attractive to the fish (Portella *et al.* 2000; Cardoso and Igarashi 2009). The reduced survival rate of fish fed once a day in association with fresh water confirms the hypothesis that salt can benefit post-larvae, probably due to reduced stress and increased survival of *Artemia* nauplii (Beux and Zaniboni-Filho 2018). The lower survival of fish fed four times a day and kept in salinized water with 2 g l⁻¹ confirms that the excess of salt together with the excessive feeding frequency

is also harmful to the post-larvae. These results confirm the importance of determining the ideal amount of salt in the water according to the frequency of feeding used.

The post-larvae production cost was significantly higher for higher feeding frequencies. This result was expected, as the labor increases with the frequency of feeding. These results corroborate those found for breeding male *B. splendens*, with fish fed at higher frequencies having the highest production costs (Santos *et al.* 2019). Labor cost results were used to determine the best feeding frequency for *P. brevis*, since the fish length variables allowed the choice of three or four feedings per day. Furthermore, when taking into account the fish weight variables, two, three or four feedings per day can be used. However, labor costs were 23.3% lower with three feedings per day compared to four feedings per day, and 33.4% lower with two feedings per day compared to three feedings per day. Therefore, our results indicate that the best feeding frequency for *P. brevis* post-larvae is three feedings per day, which maximizes growth performance parameters of the post-larvae while minimizing the production costs.

CONCLUSIONS

A feeding frequency of three times a day in salinized water at 1 g L⁻¹ optimized the early post-larval development of *Pyrrhulina brevis*. Our results suggest that, although higher feeding frequencies imply higher labor costs, the cultivation in lower levels of water salinization (in the tested range of 0 to 2 g L⁻¹) and intermediate feeding frequency (in the tested range of one to four bouts per day) may also usefully favor increased survival rates of the post-larvae.

ACKNOWLEDGMENTS

The authors thank the Fish Farming Laboratory (LAPIS) of the Institute of Coastal Studies of Universidade Federal do Pará (UFPA) for providing the infrastructure to conduct the study. The study was funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and the Dean's Office of Research and Graduate Studies of UFPA (PROPEP/UFPA).

REFERENCES

- Abe, H.A.; Dias, A.R.; Reis, R.G.A.; Sousa, N.C.; Ramos, F.M.; Fujimoto, R.Y. 2016. Manejo alimentar e densidade de estocagem na larvicultura do peixe ornamental amazônico *Heros severus*. *Boletim do Instituto de Pesca*, 42: 514-522.
- Abe, H.A.; Dias, J.A.R.; Cordeiro, C.A.M.; Ramos, F.M.; Fujimoto, R.Y. 2015. *Pyrrhulina brevis* (Steindachner, 1876) como uma nova opção para a piscicultura ornamental nacional: larvicultura. *Boletim do Instituto da Pesca*, 41: 113-122.
- Abe, H.A.; Reis, R.G.A.; Paixão, P.E.G.; Fujimoto, R.Y. 2021. Effect of short-term fasting and re-feeding on growth performance of

- larvae and juveniles *Pyrhulina brevis*, an amazon ornamental fish. *Journal of Applied Aquaculture*, 1: 1-13.
- Araújo, L.M.; Junior, L.P.G.; de Souza, W.; Luz, R.K. 2021. Salinity and prey concentration on larviculture of killifish *Hypsolebias radiseriatus* (Cyprinodontiformes: Rivulidae). *Acta Scientiarum. Animal Sciences*, 43: e52075.
- Arias, J.; Rossi, L. 2005. Dieta y estrategia alimentaria de *Pyrhulina australis* (Pisces, Lebiasinidae). *Revista FABICIB*, 9: 197-205.
- Baldisserotto, B.; Mancera, J.M.; Kapoor, B.G. 2007. *Fish osmoregulation*. 1st ed. CRC Press, Boca Raton, 527p.
- Barton, B.A. 2002. Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and Comparative Biology*, 42: 527-525.
- Beux, L.F.; Zaniboni Filho, E. 2018. Influência da baixa salinidade na sobrevivência de náuplios de *Artemia* sp. *Boletim do Instituto de Pesca*, 32: 73-77.
- Boeuf G.; Payan, P. 2001. How should salinity influence fish growth? *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 130: 411-423.
- Campelo, D.A.V.; Dos Reis, R.E.; Dias, B.C.B., Pinheiro Junior, A.D.S.; Dos Santos, L.D.; Brabo, M.F.; Veras, G.C. 2020. Utilização de diferentes alimentos na larvicultura do peixe ornamental amazônico *Acará Severo*/Use of different feeds in larviculture of the amazon ornamental fish *Acará Severo*. *Brazilian Journal of Development*, 6: 14035-14049.
- Cardoso, R.S.; Igarashi, M.A. 2009. Aspectos do agronegócio da produção de peixes ornamentais no Brasil e no mundo. *PUBVET*, 3: 1-22.
- Coraspe Amaral, M.V.; Pedreira, M.M.; Moura, G.S.; Costa, D.C.; Santos, A.E.; Duarte, E. 2017. Combination of diet and water salinity in larviculture of piabanha-do-Pardo (*Brycon vonoi*, Lima 2017). *Revista Ceres*, 64: 384-391.
- Dias, J.A.R.; Abe, H.A.; Sousa, N.C.; Ramos, F.M.; Cordeiro, C.A.M.; Fujimoto, R.Y. 2016. Uso do sal comum (NaCl) e densidade de estocagem durante a larvicultura de *Betta splendens*. *Boletim do Instituto de Pesca*, 42: 719-726.
- Eiras, B.J.; Veras, G.C.; Alves, A.X.; Da Costa, R.M. 2019. Effect of artificial seawater and feeding frequency on the larval culture of freshwater Amazonian ornamental fish banded cichlid *Heros severus* (Heckel, 1840) and angelfish *Pterophyllum scalare* (Schultze, 1823). *Spanish Journal of Agricultural Research*, 17: e0604.
- Fabregat, T.E.H.P.; Damian, J.; Fialho, N.S.; Costa, D., Broggi, J.A.; Pereira, R.G.; Takata, R. 2015. Toxicidade aguda ao sal comum e larvicultura intensiva do jundiá *Rhamdia quelen* em água salobra. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 67: 547-554.
- Fosse, P.J.; Mattos, D.C.; Cardoso, L.D.; Motta, J.H.S.; Jasper, A.P.S.; Radael, M.C.; Andrade, D.R.; Vidal Júnior, M.V. 2013. Estratégia de coalimentação na sobrevivência e no crescimento de larvas de *Betta splendens* durante a transição alimentar. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 65: 1801-1807.
- Furuya, W.M.; Souza, S.R.D.; Furuya, V.R.B.; Hayashi, C.; Ribeiro, R.P. 1998. Dietas peletizada e extrusada para machos revertidos de tilápias do Nilo (*Oreochromis niloticus* L.), na fase de terminação. *Ciência Rural*, 28: 483-487.
- Hayashi, C.; Meurer, F.; Boscolo, W.R.; Lacerda, C.H.F.; Kavata, L.C.B. 2004. Frequência de arraçoamento para alevinos de lambari do rabo-amarelo (*Astyanax bimaculatus*). *Revista Brasileira de Zootecnia*, 33: 21-26.
- Jomori, R.K.; Luz, R.K.; Portella, M.C. 2012. Effect of salinity on larval rearing of pacu, *Piaractus mesopotamicus*, a freshwater species. *Journal of the World Aquaculture Society*, 43: 423-432.
- Jomori, R.K.; Luz, R.K.; Takata, R.; Fabregat, T.E.H.P.; Portella, M.C. 2013. Água levemente salinizada aumenta a eficiência da larvicultura de peixes neotropicais. *Pesquisa Agropecuária Brasileira*, 48: 809-815.
- Kestemont, P.; Xueliang, X.; Hamza, N.; Maboudou, J.; Toko, I.M.; 2007. Effect of weaning age and diet on pikeperch larviculture. *Aquaculture*, 264: 197-204.
- Lee, S.M.; Cho, S.H.; Kim, D.J. 2000. Effects of feeding frequency and dietary energy level on growth and body composition of juvenile flounder, *Paralichthys olivaceus* (Temminck & Schlegel). *Aquaculture Research*, 31: 917-921.
- Luz, R. K.; Dos Santos, J. C. E. 2010. Effect of salt addition and feeding frequency on cascudo preto *Rhinelepis aspera* (Pisces: Loricariidae) larviculture. *Journal of Applied Ichthyology*, 26: 453-455.
- Luz, R.K. 2007. Resistência ao estresse e crescimento de larvas de peixes neotropicais alimentadas com diferentes dietas. *Pesquisa Agropecuária Brasileira*, 42: 65-72.
- Luz, R.K.; Portella, M.C. 2005. Frequência alimentar na larvicultura do trairão (*Hoplias lacerdae*). *Revista Brasileira de Zootecnia*, 34: 1442-1448.
- Luz, R.K.; Portella, M.C. 2015. Effect of prey concentrations and feed training on production of *Hoplias lacerdae* juvenile. *Anais da Academia Brasileira de Ciências*, 87: 1125-1132.
- Luz, R.K.; Santos, A.E.H.; Melillo-Filho, R.; Turra, E.M.; Teixeira, E.A. 2013. Larvicultura de tilápia em água doce e água salinizada. *Pesquisa Agropecuária Brasileira*, 48: 1150-1153.
- Luz, R.K.; Santos, J.C.E. 2008. Densidade de estocagem e salinidade da água na larvicultura do pacamã. *Pesquisa Agropecuária Brasileira*, 43: 903-909.
- Luz, R.K.; Silva, W.S.; Melillo Filho, R.; Santos, A.E.H.; Rodrigues, L.A.; Takata, R.; Alvarenga, E.R.; Turra, E.M. 2012. Stocking density in the larviculture of Nile tilapia in saline water. *Revista Brasileira de Zootecnia*, 41: 2385-2389.
- Mankiw, N.G. 2020. *Principles of Microeconomics*. 9th ed. Cengage Learning, Stamford, 528p.
- Okomoda, V.T.; Aminem, W.; Hassan, A.; Martins, C.O. 2019. Effects of feeding frequency on fry and fingerlings of African catfish *Clarias gariepinus*. *Aquaculture*, 511: e734232.
- Oliveira, L.C.C.; Neto, E.D.A.S.; Junior, A.S.P.; Eiras, B.J.C.F.; Veras, G.C.; de Moura, L.B.; Campelo, D.A.V. 2020. Effect of prey concentrations and salinized water on initial development of *Pyrhulina brevis* (Steindachner, 1876), an Amazonian ornamental fish. *Research, Society and Development*, 9: e381985582.

- Paixao, D.J.M.R.; Brabo, M.F.; Soares, L.M.O.; Campelo, D.A.V.; Veras, G.C. 2019. Optimal feeding frequency for *Heros severus* (Heckel, 1840), an Amazon ornamental fish. *Brazilian Journal of Animal Science*, 48: e20170055
- Portella, M.C.; Verani, J.R.; Cestarolli, M.A. 2000. Use of live and artificial diets enriched with several fatty acid sources to feed *Prochilodus scrofa* larvae and fingerlings. Effects on survival and growth. *Journal of Aquaculture in the Tropics*, 15: 45–58.
- Pouey, J.L.O.F.; Rocha, C.B.; Tavares, R.A.; Portelina, M.K.; Piedras, S.R.N. 2012. Frequência alimentar no crescimento de alevinos de peixe-rei *Odontesthes humensis*. *Ciências Agrárias*, 33: 2423-2427.
- Riche, M.; Haley, D.I.; Oetker, M.; Garbrecht, S.A.R.A.H.; Garling, D.L. 2004. Effect of feeding frequency on gastric evacuation and the return of appetite in tilapia *Oreochromis niloticus* (L.). *Aquaculture*, 234: 657-673.
- Salaro, A.L.; Campelo, D.A.V.; Pontes, M.D.; Tavares, M.M.; Zuanon, J.A.S.; Luz, R.K. 2012. Saline water for juvenile giant trahira during feed training. *Revista Brasileira de Zootecnia*, 41: 1342-1345.
- Sales, A.D.; Veras, G.C.; Brabo, M.F.; Abrunhosa, F.A.; Alves, A.X.; Dias, B.C.B.; Campelo, D.A.V.; Ferreira, M.S.S.; 2016. Fotoperíodo e frequência alimentar na larvicultura do peixe beta. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 68: 1062-1068.
- Sampaio, L.A.; Bianchini, A. 2002. Salinity effects on osmoregulation and growth of the euryaline flounder *Paralichthys orbignianus*. *Journal of Experimental Marine Biology and Ecology*, 269: 187–196.
- Santos, E.L.; da Silva, L.L.A.; Soares, E.C.; de Oliveira, W.D.S.; de Lima, M.R.; de Oliveira, J.M.; Santos, E.L. 2019. Frequência e horário de arraçamento sobre o desempenho de machos *Betta splendens*. *Biota Amazônia*, 9: 30-34.
- Silva, A.P.; Mendes, P.P. 2006. Influência de duas dietas na qualidade de água dos tanques berçário, utilizados no cultivo de camarão marinho *Litopenaeus vannamei* (Boone, 1931). *Acta Scientiarum Animal Sciences*, 28: 105-111.
- Silva, N.C.S.; Costa, A.J.L.; Louvise, J.; Soares, B.E.; Reis, V.C.S.; Albrecht, M.P.; Caramaschi, E.P. 2016. Resource partitioning and ecomorphological variation in two syntopic species of Lebiasinidae (Characiformes) in an Amazonian stream. *Acta Amazonica*, 46: 25-36.
- Silveira, U.S.; Logato, P.V.R.; Pontes, E.C. 2009. Fatores estressantes em peixes. *Revista Eletrônica Nutritime*, 6: 1001-1017.
- Thomassen, J.M.; Fjaera, S.O. 1996. Studies of feeding frequency for atlantic salmon (*Salmo salar*). *Aquacultural Engineering*, 15: 49–157.
- Varsamos, S.; Nebel, C.; Charmantier, G. 2005. Ontogeny of osmoregulation in postembryonic fish: a review. *Comparative Biochemistry and Physiology*, 141: 401-429.
- Weitzman, M.; Weitzman, S.H. 2003. Family Lebiasinidae (Pencil fishes). In: Reis, R.E. (Ed.). *CLOFFSCA-Check list of the freshwater fishes of South and Central America*. Edipucrs, Porto Alegre. p.241-251.
- Xie, F.; Ai, Q.; Mai, K.; XU, W.; Ma, H. 2011. The optimal feeding frequency of large yellow croaker (*Pseudosciaena crocea*, Richardson) larvae. *Aquaculture*, 311: 1-284.
- Zuanon, J.A.S.; Salaro, A.L.; Furuya, W.M. 2011. Produção e nutrição de peixes ornamentais. *Revista Brasileira de Zootecnia*, 40: 165-174.

RECEIVED: 21/02/2022

ACCEPTED: 12/11/2022

ASSOCIATE EDITOR: Rodrigo R. do Valle



This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.