



Complete assignments of NMR data and assessment of trypanocidal activity of new eremantholide C derivatives

DÊNIA A. SAÚDE-GUIMARÃES¹, DÉLIO S. RASLAN²,
EGLER CHIARI³ and ALAÍDE B. DE OLIVEIRA⁴

¹LAPLAMED, DEFAR, Escola de Farmácia, Universidade Federal de Ouro Preto, Campus Morro do Cruzeiro, s/n, Bauxita, 35400-000 Ouro Preto, MG, Brasil

²Departamento de Química, Universidade Federal de Minas Gerais, Av. Antônio Carlos, 6627, Pampulha, 31207-901 Belo Horizonte, MG, Brasil

³Departamento de Parasitologia, ICB, Universidade Federal de Minas Gerais, Av. Antônio Carlos, 6627, Pampulha, 31207-901 Belo Horizonte, MG, Brasil

⁴Departamento de Produtos Farmacêuticos, Faculdade de Farmácia, Universidade Federal de Minas Gerais, Av. Antônio Carlos, 6627, Pampulha, 31270-901 Belo Horizonte, MG, Brasil

Manuscript received on April 4, 2014; accepted for publication on July 8, 2014

ABSTRACT

Chemical transformations of eremantholide C (**1**), a sesquiterpene lactone that was isolated from *Lychnophora trichocarpha* Spreng. led to five new derivatives: 1',2'-epoxyeremantholide C (**2**), 5-n-propylamine-4,5-dihydro-1',2'-epoxyeremantholide C (**3**), 5-n-propylammonium-4,5-dihydro-1',2'-epoxyeremantholide C chloride (**4**), 5-n-propylammonium-4,5-dihydroeremantholide C chloride (**5**) and 16-O-ethyl eremantholide C (**6**). The structures of all these derivatives were assigned on the basis of IR, MS, ¹H and ¹³C NMR data by 1D and 2D techniques. Eremantholide C and the derivatives **2**, **4** and **5** were evaluated against trypomastigotes Y and CL strains of *Trypanosoma cruzi*. Eremantholide C completely inhibited the growth of both the parasites strains while all derivatives were partially active against the CL strain and inactive against the Y strain.

Key words: Eremantholide C derivatives, *Lychnophora trichocarpha*, NMR, sesquiterpene lactones, trypanocidal activity.

INTRODUCTION

Sesquiterpene lactones are chemical markers of certain plant families such as *Asteraceae*. The large number of biological activities experimentally described up to now, raised great interest on this group of substances. The sesquiterpene lactones of the furanheliangolide type are biogenetically derived from heliangolides and are often found in

species of the genus *Lychnophora*, that is native from Brazil (Bohlmann and Jakupovic 1990). In previous studies, eremantholide C and its oxidized derivatives showed activity against *Trypanosoma cruzi* (Oliveira et al. 1996, Saúde-Guimarães et al. 2007). Other biological/pharmacological activities were reported for eremantholide C, such as antibacterial, anti-hyperuricemic, anti-gouty arthritis, anti-inflammatory and antitumor (Barrero et al. 2000, Saúde et al. 2002, de Souza et al. 2012,

Correspondence to: Dênia Antunes Saúde-Guimarães
E-mails: saude@ef.ufop.br / saudeguima@gmail.com

Ferrari et al. 2013, Saúde-Guimarães et al. 2014). Aiming to obtain new bioactive derivatives of eremantholide C (Figure 1), this sesquiterpene lactone was submitted to chemical modifications.

MATERIALS AND METHODS

GENERAL PROCEDURES

Mass spectra were obtained with VG Autospec (electron impact) and HP 5988A (chemical ionization with methane) spectrometers by direct injection (ionization chamber at 200 °C). IR spectra were taken at Galaxy 3000-FTIR spectrophotometer (Mattson Instruments). NMR spectra were taken at Bruker *Avance* DPX (4.7T), DRX (9.4T), equipped with a 5mm dual probe, at 300 K, with TMS as internal reference. One-dimensional ^1H and ^{13}C NMR spectra were acquired under standard conditions, with 90° pulse widths of 8.00 μs and 8.50 μs for ^1H and ^{13}C , respectively. ^1H NMR spectra were obtained using a sweep width of 3 kHz over 32k data points. ^{13}C NMR spectra were obtained

using a sweep width of 31 kHz. DEPT, ^1H , ^1H COSY and ^1H , ^{13}C HETCOR techniques were performed using standard pulse sequences supplied by the spectrometer manufacturer.

SYNTHESIS OF EREMANTHOLIDE C DERIVATIVES 2 TO 6

Eremantholide C (**1**) was obtained from aerial parts of *L. trichocarpa* Spreng. that were collected in Minas Gerais, Brazil, as previously described (Saúde et al. 1998, Ferrari et al. 2013). Its structural characterization is described elsewhere (Le Quesne et al. 1978, Saúde et al. 1998, Saúde-Guimarães et al. 2007).

1',2'-Epoxyeremantholide C (**2**) was prepared by reaction of **1** (2.9 mmol) with *m*-chloroperbenzoic acid (5.3 mmol) in 50 mL of anhydrous CHCl_3 . The mixture was stirred at room temperature for 2 h. The reaction mixture was worked out according to Carda et al. 1986. The resulting residue (1.14 g) was purified by column chromatography (silica gel, hexane: EtOAc 1:1), yielding 970 mg of **2** (93% yield, m.p. 213-217 °C).




Figure 1 - Eremantholide C and derivatives 2-6.

1',2'-Epoxyeremantholide C (2): 6,9-Epoxy-2*H*-1,4-dioxacyclodeca[*c,d*]pentalene-2,7(4*aH*)-dioxane,2*a*,3,5,6,11*a*,11*b*-hexahydro-3-hydroxy-2*a*,6,10-trimethyl-3-(1',2'-epoxy)-2*aR**,3*S**,4*aR**,6*S**,10*Z*,11*aS**,11*bS**). White solid, m.p. 213-217°C. IR (KBr) ν_{\max} (cm⁻¹): 3400 (OH), 1770 (C=O, γ -lactone), 1700 (ketone C=O), 1650 (C=C), 1590 (C=COR, furanone), 1450, 1350, 1300, 1275, 1150, 1100, 1050, 1000, 900. MS (EI), *m/z* (rel. int.): 362 [M⁺, C₁₉H₂₂O₇] (20), 344 (M - H₂O, 17), 234 (19), 205 (19), 189 (20), 177 (29), 165 (32), 149 (25), 138 (55), 122 (53), 95 (**100**), 69 (58), 57 (22).

5-*n*- Propylamine -4,5-dihydro-1',2'-epoxyeremantholide C (**3**) was obtained by reaction of **2** (0.55 mmol) with *n*-propylamine (5.94 mmol), at -18 °C for 16 h (Kirk 1973). After this period, the excess of the amine was removed by evaporation under reduced pressure, at room temperature. The residue obtained was purified by PTLC on silica gel (0.5 mm thickness, eluent - hexane: EtOAc 1:1) resulting in 82 mg (35% yield) of **3** as a white solid. This compound was dissolved in dry THF and treated with gaseous HCl leading to the formation of **4** as a white hydrosoluble solid (quantitative yield).

5-*n*-Propylamine-4,5-dihydro-1',2'-epoxyeremantholide C (3): White solid, m.p. 183-189 °C. IR (KBr) ν_{\max} (cm⁻¹): 3400 (OH), 1770 (C=O, γ -lactone), 1700 (C=O, ketone), 1590 (C=COR, furanone), 1450, 1380, 1300, 1200, 1150, 1100, 1050, 1000, 900, 850. MS (EI), *m/z* (rel. int.): 421 (M⁺, C₂₂H₃₁O₇N, **99**), 403 (M - H₂O, 28), 238 (25), 195 (64), 168 (31), 151 (25), 138 (33), 125 (**100**), 99 (23), 72 (35).

5-*n*-Propylammonium-4,5-dihydro-1',2'-epoxyeremantholide C chloride (4): white solid, m.p. 176-179 °C. MS (EI), *m/z* (rel. int.): 457 (M⁺, C₂₂H₃₂O₇NCl, 72), 439 (M - H₂O, 11), 390 (18), 364 (31), 336 (21), 318 (59), 195 (79), 168 (44), 152 (19), 138 (45), 125 (**100**), 99 (45), 81 (18), 72 (46).

5-*n*-Propylammonium-4,5-dihydroeremantholide C chloride (**5**) was obtained from the reaction of **1** (0.58 mmol) with *n*-propylamine (5.94 mmol),

at -18 °C for 10 h (Kirk 1973). After this period, the excess of amine was removed by evaporation under reduced pressure, at room temperature. The residue obtained was purified by PTLC on silica gel (0.5 mm thickness, eluent - hexane: EtOAc 1:1) affording 120 mg of a yellowish pasty material. Silica TLC of this material showed three spots when revealed by iodine. After development with ninhydrin, the TLC showed only a rosy spot, characteristic of amines. This material was then dissolved in dry THF and treated with gaseous HCl to give the hydrochloride **5** as a white hydrosoluble solid (70 mg).

5-*n*-Propylammonium-4,5-dihydroeremantholide C chloride (5): White solid, m.p. 195-197 °C. MS (EI), *m/z* (rel. int.): 405 (60), 387 (M - H₂O, 5), 195 (40), 168 (32), 135 (21), 125 (**100**), 69 (35).

16-*O*-Ethyleremantholide C (**6**) was obtained, according to the methodology described by Partwardhan et al. 1974, by reacting **1** (0.28 mmol) with 0.08 mL of triethyl orthoformate and 25 mg of Amberlyst resin 15. The reaction mixture was stirred at room temperature for 3 days. Then it was neutralized with aqueous K₂CO₃ solution and filtered. The product was extracted with diethyl ether. The organic layer was washed with water, filtered on anhydrous sodium sulfate and concentrated under reduced pressure, yielding 72 mg of ether **6** as a white solid (66% yield).

16-*O*-Ethyleremantholide C (6): White solid. IR (KBr) ν_{\max} (cm⁻¹): 1770 (C=O, γ -lactone), 1700 (C=O, ketone), 1660 (C=C), 1590 (C=COR, furanone), 1450, 1380, 1300, 1280, 1260, 1210, 1200, 1150, 1140, 1100, 1060, 1000, 870, 750. MS (EI), *m/z* (rel. int.): 375 (M+1)⁺, C₂₁H₂₆O₆, 3), 329 (54), 285 (17), 199 (9), 165 (49), 95 (85), 69 (**100**), 67 (18), 45 (23), 43 (39).

IN VITRO ASSAYS WITH *TRYPANOSOMA CRUZI*

TRYPOMASTIGOTES

Albino mice infected with the Y or CL strains of *T. cruzi* were used to provide trypomastigotes.

Samples of eremantholide **1** and of the derivatives **2**, **4** and **6** were dissolved or suspended in dimethyl sulfoxide (DMSO) (0.2 mL) and plus Krebs-Ringer-glucose (2.0 mL) and mixed with an equal volume of parasitized whole blood diluted in bovine calf serum. A parasite density of 2×10^5 trypomastigotes/0.1 mL was calculated for each flat-bottomed test tube (4 mL, 56 x 13 mm); control tubes without the test extracts were included. After incubation at 4 °C for 24 h the suspensions were examined microscopically. Only those samples that killed 100% of the parasites were considered active. Samples that inhibited 50% of parasite growth compared to control were considered partially active.

RESULTS AND DISCUSSION

Data of the ^1H and ^{13}C NMR spectra of derivatives **2-6** are given at on Tables I and II, respectively. The chemical shifts were assigned by consideration of known substituent effects of the groups concerned and with the aid of both ^1H , ^1H COSY and ^1H , ^{13}C HETCOR contour maps. The stereochemistry of the carbons were determined based on coupling constants and ^1H , ^1H NOESY contour maps.

From the reaction of **1** with *m*-chloroperbenzoic acid, the main product **2** was obtained and it was characterized by ^1H NMR data. The presence of an epoxy group in **2** was highlighted in the ^1H NMR spectrum by the two doublets at $\delta 3.11$ ($J = 5.5$ Hz) and $\delta 2.70$ ($J = 5.5$ Hz), attributed to H-2'a and H-2'b, which appear in the ^1H NMR spectrum of **1** at $\delta 5.31$ (br s) and $\delta 5.07$ (t, $J = 1.6$ Hz), respectively.

The signals at $\delta 130.00$ and $\delta 115.80$, attributed to the olefinic carbons C-1' and C-2', respectively, in the ^{13}C NMR spectrum of **1**, are replaced in the spectrum of **2** by the signals at $\delta 59.50$ and $\delta 58.47$, typical chemical shifts of oxygenated carbons with hybridization sp^3 . In addition, the signal attributed to C-3 in the spectrum of **1** ($\delta 19.00$) is shifted to $\delta 11.44$ in the spectrum of **2**, confirming the epoxidation of the Δ^1 double bond of **1**.

The mass spectrum of **2** showed the molecular ion peak at m/z 362, which represents an increment of 16 mass units to the molar mass of **1**, corresponding to the addition of an oxygen atom. This mass is consistent with the molecular formula $\text{C}_{19}\text{H}_{22}\text{O}_7$.

Data of IR, MS, ^1H NMR and ^{13}C NMR of **2**, are in agreement with the new derivative 1',2'-epoxyeremantholide **2**.

Epoxide **2** was reacted with methylamine, cyclohexylamine, diethylamine and *n*-propylamine. Reactions with the first three amines led to mixtures of many products, in such a way that no derivative with suitable purity was obtained. The reaction of **2** with *n*-propylamine afforded a less complex mixture of products which, after separation by column chromatography on silica gel, led to the amino product **3** with a 35% yield. The amine **3** was transformed into its hydrochloride salt by reaction with gaseous HCl in dry THF, leading to the hydrosoluble compound **4**.

The multiplets in $\delta 5.02$ - 4.98 and $\delta 6.03$ - 6.00 attributed, respectively, to H-6 and H-5, in the ^1H NMR spectrum of **2** appear in the spectrum of **3** at $\delta 4.26$ (d, $J = 6.5$ Hz) and at $\delta 2.99$ (brs), respectively. The quartet at $\delta 2.89$ ($J = 7.2$ Hz), that appears in the ^1H NMR spectrum of **2**, was assigned to H-4. A correlation of this signal with the doublets at $\delta 1.37$ ($J = 7.2$ Hz), assigned to H-15, is indicated by the values of the coupling constants and by the COSY ^1H - ^1H contour map. The signals at $\delta 134.27$ and $\delta 130.24$ which are attributed to the olefinic carbons C-5 and C-4, respectively, as well as the signal attributed to C-15 ($\delta 20.39$) in the ^{13}C NMR spectrum of **2**, are shifted in the ^{13}C NMR spectrum of **3** to $\delta 65.24$, $\delta 38.20$ and $\delta 16.70$, respectively, allowing the proposition that the amine group was linked to the C-5 of compound **2**.

The double triplets at $\delta 2.80$ ($J = 7.0$ and 11.2 Hz) and $\delta 2.53$ (6.9 and 11.2 Hz) assigned to H-4'a and b, respectively, the multiplet at $\delta 1.50$ assigned to H-5', and the triplet at $\delta 0.93$ ($J = 7.3$ Hz) assigned to the atoms H-6' of the methyl group,

TABLE I
¹H NMR data for compounds 1, 2, 3, 4, 5 and 6, δ, J (Hz).

H	1*,1	2**,1	3**,1	4**,2	5**,2	6**,1
2	5.63 s	5.62 s	5.56 s	5.99 s	5.93 s	5.62 s
4	-	-	2.89 q J _{4,15} = 7.2	3.56 q J _{4,15} = 7.2	3.45 q J _{4,15} = 7.2	-
5	6.04-6.03 m	6.03-6.00 m	2.99 brs	3.97 brs	3.88 brs	6.03-6.01 m
6	5.02-4.98 m	5.02-4.97 m	4.26 d J _{6,7} = 6.5	4.74 a	4.73 d J _{6,7} = 7.9	4.90-4.86 m
7	2.82 dd J _{6,7} = 7.1; J _{7,8} = 4.2	2.87 dd J _{6,7} = 7.3; J _{7,8} = 4.2	2.95 dd J _{6,7} = 6.5; J _{7,8} = 4.9	3.09 dd J _{6,7} = 7.8; J _{7,8} = 4.2	3.04 dd J _{6,7} = 7.8; J _{7,8} = 4.3	2.84 dd J _{6,7} = 7.2; J _{7,8} = 4.4
8	4.09 ddd J _{7,8} = 4.2; J _{8,9a} = 2.5; J _{8,9b} = 12.0	4.07 ddd J _{7,8} = 4.2; J _{8,9a} = 2.6; J _{8,9b} = 11.9	3.91 ddd J _{7,8} = 4.9; J _{8,9a} = 2.3; J _{8,9b} = 11.9	3.92 m	3.96 ddd J _{7,8} = 4.3; J _{8,9a} = 2.8; J _{8,9b} = 11.3	3.85 ddd J _{7,8} = 4.4; J _{8,9a} = 2.4; J _{8,9b} = 12.0
9a	2.48 dd J _{8,9a} = 2.5; J _{9a,9b} = 13.5	2.39 dd J _{8,9a} = 2.6; J _{9a,9b} = 13.6	2.39 dd J _{8,9a} = 2.3; J _{9a,9b} = 13.5	2.32 dd J _{8,9a} = 2.3; J _{9a,9b} = 13.6	2.27 dd J _{8,9a} = 2.8; J _{9a,9b} = 13.8	2.41 dd J _{8,9a} = 2.4; J _{9a,9b} = 13.6
9b	2.00 dd J _{8,9b} = 12.0; J _{9a,9b} = 13.5	1.94 dd J _{8,9b} = 11.9; J _{9a,9b} = 13.6	1.90 dd J _{8,9b} = 11.9; J _{9a,9b} = 13.5	2.39 J _{8,9b} = 11.5; J _{9a,9b} = 13.6	2.34 dd J _{8,9b} = 11.3; J _{9a,9b} = 13.8	2.06 dd J _{8,9b} = 12.0; J _{9a,9b} = 13.6
13	1.18 s	1.33 s	1.38 s	1.39 s	1.21 s	1.16 s
14	1.45 s	1.48 s	1.43 s	1.50 s	1.45 s	1.50 s
15	2.05 t J _{5,15} = 6,15 = 1.9	2.05 t J _{5,15} = 6,15 = 2.0	1.37 d J _{4,15} = 7.2	1.52 d J _{4,15} = 7.2	1.48 d J _{4,15} = 7.2	2.06 dd J _{5,15} = 2.0; J _{6,15} = 1.6
2'a	5.31 brs	3.11 d J _{2'a,2'b} = 5.5	3.12 d J _{2'a,2'b} = 5.5	4.04 d J _{2'a,2'b} = 11.9	5.22 brs	5.28 d J _{2'a,2'b} = 1.2
2'b	5.07 t J _{2'a,2'b} = 1.6	2.70 d J _{2'a,2'b} = 5.5	2.71 d J _{2'a,2'b} = 5.5	3.75 d J _{2'a,2'b} = 11.9	5.13 t J _{2'a,2'b} = 1.4	5.13 brs
3'	1.91 s	1.55 s	1.57 s	1.50 s	1.77 s	1.79 s
4'a	-	-	2.8 dt J _{4'a,4'b} = 11.2; J _{4'a,5'} = 7.0	3.28 td J _{4'a,4'b} = 11.2; J _{4'a,5'} = 5.8;	3.18 dt J _{4'a,4'b} = 10.5; J _{4'a,5'} = 5.7	3.36 dq J _{4'a,4'b} = 9.2; J _{4'a,5'} = 7.2
4'b	-	-	2.53 dt J _{4'a,4'b} = 11.2; J _{4'b,5'} = 6.9;	3.17 td J _{4'a,4'b} = 11.2; J _{4'b,5'} = 5.8	3.09 dt J _{4'a,4'b} = 10.5; J _{4'b,5'} = 5.7	3.20 m
5'	-	-	1.50 m	1.80 m	1.75 m	1.07 t J _{4'a,5'} = 4,4b,5' = 7.2
6'	-	-	0.93 t J _{5',6'} = 7.3	0.99 t J _{5',6'} = 7.4	0.94 t J _{5',6'} = 7.3	-
OH	3.79 s	3.69 s				

x – Superimposed signal with D₂O. * - 300 MHz; ** - 400 MHz, 1 - CDCl₃, 2 - D₂O.

indicated the presence of a propylamine group at **3**. This group was confirmed by the ^{13}C NMR signals at $\delta 50.51$, $\delta 17.52$ and $\delta 11.65$, attributed to carbons 4', 5' and 6', respectively.

TABLE II
 ^{13}C NMR data (δ) for **1**, **2**, **3**, **4**, **5** and **6**.

C	1 ^{*.1}	2 ^{**.1}	3 ^{**.1}	4 ^{**.2}	5 ^{**.2}	6 ^{**.1}
C-1	205.89	205.23	204.53	209.06	209.34	205.24
C-2	104.54	104.50	104.03	105.93	106.08	104.45
C-3	187.27	186.64	192.93	192.39	192.53	186.91
C-4	130.00	130.15	38.14	33.44	33.68	139.10
C-5	134.77	134.28	57.18	58.65	57.30	134.75
C-6	81.46	81.34	82.42	80.77	81.00	81.75
C-7	62.53	63.06	65.24	63.79	63.99	63.10
C-8	78.37	78.38	77.45	76.47	77.05	78.43
C-9	43.46	43.25	43.84	42.38	42.60	43.89
C-10	90.24	89.95	89.88	91.88	91.99	89.98
C-11	59.88	58.43	58.51	61.65	60.61	60.00
C-12	175.72	174.70	175.09	177.21	178.19	175.76
C-13	21.94	20.64	21.16	21.52	20.83	21.66
C-14	20.48	20.39	20.51	20.16	20.22	20.64
C-15	20.30	20.33	16.70	15.06	15.17	20.49
C-16	106.09	104.57	104.94	107.06	107.27	109.59
C-1'	142.22	59.60	59.71	76.67	141.39	139.28
C-2'	115.80	53.59	53.80	50.06	116.63	117.61
C-3'	19.00	11.44	17.55	20.07	19.19	18.46
C-4'	-	-	50.51	49.11	49.22	57.26
C-5'	-	-	17.52	19.04	18.70	15.29
C-6'	-	-	11.65	10.52	10.62	-

* - 75 MHz; ** - 100 MHz; 1 - CDCl_3 ; 2 - D_2O .

The two doublets at $\delta 2.71$ ($J = 5.5$ Hz) and $\delta 3.12$ ($J = 5.5$ Hz) assigned to H-2'a and b, respectively, and the signals at $\delta 59.71$ and $\delta 53.80$ on the ^{13}C NMR spectrum, due to carbons 1' and 2', respectively, showed that the epoxy group remained unchanged in the molecule of **3**.

The assignments made to the signals in the ^1H NMR spectrum of **3** were confirmed by the values of the coupling constants and by the ^1H - ^1H COSY contour map which showed the following correlations: H-4 and H-15; H-4'a, H-4'b and H-5'; H-5' and H-6'; H-6 and H-7, and H-2'a and b. Chemical shifts for ^{13}C NMR spectrum of **3** were assigned by comparison with the spectrum of **2** and based on the DEPT spectrum.

The mass spectrum of **3** presented the molecular ion peak at m/z 421 u, corresponding to 59 mass units higher than the molecular ion of **2**, and it corresponds to n-propylamine group ($\text{C}_3\text{H}_7\text{NH}_2$). The molar mass of **3** was compatible with the molecular formula $\text{C}_{22}\text{H}_{31}\text{O}_7\text{N}$.

Based on data retrieved from IR, MS, ^1H and ^{13}C NMR it was concluded that the product obtained was that resulting from the amine addition to carbon-5 of the epoxide **2**, through a Michael's reaction, generating the novel derivative 5-n-propylamine-1',2'-epoxyeremantholide (**3**). This might be compared to the proposed mechanism for the antitumor activity of the eremantholides by reaction with biological nucleophiles (Mc Dougal et al. 1989).

The shifting of the signals due to the hydrogen and carbon atoms 4, 5, 15, 4'a, 4'b, 5' e 6' of compounds **3** and **4** could be observed by comparing their ^1H and ^{13}C NMR spectra.

The signals relative to H-4, H-5, H-15, H-4'a and H-4'b, H-5' and H-6' appear in the spectrum of **3**, respectively, at $\delta 2.89$ (q, $J = 7.2$ Hz), $\delta 2.99$ (brs), $\delta 1.37$ (d, $J = 7.2$ Hz), $\delta 2.80$ (dt, $J = 7.0$ and 11.2 Hz), $\delta 2.53$ (dt, $J = 6.9$ and 11.2 Hz), $\delta 1.50$ (m) and $\delta 0.93$ (t, $J = 7.3$ Hz), while in the spectrum of **4** they are shown at $\delta 3.56$ (q, $J = 7.2$ Hz), $\delta 3.97$ (brs), $\delta 1.52$ (d $J = 7.2$ Hz), $\delta 3.28$ (td, $J = 5.8$ and 11.2 Hz), $\delta 3.17$ (td, $J = 5.8$ and 11.2 Hz), $\delta 1.80$ (m) and $\delta 0.99$ (t, $J = 7.4$ Hz), respectively.

In the ^{13}C NMR spectrum of **3**, the signals at $\delta 38.20$, $\delta 57.18$, $\delta 16.70$, $\delta 50.51$, $\delta 17.52$ and $\delta 11.65$ were assigned to C-4, C-5, C-15, C-4', C-5' and C-6', respectively. These signals were observed at $\delta 33.44$, $\delta 58.65$, $\delta 15.06$, $\delta 49.11$, $\delta 19.04$ and $\delta 10.52$, respectively, in the spectrum of **4**.

The signal corresponding to H-6 in the ^1H NMR spectra of **4** overlapped that of the solvent (D_2O) at $\delta 4.74$. This was confirmed by the ^1H - ^1H COSY contour map of **4**, evidencing correlations between the signal at $\delta 4.74$ (brs), and the signals attributed to H-7 ($\delta 3.09$, dd, $J = 4.2$ and 7.8 Hz),

and to H-5 (δ 3.97, brs). The ^1H - ^1H COSY also evidenced correlations between the following hydrogen atoms: H-4 and H-15; H-8, H-9a and H-9b; H-9a and H-9b; H-4'a, H-4'b and H-5'; H-5' and H-6'; H-2'a, H-2'b and H-3'.

The relative configuration of the n-propylamine and the C-15-methyl group was defined on the basis of the ^1H , ^1H NOESY (Figure 2), that

indicated correlation between the H-5 signal at δ 3.97 (brs), with the signals at δ 4.74 and δ 3.56 (q, $J = 7.2$ Hz), attributed to H-6 and H-4, respectively. This correlation showed a *-cis* relationship between the H-4 and H-5 atoms with a β configuration. Consequently, the C-15-methyl group and the n-propylamine group should have an α configuration.




Figure 2 - Arrows represent NOESY correlations.

The mass spectrum of **4** presented a molecular ion peak m/z 457 u, compatible with a molecular formula $\text{C}_{22}\text{H}_{32}\text{O}_7\text{NCl}$. Such a molecular mass represents an increment of 36 units compared with the mass of **3**, compatible with the addition of one HCl per molecule of **4**.

Based on data from IR, MS, ^1H and ^{13}C NMR, COSY and NOESY it was concluded that compound **4** is 5-n-propylammonium-4,5-dihydro-1',2'-epoxyeremantholide C chloride, a novel compound that is firstly described in this paper.

In order to confirm the positioning of the n-propylamine group at the C-5 of compound **2**, the reaction of **1** with n-propylamine was carried out. Three spots were observed when a silica gel TLC plate of the reaction mixture was sprayed with iodine. Under ninhydrin, only a rosy spot typical of amine, was observed. Then, the mixture was treated with gaseous HCl in dry THF yielding **5** as a white hydrosoluble solid.

In the ^1H NMR spectrum of **5**, the signals at δ 3.18 (td, $J = 5.7$ and 10.5 Hz), δ 3.09 (td, $J = 5.7$ and 10.5 Hz), δ 1.75 (m) and δ 0.94 (t, $J = 7.3$ Hz) were

attributed to H-4'a, H-4'b, H-5' and H-6' of the n-propylammonium group, respectively. These data were further confirmed by the signals at δ 49.22 and δ 18.70, together with the signal at δ 10.62 in the ^{13}C NMR of **5**, which were attributed, respectively, to C-4', C-5' and C-6' of the n-propylammonium group.

The ^1H NMR spectrum of **5** presented a quartet at δ 3.45 (1H, $J = 7.2$ Hz) that was attributed to H-4. The signals for H-5 and H-15 in this spectrum appeared as a broad singlet at δ 3.88 and a doublet at δ 1.48 ($J=7.2$ Hz), respectively. In the ^1H NMR spectrum of **1** these atoms are represented by signals at δ 6.04–6.03 (m) and δ 2.05 (t, $J = 1.9$ Hz), respectively. Values of coupling constants for H-4 and H-15 indicate that these hydrogens are in vicinal positions.

The signal attributed to H-6 in the ^1H NMR spectra of **5** is probably superimposed to that of H_2O in the solvent (D_2O), which appear at δ 4.74. In order to shift the signal of H_2O and observe the one from H-6, the ^1H NMR spectra of **5** in D_2O was run at 330 K, when the signal due to the solvent was shifted to δ 4.72 and the doublet attributed to H-6 was seen at δ 5.07 ($J = 7.9$ Hz).

The stereochemistry of the n-propylammonium and of the C-15 methyl groups were inferred based on the NOESY technique (Figure 2), which indicated a correlation between the signal at δ 3.88 (brs), attributed to H-5, and the signals at δ 4.73 (d, $J = 7.9$ Hz) and δ 3.45 (q, $J = 7.2$ Hz), corresponding to H-6 and H-4, respectively. These data suggest that H-4 and H-5 have a *cis* relationship and a β configuration. Consequently, the C-15 methyl and the n-propylammonium groups have a α configuration.

The molecular ion peak m/z 441 u, corresponding to the molecular formula $C_{22}H_{32}O_6NCl$, was not observed at the mass spectrum of **5**, but it showed a peak at m/z 405 u that was compatible to the loss of a molecule of HCl from the molecular ion.

Data of IR, MS, 1H and ^{13}C NMR spectra, and the COSY and NOESY contour maps are in agreement with the new derivative 5-n-propylammonium-4,5-dihydroeremantolide C chloride, a new derivative of eremantolide C.

The infrared spectrum of **6** showed no absorption characteristic of the hydroxyl group. The 1H NMR spectrum presented signals at δ 3.36 (dq, $J = 7.2$ and 9.2 Hz), δ 3.20 (m) and δ 1.07 (t, $J = 7.2$ Hz) corresponding to H-4'a, H-4'b and H-5', respectively, and that indicates the presence of an ethoxyl group. The COSY 1H , 1H contour map evidenced correlations between H-4'a, H-4'b and H-5' atoms.

The ^{13}C NMR spectrum of **6** shows a signal at δ 57.26, characteristic for a sp^3 carbon bonded to oxygen and, therefore, it was attributed to C-4'. The signal at δ 15.29 was assigned to the methyl carbon at the 5' position. These data confirmed the presence of an ethoxyl group in the molecule of **6**.

In the mass spectrum of **6** the ion $[M+1]^+$ appeared at m/z 375 u corresponding to an increase of 29 mass units in comparison with **1**, that is compatible with the presence of an ethyl group, and corresponding to the molecular formula $C_{21}H_{26}O_6$.

Assignments of the signals in the 1H NMR spectrum of **6** were confirmed through the corre-

lations observed at the 1H , 1H COSY contour map.

Based on the modifications observed in the IR, MS, 1H and ^{13}C NMR spectra it was concluded that the compound obtained was the new derivative 16-*O*-ethyleremantolide C (**6**).

Eremantolide C derivatives **2**, **4** and **5** were tested against trypomastigote forms of Y and CL strains of *Trypanosoma cruzi*, the infectious agent of Chagas' disease, in comparison with Violet crystal (active at 125 $\mu g \cdot mL^{-1}$) that was used as reference in the *in vitro* tests. The results are shown on Table III. Eremantolide C and the derivatives **2**, **4** and **5** were evaluated against Y and CL strains of *T. cruzi*. Eremantolide C completely inhibited the growth of both the parasite strains in the concentrations of 3,600 $\mu g/mL$ (Y strain) and 1,800 $\mu g/mL$ (CL strain), respectively, while all the derivatives were partially active against the CL strain and inactive against the Y strain, in the concentrations assayed.

TABLE III
Results of *in vitro* tests of compounds **1**, **2**, **4** and **5** against Y and CL trypomastigote strains of *T. cruzi*. Inhibition of growth (%).

Compound	Y Strain ($\mu g/mL$)	CL Strain ($\mu g/mL$)
1	3600 (100%)	1800 (100%)
2	3290 (NI)	823 (50%)
4	4155 (NI)	1039 (50%)
5	4009 (NI)	1002 (50%)

NI - Growth was not inhibited.

CONCLUSIONS

The present paper describes the synthesis of five new eremantolide C derivatives (**2-6**) that were spectroscopically characterized and had their activity evaluated *in vitro* against trypomastigotes of Y and CL strains of *T. cruzi*. All the eremantolide C derivatives tested (**2**, **4** and **5**) showed 50% growth inhibition of the CL strain.

ACKNOWLEDGMENTS

The authors acknowledge Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for financial support and fellowships.

RESUMO

Transformações químicas realizadas em eremantolida C (1), uma lactona sesquiterpênica isolada de *Lychnophora trichocarpha* Spreng. originaram cinco novos derivados: 1',2'-epoxieremantolida C (2), 5-n-propilamino-4,5-diidro-1',2'-epoxieremantolida C (3), cloreto de 5-n-propilamônio-4,5-diidro-1',2'-epoxieremantolida C (4), cloreto de 5-n-propilamônio-4,5-diidroeremantolida C (5) e 16-O-etileremantolida C (6). As estruturas químicas de todos estes derivados foram elucidadas com base nos espectros de IV, EM, RMN de ¹H e de ¹³C e por meio de técnicas 1D e 2D. Eremantolida C e os derivados 2, 4 e 5 foram avaliados frente a cepas tripomastigotas Y e CL de *Trypanosoma cruzi*. Eremantolida C inibiu completamente o crescimento de ambas as cepas de parasitas, enquanto todos os derivados foram parcialmente ativos contra a cepa CL e inativos contra a cepa Y.

Palavras-chave: Derivados de eremantolida C, *Lychnophora trichocarpha*, RMN, lactonas sesquiterpênicas, atividade anti-*Trypanosoma cruzi*.

REFERENCES

- BARRERO AF, OLTRA JE, ÁLVAREZ M, RASLAN DS, SAÚDE DA AND AKSSIRA M. 2000. New sources and antifungal activity of sesquiterpene lactones. *Fitoterapia* 71: 60-64.
- BOHLMANN F AND JAKUPOVIC J. 1990. Progress in the chemistry of the *Vernoniae*. *Pl Syst Ecol* 4: 3-43.
- CARDA M, ARNÓ M AND MARCO JA. 1986. Total synthesis of Rothin -A and Rothin-B. *Tetrahedron* 42: 3655-3662.
- DE SOUZA MR, DE PAULA CA, RESENDE MLP, GRABE-GUIMARÃES A, DE SOUZA FILHO JD AND SAÚDE-GUIMARÃES DA. 2012. Pharmacological basis for use of *Lychnophora trichocarpha* in gouty arthritis: anti-hyperuricemic and anti-inflammatory effects of its extract, fraction and constituents. *J Ethnopharmacol* 142: 845-850.
- FERRARI FC, FERREIRA LC, SOUZA MR, GRABE-GUIMARÃES A, PAULA CA, REZENDE SA AND SAÚDE-GUIMARÃES D A. 2013. Anti-Inflammatory Sesquiterpene Lactones from Spreng. (Brazilian Arnica). *PTR Phytother Res* 27: 384-389.
- KIRK DN. 1973. Selectivity in reactions of epoxides. *Chem Ind* 3:109-112.
- LE QUESNE PW, LEVERY SB, MENACHERY MD, BRENNAN TF AND RAFFAUF RF. 1978. Novel Modified Germacranolides and Other Constituents of *Eremanthus elaeagnus* Schultz-Bip. (Compositae). *J Chem Soc Perkin Trans I* 12:1572-1580.
- MC DOUGAL PG, OH YI AND VAN DERVEER DJ. 1989. Synthesis of the Furanoheliangolide Ring Skeleton. *J Org Chem* 54: 91-97.
- OLIVEIRA AB, SAÚDE DA, PERRY KSP, DUARTE DS, RASLAN DS, BOAVENTURA MAD AND CHIARI E. 1996. Trypanocidal sesquiterpenes from *Lychnophora* species. *Phytother Res* 10: 292-295.
- PATWARDHAN AS AND DEV S. 1974. Amberlyst-15, a superior catalyst for the preparation of enol ethers and acetals. *Synthesis* 5: 348-349.
- SAÚDE DA, BARRERO AF, OLTRA JE, JUSTICIA J, RASLAN DS AND SILVA EA. 2002. Atividade Antibacteriana de Furanoheliangolidos. *Rev Bras Farmacogn* 12: 7-10.
- SAÚDE DA, RASLAN DS, DE SOUZA FILHO JD AND DE OLIVEIRA AB. 1998. Constituents from the aerial parts of *Lychnophora trichocarpha*. *Fitoterapia* 69: 90-91.
- SAÚDE-GUIMARÃES DA, PERRY KSP, RASLAN DS, CHIARI E, BARRERO AF AND OLTRA JE. 2007. Complete assignments of ¹H and ¹³C NMR data for trypanocidal eremantholide C oxide derivatives. *Magn Reson Chem* 45: 1084-1087.
- SAÚDE-GUIMARÃES DA, RASLAN DS AND OLIVEIRA AB. 2014. *In vitro* antitumor activity of sesquiterpene lactones from *Lychnophora trichocarpha*. *Rev Bras Plant Med* 16: 275-282.

