



---

## Edition of March 2014, vol. 102 (3), Supl. 1, pg. 1-61

The “Brazilian Society of Cardiology Guidelines on Unstable Angina and Non-ST-Segment Elevation Myocardial Infarction (2nd Edition, 2007) – 2013 Updating”, published as supplement number 3 of the March 2014 edition of the *Arquivos Brasileiros de Cardiologia* [Arq Bras Cardiol. 2014;102(3Supl.1):1-61], has undergone the following corrections:

On page 46, item “Lipid Approach – Summary of Recommendations and Evidence”, sub-item “Class 1 recommendations”, the following text has been considered second topic:

For patients with unstable myocardial ischemic syndromes with no ST-segment elevation and LDL-C  $\geq$  100 mg/dL, statins should be used if no contraindications apply, aiming at LDL-C < 70 mg/dL (level of evidence: A). If tolerated, statin should be maintained indefinitely.

On page 46, item “Lipid Approach – Summary of Recommendations and Evidence”, sub-item “Class 1 recommendations”, the third topic has been inserted:

If the patient is already on statin, the drug should not be suspended regardless of the LDL-C level (level of evidence: B).

On page 46, item “Lipid Approach – Summary of Recommendations and Evidence”, sub-item “Class 2 recommendations” has been removed.