

Smoking is Associated with Remodeling of Gap Junction in the Rat Heart: Smoker's Paradox Explanation?

Rosângela Novo¹, Cristiana M. Freire¹, Sergio Felisbino², Marcos F. Minicucci, Paula S. Azevedo¹, Leonardo A. M. Zornoff¹, Sergio A. R. Paiva¹

Faculdade de Medicina de Botucatu¹; Instituto de Biociências de Botucatu², Botucatu, SP – Brazil

Abstract

Background: In a previous study utilizing the rat model, exposure to tobacco smoke for 5 weeks increased survival after AMI, despite similar age and infarct size between the smokers and nonsmokers, and absence of reperfusion.

Objective: Thus, this study aimed to analyze the effects of exposure to tobacco smoke on intensity, distribution or phosphorylation of connexin 43 in the rat heart.

Methods: Wistar rats weighing 100 g were randomly allocated into 2 groups: 1) Control (n = 25); 2) Exposed to tobacco smoke (ETS), n = 23. After 5 weeks, left ventricular morphometric analysis, immunohistochemistry and western blotting for connexin 43 (Cx43) were performed.

Results: Collagen volume fraction, cross-sectional areas, and ventricular weight were not statistically different between control and ETS. ETS showed lower stain intensity of Cx43 at intercalated disks (Control: 2.32 ± 0.19 ; ETS: 1.73 ± 0.18 ; $p = 0.04$). The distribution of CX43 at intercalated disks did not differ between the groups (Control: 3.73 ± 0.12 ; ETS: 3.20 ± 0.17 ; $p = 0.18$). ETS rats showed higher levels of dephosphorylated form of Cx43 (Control: 0.45 ± 0.11 ; ETS: 0.90 ± 0.11 ; $p = 0.03$). On the other hand, total Cx43 did not differ between control and ETS groups (Control: 0.75 ± 0.19 ; ETS: 0.93 ± 0.27 ; $p = 0.58$).

Conclusion: Exposure to tobacco smoke resulted in cardiac gap junction remodeling, characterized by alterations in the quantity and phosphorylation of the Cx43, in rats hearts. This finding could explain the smoker's paradox observed in some studies (Arq Bras Cardiol. 2013;100(3):274-280).

Keywords: Smoking/physiopathology; Smoke Inhalation Injury/mortality; Arrhythmias, Cardiac; Rats.

Introduction

Despite convincing data on the harmful effects of smoking, several studies have reported low mortality after acute myocardial infarction (AMI) for smokers compared with nonsmokers. This phenomenon has been referred as the smoker's paradox¹⁻⁴.

Many explanations have been proposed for the more favorable outcomes in smokers after AMI including: 1) smokers are younger and present fewer co-morbidities; 2) they have higher pre-hospital mortality; 3) smokers have a larger thrombus burden, inducing a greater effectiveness of intravenous thrombolysis; 4) ischemic preconditioning-like effect is induced by tobacco smoke. Ischemic preconditioning is a phenomenon

whereby a brief period of ischemia renders the myocardium resistant to infarction from a subsequent ischemic insult. Ischemic preconditioning limits infarct size, reduces the incidence of fatal arrhythmias, and protects against stunning myocardium^{5,6}.

In a previous study using the rat model, exposure to tobacco smoke for 5 weeks increased survival after AMI, despite similar age and infarct size between the smokers and nonsmokers, and absence of reperfusion. Importantly, the reduction of mortality was observed within the initial 24 hours after coronary occlusion⁷. In this model, 87% of early deaths were a consequence of ventricular arrhythmias⁸. Therefore, we assumed that the smoker's paradox in this model could be a result of a preconditioning-like effect. However, the potential mechanisms of this protection induced by exposure to tobacco smoke are still unknown.

Ischemic preconditioning protection depends on functional channels of gap junction intercellular communication⁹. The gap junctions in myocardium are specialized intercellular contacts that allow electrical impulse propagation among cardiomyocytes. Connexin 43 (Cx43) is the principal gap junction protein in ventricles of the heart¹⁰ and plays a major role in ischemic preconditioning. This fact is evidenced by

Mailing Address: Leonardo Zornoff •

Departamento de Clínica Médica, Rubião Jr, S/N, Postal Code 18618-970, Botucatu, SP – Brazil

E-mail: lzornoff@fmb.unesp.br, lzornoff@cardiol.br

Manuscript received April 18, 2012, revised manuscript October 08, 2012; accepted October 18, 2012.

DOI: 10.5935/abc.20130065

the finding that ischemic preconditioning cannot induce protection in heterozygous Cx43-deficient mice¹¹. However, the effects of smoking on cardiac Cx-43 remain unknown¹². Therefore, to address this issue, we tested the hypothesis that exposure to tobacco smoke would interfere with the intensity, distribution or phosphorylation of Cx43 in the rat heart.

Materials and Methods

Groups and Treatment

All experiments and procedures were performed in accordance with NIH guidelines for the Care and Use of Laboratory Animals and were approved by the Animal Ethics Committee of our Institution (protocol number 399).

Male Wistar rats weighing 100 g, were randomly allocated into 2 groups: 1) Control group (n = 25), control rats without exposure to tobacco smoke; 2) Group exposed to tobacco smoke (ETS), n = 23. The tobacco smoke exposure lasted for 5 weeks.

The ETS rats were exposed to cigarette smoke in a chamber (dimensions 95 x 80 x 65 cm) connected to a smoking device based on a model adapted by Paiva et al^{7,13-17}. The smoke was drawn out of filtered commercial cigarettes (composition per unit: 1.1 mg of nicotine; 14 mg of tar; and 15 mg of carbon monoxide) with a vacuum pump and was exhausted into the smoking chamber. During wk 1, the number of cigarettes was gradually increased from 5 to 10 over a 30-min period, twice in the afternoon. For the next approximately 3–4 wk, until the rats reached the target weight, 10 cigarettes were used in each smoking trial, repeated 4 times/d, twice in the morning and twice in the afternoon. In a previous study, with similar tobacco exposure time, the carboxyhemoglobin levels in smoke-exposed rats were greater than in the control group (control rats: $0.9 \pm 0.7\%$ and ETS rats: $5.3 \pm 2.8\%$, $p = 0.008$)¹⁴. Thus, the latter study confirmed the efficacy of the exposure of smoking animals to cigarette smoke. All animals were fed the same amount of diet, which was based on the food ingestion of the group ETS. After this period of tobacco smoke exposure, rats were euthanized by an excessive dosage of anesthesia (sodium pentobarbital 100 mg/kg of body weight). Rats in the ETS group were not exposed to tobacco smoke during the 12 h before euthanasia.

Morphometric Analysis

At the completion of the treatment period, the right and left ventricles (including the interventricular septum) were dissected, separated, and weighed. Coronal sections, approximately 3 mm in thickness, taken 5 mm from the tip of the ventricles were fixed in a 10% buffered formaldehyde solution for 48 hours followed by immersion in a 70% alcohol solution for 48 hours. After fixation, the tissues were embedded in paraffin. Histological sections four and six microns in thickness were stained, respectively, with Hematoxylin-Eosin (HE) and collagen-specific Picro Sirius red. The myocyte cross-sectional areas (MA) were measured in the HE-stained sections by a LEICA DM LS microscope connected to a video camera and IBM compatible computer, equipped with the image analysis program Image Pro-plus (Media

Cybernetics, Silver Spring, Maryland, USA). Fifty to seventy cells were analyzed per slide. The selected cells presented a circular shape and were located in the subendocardial layer of the ventricular walls. This precaution was taken in order to optimize myocyte shape uniformity among the groups. The average cross-sectional area obtained for each group was used as an indicator of cell hypertrophy. The slides stained with Picro Sirius red were used to quantify the interstitial collagen volume fraction using video densitometry. Images of the cardiac tissue were taken and analyzed using the system described above. The elements of the cardiac tissue were identified according to color level. Therefore, the collagen fibers were visualized in red and the myocytes in yellow. The collagen volume fraction (CVF) was calculated automatically and corresponded to the sum of the collagen areas divided by the sum of the collagen tissue and myocyte areas. An average of thirty fields was analyzed using a lens with 40X magnification. Perivascular collagen was excluded from the analysis^{14,16}.

Immunohistochemistry for connexin 43

Immunohistochemistry experiments were performed according to Saffitz et al¹⁸ with some modifications¹⁹. All hearts, transversally cut, were fixed, successively, in 10% buffered formaldehyde solution for 24 hours, running water for 24 hours and 70 percent alcohol for 24 hours. After fixation, tissues were embedded in paraffin and sliced into 3- μ m sections. Before immunolabeling, tissue characterization and orientation were recorded by hematoxylin-eosin staining.

The sections (3 μ m) were placed on poly-D-lysine-coated glass slides and successively dewaxed three times, for 5 minutes each, in baths of xylol, 100% alcohol, 95% alcohol and 70% alcohol followed by rehydration in distilled water for 10 minutes. Antigenic sites were retrieved by treatment of slices in 0.01 M acid citric buffer in phosphate-saline buffer (PBS) pH 6.0 for 10 minutes in microwave environment. After this period, the slices were kept ice-cold for 20 minutes followed by washing in PBS pH 7.2. The sections were blocked with 3% serum albumin bovine (BSA) solution in PBS for 30 minutes at room temperature. After this procedure, the sections were incubated overnight with rabbit polyclonal anti-connexin 43 antibody (Abcam Inc. Cambridge) diluted 1:100 in 2% BSA in PBS at 4 °C. On the next morning, the sections were washed in PBS for 15 minutes three times and incubated in secondary antibody (Texas-red anti-rabbit IgG, Vector) at 1:50 dilution in 2% BSA in PBS at room temperature for 1 hour.

The sections were washed in PBS for 15 minutes three times and mounted in glycerol medium. Ten sections were processed simultaneously in each immunolabeling run while negative controls, in which the primary antibody was omitted, were routinely run in parallel²⁰ with the objective of monitoring any occurrence of unspecific stain. Immunolabeling of Cx43 was examined by epi-fluorescent microscopy (Carl-Zeiss, Inc. North America) equipped with a WG filter cube (excitation = 550 nm and emission = 650 nm) connected to a video camera and compatible computer (AxioPlan 4.1; Carl-Zeiss Inc.). All images were recorded on the same day as labeling and the tissues were coded and examined blindly by the same person.

From each heart, as described by Kostin et al²¹, 10 optical fields were recorded and analyzed for distribution of Cx43 in the myocytes (i.e. presence of Cx43 within or outside the intercalated disk) and intensity of Cx43 staining at intercalated disks.

Western blotting for connexin 43

The western blotting experiments were performed according to Rolim et al²². The samples were homogenized in ice-cold lyses buffer (50 mM potassium phosphate buffer pH 7.0; 0.3 mM sucrose; 0.5 mM DTT; 1 mM EDTA pH 8.0; 0.3 mM PMSF; 10 mM NaF and protease inhibitor cocktail).

The supernatant was collected by centrifugation at 12000 rpm for 20 minutes and stored at -80 °C. The determination of proteins was performed by BSA protein standard (Bio-Rad, Hercules, CA, USA) curves and the Bradford method. The protein samples were diluted in Laemmli buffer (240 mM Tris-HCl; 0.8% SDS; 40% glycerol; 0.02% bromophenol blue and 200 mM β-mercapthoethanol). Then 50 μg of each sample was separated on 10% polyacrylamide stacking gels (240 mM Tris-HCl pH 6.8; 30% polyacrylamide; APS and TEMED) and resolving gels (240 mM Tris-HCl pH 8.8; 30% polyacrylamide; APS and TEMED). Electrophoresis was run at 120 V for 90 minutes.

The transfer onto nitrocellulose membrane (Bio-Rad) was performed in the Mini-Trans Blot system (Bio-Rad, Hercules, CA, USA) with transfer buffer (25 mM Tris; 192 mM glycine; 20% methanol and 0.1% SDS) for 90 minutes at 120 mA. The blots were blocked with 5% non-fat milk for 2 hours at room temperature with shaking and washing three times in PBS Tween (1mM Tris pH 2.8; 5 mM NaCl and Tween 20). Afterwards, the blots were immunolabeled overnight with a monoclonal mouse anti-Cx43 primary antibody (Ab11369 - Abcam), corresponding to amino acids 130–143 of Cx43 diluted to 1: 250 in PBS Tween (1mM Tris pH 2.8; 5 mM NaCl and Tween 20) or monoclonal mouse anti-Cx43 (13-8300 – Zymed) diluted to 3μg/mL, specific to serine 368, an unphosphorylated residue of Cx43.

On the next morning, the membranes were washed three times in PBS Tween (1mM Tris pH 2.8; 5 mM NaCl and Tween 20). This was followed by incubation for 2 hours at room temperature with an HRP- conjugated anti-mouse antibody (Sigma) diluted to

1:10000. After rinsing three times in PBS Tween (1mM Tris pH 2.8; 5 mM NaCl and Tween 20), the membranes were incubated for 1 minute in ECL solution (Amersham) (*Solution I*: 2.5mM luminol, 400 μM p-cumaric acid, 100 μM tris pH 8.5; *solution II*: 5.4 mM 30% hydrogen peroxide, 100 μM tris pH 8.5). After this period, the membranes were exposed to x-ray film (Eastman Kodak Co, USA) for 10 minutes in a darkroom and then successively dipped in revealing solutions, water and fixative for 1 minute each. The reaction between ECL reagent and the HRP- conjugated anti-mouse antibody on the membranes emits light that appears as dark bands on the x-ray film after revelation.

After this procedure, densities of the dark bands obtained were analyzed by densitometric quantification of signal intensity by Scion Image (Scion Corporation, Frederick, Maryland, USA) software with the background measurements of signal intensity being subtracted individually from each Lane¹⁹.

Statistical analysis

Data were expressed as means ± SD. The statistical analysis was performed by Student's *t* test. Differences were considered significant at level of 5%. Statistical analyses were performed using SigmaStat for Windows v2.03 (SPSS Inc, Chicago, IL).

Results

The morphometric variables did not differ statistically between control and ETS groups (Table 1).

Smoke exposure induced lower stain intensity of Cx43 at ID when compared to control rats (Control: 2.32 ± 0.19; ETS: 1.73 ± 0.18; *p* = 0.04). The distribution of Cx43 at ID did not differ between control and ETS rats (Control: 3.73 ± 0.12; ETS: 3.20 ± 0.17; *p* = 0.18); (Figure 1).

ETS rats showed higher values of dephosphorylated form of Cx43 than control rats (Control: 0.45 ± 0.11; ETS: 0.90 ± 0.11; *p* = 0.03). On the other hand, total Cx43 did not differ between control and ETS groups (Control: 0.75 ± 0.19; ETS: 0.93 ± 0.27; *p* = 0.58); (Figure 2).

Table 1 - Morphometric data

	Control (n = 25)	ETS (n = 23)	p
BW (g)	237 ± 20.5	230 ± 19.6	0.30
LV (g)	0.52 ± 0.05	0.52 ± 0.05	0.97
RV (g)	0.15 ± 0.015	0.15 ± 0.014	0.45
LV/BW (g)	2.21 ± 0.15	2.27 ± 0.14	0.15
RV/BW (g)	0.64 ± 0.05	0.67 ± 0.10	0.11
Atrium (g)	0.05 ± 0.015	0.045 ± 0.014	0.16
Myocyte area (μ ²)	190 ± 51.6	184 ± 30.8	0.68
Collagen (%)	4.34 ± 2.75	4.41 ± 0.96	0.90

Values are mean ± SD. n: number of rats; BW: body weight; LV: left ventricle; RV: right ventricle; LV/BW: left ventricle normalized by BW; RV/BW: right ventricle normalized by BW. Student's *t* test.




Figure 1 - Cx43 immunofluorescence micrographs from sections of rat myocardium: control (A), exposed to tobacco smoke (B and C) and negative control (D). Arrows in A: Intercalated disks with immunoreactive signal for Cx43; in B and C: immunoreactive signal for Cx43 at intercalated disks and plasma membrane; in D: intercalated disks free of stain for Cx43. Bar = 20 μ m.

Discussion

This study aimed to analyze the effects of exposure to tobacco smoke on the intensity, distribution, and phosphorylation of Cx43 in the rat heart. Our data indicated that tobacco smoke exposure induced remodeling of the cardiac gap junctions.

An important issue to be considered is that, in addition to the already well-known effects on the vascular endothelium

and lungs, previous studies showed that cigarette smoke exposure that lasts at least two months induces cardiac remodeling, characterized by morphological and functional alterations. Indeed, smoking produced augmentations in the left ventricular diastolic and systolic diameters, and induced left ventricular hypertrophy. These morphological alterations were associated with reduction of left ventricular systolic function^{7,13,14,16,19,23-27}. However, the effects of tobacco smoke exposure on the cardiac gap junctions are still unknown.




Figure 2 - Western blot analysis. A: Blot with anti-total-Cx43, B: non-phosphorylated form of Cx43 detected by monoclonal mouse anti-np-Cx43. C: GAPDH bands.

The main finding of the present study was that exposure to tobacco smoke induced remodeling of the cardiac gap junctions. Indeed, the remodeling process was characterized by a lower quantity of Cx43 at ID and an increase in the non phosphorylated form of Cx43 in the myocardium.

Another important issue is that cardiac remodeling characteristically starts with genetic, molecular, cellular, biochemical, and then structural alterations²⁸⁻³¹. In this context, our findings of Cx43 alterations and no differences in the left and right ventricular morphological variables can be interpreted as the observation of an early stage of the cardiac remodeling. Therefore, we assume that Cx43 remodeling precedes the morphological and functional alterations induced by tobacco smoke exposure.

Gap junctions are clusters of transmembrane channels that connect the cytoplasmic compartments of neighboring cells. The component proteins of gap junction channels are termed connexins. There are many types of connexins, but the predominant isoform in the heart, mainly in the ventricle, is Cx43¹⁰. A major role of gap junctions in the myocardium is to coordinate electrical excitation and to facilitate intercellular change of small proteins, such as signal transduction proteins. Importantly, the normal gap junction functions are related to normal expression and distribution of Cx43, given that Cx43 is located mainly in the intercalated disc of cardiomyocytes. In addition, the degree of Cx43 phosphorylation is critical to gap junction function¹⁰.

Alterations in the quantity, localization and phosphorylation of Cx43 have been observed when the heart was submitted to ischemic or hypoxic events^{21,32,33} and when these events are preceded by ischemic preconditioning^{34,35}. Such ischemia induces reduction of cardiac gap junction, characterized by lower expression of Cx43^{32,36}. Thus, the observed alterations in Cx43 intensity provoked by smoking in the present study are similar to those induced by ischemia.

The diminished Cx43 staining intensity at ID induced by exposure to tobacco smoke in the present study resembles the effects of ischemic preconditioning, which initiates a process of Cx43 occurrence in the free plasma membrane, outside of the intercalated disks. As a consequence of this effect, ischemic preconditioning may improve the cell's chance of survival following prolonged ischemia³⁵.

With respect to the increase in the non-phosphorylated form of Cx43 induced by exposure to tobacco smoke, many studies have evidenced that the dephosphorylation of Cx43 causes electrical uncoupling in the cardiac myocytes³⁴ and the inhibition of Cx43 dephosphorylation lead to protective effects on cardiac function during hypoxic injury³⁷. In this context, studies found that gap junction-uncoupling agents administered prior to ischemia preserve the electrical coupling of cardiac myocytes during subsequent ischemic insult, similarly to ischemic preconditioning³⁸. These authors suggest that the partial and reversible uncoupling of gap junctions prior to ischemia results in an antiarrhythmic effect during a prolonged ischemic insult, similarly to ischemic preconditioning.

In this context, the present study found that exposure to tobacco smoke produced a remodeling process in the cardiac gap junction, an effect similar to ischemic preconditioning. Therefore, our data strongly suggest that exposure to tobacco smoke may induce an ischemic preconditioning-like effect in the rat heart. Importantly, this phenomenon could explain the smoker's paradox observed in some studies¹⁻⁴.

The mechanisms involved in the effects of tobacco smoke exposure on cardiac gap junctions are unknown. Among the possible mechanisms, ventricular remodeling induced by such exposure is associated with augmented activation of mitogen-activated protein kinases²³. In addition, there is evidence that the PKC p38 and p42/44 MAPK pathways regulate Cx43 expression³⁹.

Our study should be analyzed considering a potential limitation. We did not include a preconditioning group. However, the alterations in the Cx43 induced by the preconditioning are well established. Therefore we strongly believe that our data add important information about this subject.

Conclusion

This study has shown that exposure to tobacco smoke resulted in cardiac gap junction remodeling, characterized by alterations in the quantity and phosphorylation of Cx43, in rats hearts. This finding could explain the smoker's paradox observed in some studies.

Author contributions

Conception and design of the research: Novo R, Paiva SAR; Acquisition of data: Novo R, Freire CM, Felisbino S, Minicucci MF, Azevedo PS; Analysis and interpretation of the data: Novo R, Freire CM, Felisbino S, Minicucci MF, Azevedo PS, Zornoff LAM, Paiva SAR; Statistical analysis: Novo R, Minicucci MF, Paiva SAR; Writing of the manuscript: Novo R, Zornoff LAM, Paiva SAR; Critical revision of the manuscript for intellectual content: Freire CM, Felisbino S, Minicucci MF, Azevedo PS, Zornoff LAM, Paiva SAR.

Potential Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Sources of Funding

There were no external funding sources for this study.

Study Association

This article is part of the thesis of doctoral submitted by Rosângela Novo, from Faculdade de Medicina de Botucatu.

References

- Barbash GI, Reiner J, White HD, Wilcox RG, Armstrong PW, Sadowski Z, et al. Evaluation of paradoxical beneficial effects of smoking in patients receiving thrombolytic therapy for acute myocardial infarction: mechanism of the "smoker's paradox" from the GUSTO-I trial, with angiographic insights. Global Utilization of Streptokinase and Tissue-Plasminogen Activator for Occluded Coronary Arteries. *J Am Coll Cardiol.* 1995;26(5):1222-9.
- Kievit PC, Brouwer MA, Veen G, Aengevaeren WR, Verheugt FW. The smoker's paradox after successful fibrinolysis: reduced risk of reocclusion but no improved long-term cardiac outcome. *J Thromb Thrombolysis.* 2009;27(4):385-93.
- Barua RS, Sy F, Srikanth S, Huang G, Javed U, Buhari C, et al. Acute cigarette smoke exposure reduces clot lysis--association between altered fibrin architecture and the response to t-PA. *Thromb Res.* 2010;126(5):426-30.
- Aune E, Roislien J, Mathisen M, Thelle DS, Otterstad JE. The "smoker's paradox" in patients with acute coronary syndrome: a systematic review. *BMC Med.* 2011;9:97.
- Evrengul H, Selesi D, Tanriverdi H, Kaftan A. The antiarrhythmic effect and clinical consequences of ischemic preconditioning. *Coron Artery Dis.* 2006;17(3):283-8.
- Yellon DM, Downey JM. Preconditioning the myocardium: from cellular physiology to clinical cardiology. *Physiol Rev.* 2003;83(4):1113-51.
- Paiva SA, Novo R, Matsubara BB, Matsubara LS, Azevedo PS, Minicucci MF, et al. beta-carotene attenuates the paradoxical effect of tobacco smoke on the mortality of rats after experimental myocardial infarction. *J Nutr.* 2005;135(9):2109-13.
- Opitz CF, Mitchell GF, Pfeffer MA, Pfeffer JM. Arrhythmias and death after coronary artery occlusion in the rat: continuous telemetric ECG monitoring in conscious, untethered rats. *Circulation.* 1995;92(2):253-61.
- Schulz R, Heusch G. Connexin 43 and ischemic preconditioning. *Cardiovasc Res.* 2004;62(2):335-44.
- Severs NJ, Bruce AF, Dupont E, Rothery S. Remodelling of gap junctions and connexin expression in diseased myocardium. *Cardiovasc Res.* 2008;80(1):9-19.
- Schwane U, Konietzka I, Duschin A, Li X, Schulz R, Heusch G. No ischemic preconditioning in heterozygous connexin43-deficient mice. *Am J Physiol Heart Circ Physiol.* 2002;283(4):H1740-2.
- Azevedo PS, Minicucci MF, Matsubara BB, Matsubara LS, Duarte DR, Paiva SA, et al. [Remodeling pattern and ventricular function in rats exposed to cigarette smoke]. *Arq Bras Cardiol.* 2010;94(2):224-8.
- Castardeli E, Duarte DR, Minicucci MF, Azevedo PS, Matsubara BB, Matsubara LS, et al. Tobacco smoke-induced left ventricular remodelling is not associated with metalloproteinase-2 or -9 activation. *Eur J Heart Fail.* 2007;9(11):1081-5.
- Castardeli E, Paiva SA, Matsubara BB, Matsubara LS, Minicucci MF, Azevedo PS, et al. Chronic cigarette smoke exposure results in cardiac remodeling and impaired ventricular function in rats. *Arq Bras Cardiol.* 2005;84(4):320-4.
- Oliveira LC, Azevedo PS, Minicucci ME, Rafacho BP, Duarte DR, Matsubara LS, et al. Retinoic acid prevents ventricular remodelling induced by tobacco smoke exposure in rats. *Acta Cardiol.* 2011;66(1):3-7.
- Paiva SA, Zornoff LA, Okoshi MP, Okoshi K, Cicogna AC, Campana AO. Behavior of cardiac variables in animals exposed to cigarette smoke. *Arq Bras Cardiol.* 2003;81(3):221-8.
- Rafacho BP, Santos P, Assalin HB, Ardisson LP, Roscani MC, Polegato BF, et al. Role of vitamin D in the cardiac remodeling induced by tobacco smoke exposure. *Int J Cardiol.* 2012;155(3):472-3.
- Saffitz JE, Green KG, Kraft WJ, Schechtman KB, Yamada KA. Effects of diminished expression of connexin43 on gap junction number and size in ventricular myocardium. *Am J Physiol Heart Circ Physiol.* 2000;278(5):H1662-70.
- Duarte DR, Minicucci MF, Azevedo PS, Chiuseo-Minicucci F, Matsubara BB, Matsubara LS, et al. Influence of lisinopril on cardiac remodeling induced by tobacco smoke exposure. *Med Sci Monit.* 2010;16(8):BR255-9.
- Uzzaman M, Honjo H, Takagishi Y, Emdad L, Magee AI, Severs NJ, et al. Remodeling of gap junctional coupling in hypertrophied right ventricles of rats with monocrotaline-induced pulmonary hypertension. *Circ Res.* 2000;86(8):871-8.
- Kostin S, Rieger M, Dammer S, Hein S, Richter M, Klovekorn WP, et al. Gap junction remodeling and altered connexin43 expression in the failing human heart. *Mol Cell Biochem.* 2003;242(1-2):135-44.
- Rolim NP, Medeiros A, Rosa KT, Mattos KC, Irigoyen MC, Krieger JE, et al. Exercise training improves the net balance of cardiac Ca²⁺ handling protein expression in heart failure. *Physiol Genomics.* 2007;29(3):246-52.
- Gu L, Pandey V, Geenen DL, Chowdhury SA, Piano MR. Cigarette smoke-induced left ventricular remodelling is associated with activation of mitogen-activated protein kinases. *Eur J Heart Fail.* 2008;10(11):1057-64.
- Zornoff LA, Matsubara BB, Matsubara LS, Minicucci MF, Azevedo PS, Camapanha AO, et al. [Cigarette smoke exposure intensifies ventricular remodeling process following myocardial infarction]. *Arq Bras Cardiol.* 2006;86(4):276-82.
- Zornoff LA, Matsubara LS, Matsubara BB, Okoshi MP, Okoshi K, Dal Pai-Silva M, et al. Beta-carotene supplementation attenuates cardiac remodeling induced by one-month tobacco-smoke exposure in rats. *Toxicol Sci.* 2006;90(1):259-66.
- Rafacho BP, Azevedo PS, Polegato BF, Fernandes AA, Bertoline MA, Fernandes DC, et al. Tobacco smoke induces ventricular remodeling associated with an increase in NADPH oxidase activity. *Cell Physiol Biochem.* 2011;27(3-4):305-12.

27. Talukder MA, Johnson WM, Varadharaj S, Lian J, Kearns PN, El-Mahdy MA, et al. Chronic cigarette smoking causes hypertension, increased oxidative stress, impaired NO bioavailability, endothelial dysfunction, and cardiac remodeling in mice. *Am J Physiol Heart Circ Physiol.* 2011;300(1):H388-96.
28. Cohn JN, Ferrari R, Sharpe N. Cardiac remodeling--concepts and clinical implications: a consensus paper from an international forum on cardiac remodeling. *Behalf of an International Forum on Cardiac Remodeling. J Am Coll Cardiol.* 2000;35(3):569-82.
29. Pfeffer MA, Braunwald E. Ventricular remodeling after myocardial infarction: experimental observations and clinical implications. *Circulation.* 1990;81(4):1161-72.
30. Swynghedauw B. Molecular mechanisms of myocardial remodeling. *Physiol Rev.* 1999;79(1):215-62.
31. Zornoff LA, Paiva SA, Duarte DR, Spadaro J. Ventricular remodeling after myocardial infarction: concepts and clinical implications. *Arq Bras Cardiol.* 2009;92(2):157-64.
32. Beardslee MA, Lerner DL, Tadros PN, Laing JG, Beyer EC, Yamada KA, et al. Dephosphorylation and intracellular redistribution of ventricular connexin43 during electrical uncoupling induced by ischemia. *Circ Res.* 2000;87(8):656-62.
33. Matsushita S, Kurihara H, Watanabe M, Okada T, Sakai T, Amano A. Alterations of phosphorylation state of connexin 43 during hypoxia and reoxygenation are associated with cardiac function. *J Histochem Cytochem.* 2006;54(3):343-53.
34. Jain SK, Schuessler RB, Saffitz JE. Mechanisms of delayed electrical uncoupling induced by ischemic preconditioning. *Circ Res.* 2003;92(10):1138-44.
35. Vetterlein F, Muhlfield C, Cetegen C, Volkmann R, Schrader C, Hellige G. Redistribution of connexin43 in regional acute ischemic myocardium: influence of ischemic preconditioning. *Am J Physiol Heart Circ Physiol.* 2006;291(2):H813-9.
36. Hatanaka K, Kawata H, Toyofuku T, Yoshida K. Down-regulation of connexin43 in early myocardial ischemia and protective effect by ischemic preconditioning in rat hearts in vivo. *Jpn Heart J.* 2004;45(6):1007-19.
37. Matsushita S, Kurihara H, Watanabe M, Okada T, Sakai T, Amano A. Inhibition of connexin43 dephosphorylation is involved in protective effects of diltiazem on cardiac function during hypoxic injury. *Histol Histopathol.* 2011;26(3):315-22.
38. Papp R, Goncz M, Kovacs M, Seprenyi G, Vegh A. Gap junctional uncoupling plays a trigger role in the antiarrhythmic effect of ischaemic preconditioning. *Cardiovasc Res.* 2007;74(3):396-405.
39. Salameh A, Krautblatter S, Baessler S, Karl S, Rojas Gomez D, Dhein S, et al. Signal transduction and transcriptional control of cardiac connexin43 up-regulation after alpha 1-adrenoceptor stimulation. *J Pharmacol Exp Ther.* 2008;326(1):315-22.