

Schwannoma of the lower lip mimicking a mucocele in children

Bruna Noemia Ferreira de Menezes^a , John Lennon Silva Cunha^b ,
Samuel de Carvalho Chaves-Júnior^c , Bruno Torres Bezerra^a 

How to cite: Menezes BNF, Cunha JLS, Chaves-Júnior SC, Bezerra BT. Schwannoma of the lower lip mimicking a mucocele in children. *Autops Case Rep* [Internet]. 2020;10(1):e2019134. <https://doi.org/10.4322/acr.2019.134>

Dear Editor,

Schwannoma or neurilemmoma is a benign tumor of neural origin, uncertain etiology, firstly described in 1910 by Verocay (1910).¹ Schwannomas tend to occur in adults between the third and fifth decades of life and do not show sex predilection.² Most of the lesions occur in the head and neck regions (25-40%).³ However, despite the lips and the oral cavity are highly innervated anatomic sites, they are rarely found in these locations. Only 1% of all schwannomas arise in the oral cavity, being the tongue the most common intraoral site of occurrence.⁴ Moreover, schwannomas in the lower lip are excessively rare during childhood and adolescence, with only a few well-documented cases published in the English-language literature. Herein, we presented a rare case of schwannoma in the lower lip mimicking a mucocele in a 7-year-old male child.

The patient, a 7-year-old Caucasian male, was referred to the oral medicine service of the School of Dentistry at Tiradentes University, complaining of a painless lesion on the lip over the last 2 months. The extraoral examination was unremarkable. The intraoral examination revealed a fibrous resilient nodular lesion on the left lower lip mucosa, with the coloration resembling normal mucosa, measuring 0.8 cm (Figure 1).

Past medical history was not contributory, except for a local trauma. Based on the clinical

aspects, the diagnostic hypothesis was a mucocele. An excisional biopsy was performed, and the specimen was sent for histopathological analysis. Histological examination showed an encapsulated tumor composed of spindle-shaped cells with well-aligned nuclei interpreted as Schwann cells.

A palisaded arrangement surrounding central acellular areas, known as Verocay bodies, was observed in most parts, forming a typical Antoni A histological pattern. Less cellular areas comprised of diffusely arranged spindle shaped-cells were also observed, characterizing an Antoni B pattern (Figure 2A-C). Immunohistochemical analysis showed a marked positivity of neoplastic cells for S-100 protein (Figure 2D). The diagnosis was schwannoma was made. After one year of follow-up, there was no recurrence of the tumor.

Swellings of the lower lip can represent many entities, and lip schwannoma is overlooked in the initial differential diagnosis due to its rarity in this location.^{3,4} In our case, the submucosal presentation, slow growth pattern, with no hardening of surround tissues, the clearly defined limits of the tumor, and history of the habit of biting the mucosa reported during the anamnesis suggested a benign process. Therefore, our diagnostic hypothesis included mainly conditions commonly observed in young patients, such as reactive proliferative processes and mesenchymal tumors,

^aTiradentes University (UNIT), Department of Dentistry. Aracaju, SE, Brazil.

^bUniversity of Campinas (UNICAMP), Piracicaba Dental School, Department of Oral Diagnosis, Oral Pathology Section. Piracicaba, SP, Brazil.

^cUniversity of Campinas (UNICAMP), Piracicaba Dental School, Department of Pediatric Dentistry. Piracicaba, SP, Brazil.





Figure 1. Clinical aspect of the lip lesion. Observe the small nodular sessile lesion, with a similar color to the (arrowhead) mucosa in lower lip.

and the patient was misdiagnosed with mucocele. Mucocele is a common lesion of the lip, asymptomatic in most cases, similar to the mucosa in color, clinically resembling a variety of other lesions that affect oral soft tissues.⁵ Because it is the most common lesion of the lower lip, it would be a dentist's leading hypothesis as a provisional diagnosis. Thus, histopathological analysis is a necessary procedure to assure the correct diagnosis and therapeutic management.

In the oral cavity, the tongue is the most common site affected by schwannoma, and lip tumors are considered rare, with only 21 cases reported in the literature, including the current case. Lip schwannoma tends to occur in adults between the third and fifth decades of life, with a mean age of 26.4 years (range: 7-82) and do not show sex predilection.³ The diagnosis




Figure 2. Photomicrographs of the lip lesion biopsy. **A** – Well-circumscribed spindle-cell lesion located in the superficial planes (H&E, 2,5X); **B** – Proliferation of spindle shaped and ovoid cells organized in small fascicles, sometimes forming a palisading arrangement (H&E, 10X); **C** – detail of neoplastic cells, highlighting well-stained wavy nuclei, with imperceptible nucleoli and Verocay bodies typical of Antoni A pattern (H&E, 20X); **D** – Immunohistochemical analysis revealing intense and diffuse positivity for S-100 protein (20X).

of schwannoma is based on histological features. Morphologically, schwannoma spindle-shaped cells are arranged in two distinct patterns, Antoni A and B, in variable proportions, with a few or none intralésional axons, forming completely encapsulated lesions.^{5,6} The current case fulfills all the morphological criteria to be classified as schwannoma. It has also been highlighted that immunohistochemical positivity for S-100 protein, a marker of neural differentiation, is widely observed in schwannomas.⁴⁻⁶ Marked immunoexpression of S-100 protein was seen in the current case, confirming the neural derivation of the tumor.

Regarding the treatment, a conservative surgical excision is the best approach, and the prognosis is excellent.⁴ Local recurrence of schwannomas has not been reported after complete surgical excision of the tumors, and malignant transformation is exceptionally rare.^{4,6}

Although tumors of the peripheral nerve sheath are rare, they must also be included among the differential diagnoses of nodular lesions of the lower lip;⁵ familiarity with morphological and immunohistochemical aspects of these lesions is essential to avoid diagnostic mistakes and provide a satisfactory therapeutic approach.

The patient signed informed consent declaration authorizing the use of the images and medical information. The manuscript was approved by the Institutional Ethics Committee.

Authors contributions: Menezes BNF wrote the manuscript. Chaves-Júnior SC and Bezerra BT were the surgeon of the patient and provided the images. Cunha JLS performed the microscopic and the critical review of the manuscript, and provided the photomicrographs. All authors approved and proofread the final document.

Conflict of interest: None

Financial support: None

Submitted on: October 2nd, 2019

Accepted on: October 21st, 2019

Correspondence

John Lennon Silva Cunha

Oral Pathology Section - Department of Oral Diagnosis - Piracicaba Dental School - University of Campinas (UNICAMP)

Av. Limeira 901, P.O. Box. 52 – Piracicaba/SP, Brazil

CEP: 1314-903

Phone: +55 (79) 99654-0805

lennonrrr@live.com

REFERENCES

1. Verocay J. Zur Kenntnis der Neurofibrome. *Beitr Pathol Anat.* 1910;48:1-68
2. Alotaiby FM, Fitzpatrick S, Upadhyaya J, Islam MN, Cohen D, Bhattacharyya I. Demographic, clinical and histopathological features of oral neural neoplasms: A retrospective study. *Head Neck Pathol.* 2019;13(2):208-14. <http://dx.doi.org/10.1007/s12105-018-0943-1>. PMID:29931661.
3. Sitenga J, Aird G, Vaudreuil A, Huerter CJ. Clinical features and management of schwannoma affecting the upper and lower lips. *Int J Dermatol.* 2018;57(9):1047-52. <http://dx.doi.org/10.1111/ijd.13920>. PMID:29377087.
4. Desai J. An unexpected and rare outcome of a common nodular mass on upper lip in a pediatric patient with a history of trauma - Schwannoma. *Natl J Maxillofac Surg.* 2019;10(1):102-4. http://dx.doi.org/10.4103/njms.NJMS_48_18. PMID:31205398.
5. Cunha JLS, Silva AF, Rocha RS, Silveira EJD, Albuquerque RLC Jr. Palisaded encapsulated neuroma clinically mimicking a mucocele: report of a case. *J Bras Patol Med Lab.* 2018;54(3):189-92. <http://dx.doi.org/10.5935/1676-2444.20180033>.
6. Vera-Sirera B, Fernades-Ciacha L, Floria LM, Vera-Sempere F. Palatal ancient schwannoma: optical, immunohistochemical and ultrastructural study with literature review. *Eur Arch Otorhinolaryngol.* 2017;274(12):4195-202. <http://dx.doi.org/10.1007/s00405-017-4759-5>. PMID:28975399.