

Artigo: National hospice for the insane and the Brazilian Neurology in the beginning of the 20th century  
 Página 824 Fig.1, onde se lê:





Fig 1. National hospice for the insane inpatients. Left: causes for hospitalization (mainly alcoholism for men, and hysteria, for women. "Dementia precoce" and epilepsy are also relevant causes); Right: death causes (mainly tuberculosis, "general paralysis" and marasmus) (report of 1904–1905, by Afrânio Peixoto<sup>5</sup>).

Leia-se:



MEN WOMEN 1=idocy; 2=imbecility; 3=mental retardation; 4=neurasthenia; 5=hysteria; 6=epilepsy; 7=psychopathic states, degeneration, disnoias; 8=paranoia; 9=toxic psychosis, alcoholism; 10=psychose autotoxicus, exhaustion, etc.; 11= infectious psychosis, syphilis, atypical smallpox, etc; 12=manic depressive madness; 13= precocious dementia; 14=paralytic dementia, general paralysis; 15=terminal dementia; 16=senile dementia; 17=under observation; 18=not alienated.

Fig 1. Causes of hospitalization and deaths observed in 1904 at the National Hospice and Colony for the insane (in 1806 patients – 1136 men and 670 women, 196 deaths). Graphic: causes for hospitalization (mainly alcoholism for men, and hysteria, for women, but "dementia precoce" and epilepsy are also relevant causes); Table: causes of deaths (mainly tuberculosis, "general paralysis" and marasmus) (adapted from report of 1904–1905 by Afrânio Peixoto<sup>5</sup>).

← Various types of madness