

Emergence of kinky hair in Menkes disease

Desenvolvimento de kinky hair na doença de Menkes

Hannah A. Oppenheim¹  Maria A. Montenegro² 

¹ University of California San Diego, Department of Neuroscience, San Diego California, United States.

² Rady Children's Hospital, San Diego California, United States.

Address for correspondence Maria A. Montenegro
(email: mam99@health.ucsd.edu)

Arq. Neuro-Psiquiatr. 2024;82(5):s00441786761.

A three-month-old male presented with developmental delay, rib fractures, and seizures. He had normal-looking hair.¹ A magnetic resonance imaging (MRI) scan of the brain was normal. Pathogenic *ATP7A* mutation was identified, which confirmed Menkes Disease. Only at 9 months of age did he develop patches of kinky hair.

Parenteral copper histidine supplementation can modify Menkes disease progression if initiated within days after birth.^{2,3} The clinical diagnosis relies on phenotypic presentation, especially sparse and lusterless scalp

hair typically apparent by 2 to 3 months of age; however, our patient demonstrated that this feature may not appear until much later in life. Therefore, hair abnormalities should not be relied upon to initiate genetic testing (► **Figure 1**).

Authors' Contributions

HAO, MAM: conceptualization, data curation, formal analysis, writing of the original draft, and writing – review and editing.




Figure 1 (A) Normal hair at 3 months of age; and (B) hair strand at 9 months of age with classic *pili torti* seen in Menkes disease.

received
January 16, 2024
accepted
February 25, 2024

DOI <https://doi.org/10.1055/s-0044-1786761>.
ISSN 0004-282X.

Editor-in-Chief: Ayrton Roberto Massaro.
Associate Editor: Juliana Gurgel-Gianetti.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Conflict of Interest

The authors have no conflict of interest to declare.

References

- 1 Tümer Z, Møller LB. Menkes disease. *Eur J Hum Genet* 2010;18(05):511–518. Doi: 10.1038/ejhg.2009.187
- 2 Vairo FPE, Chwal BC, Perini S, Ferreira MAP, de Freitas Lopes AC, Saute JAM. A systematic review and evidence-based guideline for diagnosis and treatment of Menkes disease. *Mol Genet Metab* 2019;126(01):6–13. Doi: 10.1016/j.yimgme.2018.12.005
- 3 Kodama H, Murata Y, Kobayashi M. Clinical manifestations and treatment of Menkes disease and its variants. *Pediatr Int* 1999;41(04):423–429. Doi: 10.1046/j.1442-200x.1999.01095.x