



Influence of agrochemicals fipronil and imidacloprid on the learning behavior of *Apis mellifera* L. honeybees

Marcela Pedraza Carrillo, Thaís de Souza Bovi, Adriana Fava Negrão and Ricardo de Oliveira Orsi*

Departamento de Produção Animal, Faculdade de Medicina Veterinária Zootecnia, Universidade Estadual Paulista "Júlio de Mesquita Filho", Distrito de Rubião Júnior, s/n, 18618-970, Botucatu, São Paulo, Brazil. *Author for corresponding. E-mail: orsi@fmvz.unesp.br

ABSTRACT. Agrochemicals on crop cultivated areas is a source of contamination for bees and may cause physiological and behavioral disorders and mortality. The LD₅₀ of the pesticides fipronil and imidacloprid was determined and their effect on the learning behavior of *Apis mellifera* L. honeybee evaluated. LD₅₀ was determined by the ingestion of contaminated food with different concentrations of insecticide concentrations: Fipronil (0, 0.8, 0.4, 0.2, 0.1 and 0.05 µg bee⁻¹) and imidacloprid (0, 0.4, 0.2, 0.1, 0.05 and 0.025 µg bee⁻¹). The method of proboscis extension reflection (PER) and learning through citral odor evaluated their responses to food stimulation. LD₅₀ obtained were 0.28 ± 0.11 and 0.10 ± 0.04 µg bee⁻¹ for fipronil and imidacloprid, respectively. The PER test showed no significant difference (p < 0.05) although agrochemicals affected the learning of bees. Insecticides fipronil and imidacloprid are extremely harmful to foraging Africanized *Apis mellifera* bees.

Keywords: beekeeping, toxicity, lethal dose.

Influência dos agroquímicos Fipronil e Imidaclopride no aprendizado de abelhas *Apis mellifera* L.

RESUMO. O uso de agroquímicos nas áreas de cultivo representa uma fonte de contaminação para as abelhas, podendo ocasionar distúrbios fisiológicos, comportamentais e mortalidade. O objetivo do presente trabalho foi determinar a DL₅₀ dos agroquímicos Fipronil e Imidaclopride e avaliar o efeito destes no aprendizado de abelhas *Apis mellifera* L. adultas. A DL₅₀ foi determinada por meio de teste de ingestão de alimento contaminado, com diferentes concentrações dos inseticidas Fipronil (0; 0,8; 0,4; 0,2; 0,1 e 0,05 µg abelha⁻¹) e Imidaclopride (0; 0,4; 0,2; 0,1; 0,05 e 0,025 µg abelha⁻¹). Para avaliar as respostas ao estímulo do alimento foi utilizado o método de reflexão de extensão da probóscide (REP) e aprendizado, utilizando odor citral. As DL₅₀ obtidas foram 0,28 ± 0,11 e 0,10 ± 0,04 µg abelha⁻¹ para o Fipronil e Imidaclopride, respectivamente. O teste REP não apresentou diferenças significativas (p < 0,05); porém, os agroquímicos afetaram o aprendizado das abelhas. Dessa maneira, conclui-se que os inseticidas Imidaclopride e Fipronil são nocivos às abelhas campeiras de *Apis mellifera*.

Palavras-chave: apicultura, toxicidade, dose letal.

Introduction

As any other group of living beings, bees need chemical substances for growth, reproduction and other vital functions. These chemical substances, named nutrients, may be obtained through nutrition. Plants are the main nutrition source for these insects (EDWARDS; WRATTEN, 2000).

However, the expansion of cultivation areas, especially fields planted with a single plant species, promotes the emergence of pests and diseases which make agriculture even more dependent on agrochemicals (COUTINHO et al., 2005). These products enter the soil-water-plant chain and represent a direct and indirect dangerous source of contamination to bees, which, in their turn,

contaminate their products.

Bees poisoning may occur by contact or ingestion during flower visiting. During forage, bees get in touch with different groups of agrochemicals, such as insecticides, fungicides, acaricides and herbicides. However, the effect of insecticides on the pollinator fauna of agroecology systems has been well shown and is directly responsible for the decline of bee populations and indirectly for economic losses owing to the population decline of pollinators (RICHARDS; KEVAN, 2002). In the United States, about one third of commercial crops undergo liabilities through the decrease of natural pollinators (PAOLETTI, 1999).

Agrochemical group of insecticides acts directly upon the nervous system and causes over activity of neurons, or rather, those connected to learning and memory, and, as a consequence, may change brain structures with paralysis of leg, wings and digestive tract. Insects stop consuming water and food and, consequently, die from starvation or dehydration (MALASPINA; STORT, 1985). These products damage the return flight to the nest or the information given by the forager to the other worker bees through dance and sound (PHAM-DELÈGUE et al., 2002).

Commercially known as *Regent*, the insecticide fipronil belongs to the pyrazole class used to control agricultural pests. Nevertheless, the agrochemical affects non-target insects and causes the mortality of important pollinators such as *Apis mellifera* L. (GUNASEKARAN et al., 2007). In sub-lethal doses, fipronil may affect the gustatory perception, olfactory learning and motor activity of bees, that are essential functions to forage insects (HASSANI et al., 2005).

Within the neonicotinoid chemical group, the insecticide *Gaucho*, whose active agent is imidacloprid, is highly important. The above pesticide is extremely toxic to bees, which, even in low concentrations, causes high mortality due to the toxicity of its metabolites (FAUCON et al., 2005), whereas sub-lethal doses change foraging and communication (DECOURTYE et al., 2003).

One method to evaluate agrochemical toxicity is through LD₅₀ tests, i.e., an acute dose may cause 50% mortality in a population. Generally, LD₅₀ tests are done by contact and/or ingestion, and mortality is recorded 24/48h after exposure to the substance (DEVILLERS, 2002).

Other method of toxicity study in bees is through the proboscis extension reflex (PER) test which evaluates the bee's apprenticeship with olfactory stimulus. During bee foraging, a learning process occurs by floral parameters, such as shape, location, color, and flower odor, associated with food reward (MENZEL; MÜLLER, 1996). In this manner, the proboscis extension reflex (PER) test, based on a temporal association paired to conditioned and unconditioned stimuli, simulates honey-bee-plant interactions. During conditioning, PER is caused by contact with taste receptors of the antennae with a sucrose solution (unconditioned stimulus), and simultaneously providing an odor (conditioned stimulus). When the proboscis extension occurs, the bee is immediately rewarded

by receiving a sucrose solution (ROMERO et al., 2008).

Current research determines the lethal dose (DL₅₀) of the insecticide fipronil and imidacloprid and evaluates the effect on learning of the bee *Apis mellifera* L.

Material and methods

Experiment location and honeybees samples

The experiment was conducted at the Beekeeping Production Area, located in Lageado Experimental Farm, Faculty of Veterinary Medicine and Animal Science, UNESP, Botucatu, 22°50'30.16"S; 48°25'41.90"W.

Over 20-day-old Africanized honeybees from five hives, standardized with regard to number of brood frames and food, containing a young queen, were used. During the experiments, the hive entrance was closed for ten minutes and forager bees returning from the field were collected. They were placed in a recipient with a screen plastic and transported to the laboratory for the accomplishment of the experiment.

LD₅₀ determination

LD₅₀ toxicity was determined according to methodology described by Miranda et al. (2003), with modifications. Ten adult bees over 20-days-old were placed in wooden boxes containing screen sides (25.0 x 15.0 x 10.0 cm) and kept at room temperature. Further, 1 mL of honey was supplied in a feeder with test tube lid containing different doses µg bee⁻¹ of the pesticides: 0, 0.8, 0.4, 0.2, 0.1 and 0.05 in the case of fipronil and 0, 0.4, 0.2, 0.1, 0.05 and 0.025 in the case of imidacloprid. The number of bees with behavioral modifications or dead during 30, 60, 90 and 120 minutes was reported. A control group received only honey and the experiments were performed in triplicate. The results obtained were used to calculate LD₅₀, or rather, the concentration that would kill 50% of the bee population.

Pesticides effect on reflection and learning by proboscis extension

Proboscis Extension Reflection (PER) method was employed to evaluate the response to food stimulus (sugar syrup 50%) (SCHEINER et al., 2004). Twenty bees per treatment were used for the experiments, or rather, control (without agrochemical), fipronil and imidacloprid (LD₅₀). The bees were kept individually in plastic tubes, so that only the antennae were free to the outside. The bees remained necessarily without food for a period of three hours prior to tests. The antennae were then stimulated with a solution of sugar syrup and the number of individuals that exhibited PER was

recorded. The control group received only a solution of sugar syrup without any agrochemical.

For the performance tests of learning and memory, bee antenna was put in contact with a floral odor citral which stimulated the antennae's gustatory receptors. After the bees learned and memorized the floral odor, fipronil and imidacloprid were administered at previously obtained LD₅₀ concentrations. Reflex or its absence of the proboscis extension after stimulus with floral odor was observed.

All tests were performed in triplicate.

Statistical analysis

Probit analysis with BioStat program was employed for LD₅₀ calculation and the evaluation of bee learning was undertaken by analysis of variance with Tukey's test at 5% significance level to compare means (ZAR, 1996).

Results and discussion

Table 1 shows rates for LD₅₀ ($\mu\text{g bee}^{-1}$) obtained by ingestion test of insecticides fipronil and imidacloprid.

Table 1. Determination of mean lethal dose (LD₅₀) in $\mu\text{g bee}^{-1}$ for fipronil and imidacloprid in *Apis mellifera* honeybees by ingestion test. Results represent average and standard deviation.

Fipronil	Imidacloprid
0.28 ± 0.11	0.10 ± 0.04

Research demonstrated that LD₅₀ rates for insecticides under analysis lay below 2 $\mu\text{g bee}^{-1}$ and thus classified as highly toxic to bees, according to Johansen and Mayer (1990).

Fipronil's high toxicity is due to the mechanism which interferes with the functioning of the neurotransmitter GABA (amino-butyric acid) and which breaks the normal neuronal influx (chloride ions passage), causing excessive excitation, severe paralysis and death (GUNASEKARAN et al., 2007).

Similarly, imidacloprid acts as a competitive agonist of the insects' nicotinic receptors, triggering severe neurotoxic effects (SUCHAIL et al., 2003). Suchail

et al. (2000) registered LD₅₀ reported rates of 5 and 25 ng bee^{-1} respectively for ingestion and contact tests.

It has also been reported that honeybees exposed to LD₅₀ of the insecticides above showed no changes to the reflection of proboscis extension. However, their learning was affected, or rather, the reaction to the odor stimulant used within experimental

conditions was significantly reduced. Hassani et al. (2005) observed changes in motor activity and a significant decrease in olfactory learning of honeybees exposed to a dose of 1 ng bee^{-1} of topically applied Fipronil. Decourtye et al. (2004) found that Imidacloprid reduced the olfactory perception and flight activity in worker bees exposed to sub-lethal doses. Bees exposed to doses of Fipronil by contact and ingestion (0.1 and 0.01 ng bee^{-1} , respectively), showed an impairment on the olfactory learning. Whereas bees treated with 0.1 ng bee^{-1} died after 7 days of exposure, those treated with 0.01 ng bee^{-1} ingested more water than those in other treatments and had their movements impaired (ALIOUANE et al., 2009).

Table 2 shows data related to the proboscis extension reflex test and learning when the insecticides fipronil and imidacloprid were used.

Table 2. Percentage of negative response to proboscis extension reflex test (PER) and learning in control, fipronil and imidacloprid treatments for *A. mellifera*. Results represent averages and standard deviation.

	Control	Fipronil	Imidacloprid
PER	17.94 ± 0.12 a*	23.25 ± 0.33 a	44.71 ± 0.00 a
Learning	6.25 ± 8.80 a	59.00 ± 2.10 b	70.63 ± 15.00 b

*Different lower case letters on the same line indicate statistical difference between the averages ($p \leq 0.05$)

Exposure of bees to agrochemicals causes adverse impacts on their foraging activity and reduces the worker bees' longevity (MCKENZIE; WINSTON, 1989) leading to physiological changes, individual behavior and cell physiology alterations, due to stress induced by chemicals agents (GREGORC; ELLIS, 2011).

One factor that is linked to pollinators' loss is Colony Collapse Disorder (CCD), characterized by a great decrease in the number of worker bees. CCD was first reported in the United States in 2006-2007, which lost an average of 45% of its hives (COX-FOSTER et al., 2007). Mullin et al. (2010) have reported a direct association between exposure of bees and agrochemicals, CCD occurrence and pollinators decline. The combination of different effects by pesticides causes high liabilities to bees and is intrinsically linked to the decrease in bee population observed over the past years (ALIOUANE et al., 2009).

The effect of insecticides on the agro-ecosystems pollinator wildlife has been well evidenced, being directly responsible for the reduction of bee populations and indirectly for economic losses as a result of the insects' population decline (KEVAN, 1999).

Current results indicate the need for further

studies that deal with the toxicity of agrochemicals on bees, coupled to the importance of the use of non-toxic pesticides to wildlife pollinators.

Conclusion

Current research found insecticides fipronil and imidacloprid are extremely harmful to foraging Africanized *Apis mellifera* bees.

References

- ALIOUANE, Y.; HASSANI, A. K.; GARY, V.; ARMENGAUD, C.; LAMBIN, M.; GAUTHIER, M. Subchronic exposure of honeybees to sub-lethal doses of pesticides: effects on behavior. **Environmental Toxicology and Chemistry**, v. 28, n. 1, p. 113-122, 2009.
- COUTINHO, C. F. B.; TANIMOTO, S. T.; GALLI, A.; GARBELLINI, G. S.; TAKAYAMA, M.; AMARAL, R. B.; MAZO, L. H.; AVACA, L. A.; MACHADO, S. A. S. Pesticidas: mecanismos de ação, degradação e toxidez. **Pesticidas: Revista de Ecotoxicologia Meio Ambiente**, v. 15, p. 65-72, 2005.
- COX-FOSTER, D. L.; CONLAN, S.; HOLMES, E. C.; PALACIOS, G.; EVANS, J. D.; MORAN, N. A.; QUAN, P. L.; BRIESE, T.; HORNIG, M.; GEISER, D. M.; MARTINSON, V.; ENGELSDORP, D.; KALKSTEIN, A. L.; DRYSDALE, A.; HUI, J.; ZHAI, J.; CUI, L.; HUTCHISON, S. K.; SIMONS, J. F.; EGHOLM, M.; PETTIS, J. S.; LIPKIN, W. I. A metagenomic survey of microbes in honey bee colony collapse disorder. **Science**, v. 318, n. 5848, p. 283-287, 2007.
- DECOURTYE, A.; DEVILLERS, J.; CLUZEAU, S.; CHARRETON, M.; PHAM-DELÈGUE, M. H. Effects of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. **Ecotoxicology and Environmental Safety**, v. 57, n. 3, p. 410-419, 2004.
- DECOURTYE, A.; LACASSIE, E.; PHAM-DELÈGUE, M. H. Learning performances of honey bees are differentially affected by imidacloprid according to the season. **Pest Management Science**, v. 59, n. 3, p. 269-278, 2003.
- DEVILLERS, J. Acute toxicity of pesticides to honey bees. In: DEVILLERS, J.; PHAM-DELÈGUE, M. H. (Ed.). **Honey bees: estimating the environmental impact of chemicals**. London: Taylor and Francis, 2002. p. 56-66.
- EDWARDS, P. J.; WRATTEN, S. D. **Ecologia das interações entre insetos e plantas**. São Paulo: EPU, 2000.
- FAUCON, J. P.; AURIÈRES, C.; DRAJNUDEL, P.; MATHIEU, L.; RIBIÈRE, M.; MARTEL, A. C.; ZEGGANE, S.; CHAUZAT, M. P.; AUBERT, M. F. Experimental study on the toxicity of imidacloprid given in syrup to honey bee (*Apis mellifera*) colonies. **Pest Management Science**, v. 61, n. 2, p. 111-125, 2005.
- GREGORC, A.; ELLIS, J. D. Cell death localization in situ in laboratory reared honey bee (*Apis mellifera* L.) larvae treated with pesticides. **Pesticide Biochemistry and Physiology**, v. 99, n. 2, p. 200-207, 2011.
- GUNASEKARAN, A. S.; TRUONG, T.; GOH, S. K.; SPURLOCK, F.; TJEERDEMA, R. S. Environmental fate and toxicology of fipronil. **Journal of Pesticides Science**, v. 32, n. 3 p. 189-199, 2007.
- HASSANI, A. K.; DACHER, M.; GAUTHIER, M.; ARMENGAUD, C. Effects of sub-lethal doses of fipronil on the behavior of the honeybee (*Apis mellifera*). **Pharmacology Biochemistry and Behavior**, v. 82, n. 1, p. 30-39, 2005.
- JOHANSEN, C. A.; MAYER, D. F. Pollinator protection. **A bee and pesticide handbook**. Cheshire: Wicwas Press, 1990.
- KEVAN, P. G. Pollinators as bioindicators of the state of the environment: species, activity and diversity. **Agriculture, Ecosystems and Environment**, v. 74, n. 1-3, p. 373-393, 1999.
- MALASPINA, O.; STORT, A. C. As abelhas e os pesticidas. **Apicultura no Brasil**, v. 2, n. 10, p. 42-45, 1985.
- MCKENZIE, K. E.; WINSTON, M. L. Effects of sub-lethal exposure to diazinon on longevity and temporal division of labor in the honey bee (Hymenoptera: Apidae). **Journal of Economic Entomology**, v. 82, n. 1, p. 75-82, 1989.
- MENZEL, R.; MÜLLER, U. Learning and memory in honeybees: from behavior to neural substrates. **Annuary of Review Neuroscience**, v. 19, n. 1, p. 379-404, 1996.
- MIRANDA, J. E.; DESBONSI, H. M.; NOGUEIRA-COUTO, R. H.; DE BORTOLI, S. A.; KATO, M. J.; BOLZANI, V. S.; FURLAN, M.; MCKENZIE, K. E.; WINSTON, M. L. Susceptibility of *Apis mellifera* (Hymenoptera, Apidae) to pellitorine, na amide isolated from *Piper tuberculatum* (Piperaceae). **Apidologie**, v. 34, n. 4, p. 409-415, 2003.
- MULLIN, C. A.; FRAZIER, M.; FRAZIER, J. L. High levels of miticides and agrochemicals in North American apiaries: Implications for honey bee health. **Plos One**, v. 5, n. 3, p. 1-19, 2010.
- PAOLETTI, M. G. **Invertebrate biodiversity as bioindicators of sustainable landscapes**. Practical use of invertebrates to assess sustainable landuse. Amsterdam: Elsevier, 1999.
- PHAM-DELÈGUE, M. H.; DECOURTYE, A.; KAISER, L.; DEVILLERS, J. Behavioural methods to assess the effects of pesticides on honey bees. **Apidologie**, v. 33, n. 5, p. 425-432, 2002.
- RICHARDS, K. W.; KEVAN, P. G. Aspects of bee biodiversity, crop pollination, and conservation in Canada. In: KEVAN, P.; IMPERATRIZ FONSECA, V. L. (Ed.). **Pollinating bees - the conservation link between agriculture and nature**. Brasília: Ministério do Meio Ambiente, 2002. p. 77-94.
- ROMERO, R. R.; DESNEUX, N.; DECOURTYE, A.; CHAFFIOLD, A.; PHAM-DELÈGUE, M. H. Does cryab protein affect learning performances of the honey bee *Apis*

mellifera L. (Hymenoptera, Apidae). **Ecotoxicology and Environmental Safety**, v. 70, n. 2, p. 327-333, 2008.

SCHEINER, R.; PAGE, R. E.; ERBER, J. Sucrose responsiveness and behavioral plasticity in honey bees (*Apis mellifera*). **Apidologie**, v. 35, n. 2, p. 133-142, 2004.

SUCHAIL, S.; DEBRAUWER, L.; BEIZUNCES, L. P. Metabolism of imidacloprid in *Apis mellifera*. **Pest Management Science**, v. 60, n. 3, p. 291-296, 2003.

SUCHAIL, S.; GUEZ, D.; BELZUNCES, L. P. Characteristics of imidacloprid toxicity in two *Apis mellifera* subspecies. **Environmental Toxicology and Chemistry**, v. 19, n. 7, p. 1901-1905, 2000.

ZAR, J. H. **Bioestatistical analysis**. New Jersey: Prentice Hall, 1996.

Received on September 21, 2012.

Accepted on December 14, 2012.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.