

## Relative Growth and Sexual Maturity of the Stone Crab *Menippe nodifrons* Stimpson, 1859 (Brachyura, Xanthoidea) in Southeastern Brazil

Giovana Bertini<sup>1\*</sup>, Adriane Araújo Braga<sup>2</sup>, Adilson Fransozo<sup>2</sup>, Michéle de Oliveira Dias Alves Corrêa<sup>2</sup> and Fulvio Aurélio de Moraes Freire<sup>2</sup>

<sup>1</sup>Universidade Estadual Paulista; Campus Experimental de Registro; Rua Tamekichi Takano, 5; 11900-000; gibertini@registro.unesp.br; Registro - SP - Brasil. <sup>2</sup>Departamento de Zoologia; Instituto de Biociências; UNESP; C. P. 510; fransozo@ibb.unesp.br; 18.618-000; Botucatu - SP - Brasil

### ABSTRACT

The relative growth and size at onset of morphological sexual maturity of the stone crab *Menippe nodifrons* were investigated. A total of 399 crabs was captured on Praia Grande and Tenório beaches at Ubatuba. Carapace width (CW) and length, cheliped propodus length and height, abdomen width in females, and gonopod length in males were recorded. In females, the abdominal width showed negative allometry for juveniles and positive allometry for adults; the puberty molt occurred at 31.6 mm CW. In males, the size at onset of morphological sexual maturity was estimated as 29.7 mm CW; the gonopod growth showed positive allometry for juveniles, and an isometric relationship for adults. The gonopod length and the abdominal width were the most appropriate morphometric variables to estimate size at onset of sexual maturity in this stone crab.

**Key words:** Growth, Sexual Maturity, *Menippe*, Xanthoidea

### INTRODUCTION

Stone crabs of the genus *Menippe* occur in tropical and subtropical regions. They support important fisheries in the United States, Mexico and Cuba (Ehrhardt, 1990). In many localities, only their claws are widely sold; the animals are returned to their environment, where they are later recaptured. The biology, life history, and fisheries of stone crabs have been investigated mainly for *Menippe mercenaria* (Say, 1818) and *M. adina* Williams and Felder, 1986, which grow large and are widely exploited commercially (Simonson and Steele, 1981, Wilber, 1989, Ehrhardt, 1990, Landry, 1992, Caldwell, 1992, Stuck and Perry, 1992). Only one

species of this genus, *M. nodifrons* Stimpson, 1859, occurs along the coast of Brazil. According to Oshiro (1999), *M. nodifrons* is little exploited commercially, probably because it is smaller than the North American species. Therefore, it is caught only by a few tourists and amateur fishermen. Studies of relative growth and morphological sexual maturity are important parameters for determining the reproductive potential of a species, and for the management of populations of commercial interest (Campbell and Eagles, 1983; Guerrero-Ocampo et al., 1998). Certain species of xanthoid crabs have been investigated in relation to these subjects, notably by Caldwell (1992) with *M. mercenaria*; Finney and Abele (1981) with

\* Author for correspondence

*Trapezia ferruginea* Latreille, 1828; Huber (1985) with *Trapezia ferruginea*, *T. corallina* Gerstaecker, 1857 and *T. formosa* Smith, 1869; Vannini and Gherardi (1988) with *Eriphia smithi* MacLeay, 1838; Góes and Fransozo (1997) with *Eriphia gonagra* (Fabricius, 1781) and Guimarães and Negreiros-Fransozo (2002) with *Eurytium limosum* (Say, 1818).

Knowledge of the biology of *M. nodifrons* is limited to studies of its larval development by Scotto (1979), juvenile development and population biology and habitat utilization by Fransozo et al. (1988 and 1999) and aspects of its reproduction by Oshiro (1999). Further investigations may elucidate fundamental questions regarding the life history of this stone crab and may make possible its commercial exploitation. The goal of this study was to analyze the relative growth of *M. nodifrons*, focusing on the morphometric relationships between the carapace width and the dimensions of certain appendages. Also, the transition intervals between the immature and mature growth phases in allometric relationships were assessed, in order to estimate the size at onset of morphological maturity.

## MATERIAL AND METHODS

Crabs were collected manually from 1997 to 1999, during low tide, on the rocky shores of Praia Grande and Tenório beaches in Ubatuba, state of São Paulo, southeastern Brazil. Body dimensions were measured with a caliper to the nearest 0.01 mm. The following dimensions were selected for morphometric analyses: carapace width (CW), carapace length (CL), left cheliped propodus length (LPL) and height (LPH), right cheliped propodus length (RPL) and height (RPH), abdomen width (AW) of the fifth somite in females, and gonopod length (GL) (the exopodite of the first pair of pleopods) in males. The relative frequency of adults (%) in each size class was plotted on a graph, which was fitted as a sigmoid curve following the result of the logarithmic


equation  $y = \frac{1}{1 + e^{r(CW - CW_{50})}}$ , where  $CW_{50}$  = carapace width at which 50% of the individuals reach sexual maturity, and  $r$  = slope of the curve. The adjusted equation was fitted by the least-squares regression method (Vazzoler, 1996).

Carapace width was considered the independent variable, and the allometric equation  $Y = a \cdot x^b$  for each sex and maturation phase was calculated using the least-squares regression method (Hartnoll, 1982). The values of  $b$  define the type of allometry ( $b = 1$ : isometry;  $b < 1$ : negative allometry;  $b > 1$ : positive allometry). Departures from isometry ( $H_0: b = 1$ ) were tested using Student's t-test on the slope values obtained ( $\alpha = 0.05$ ). An analysis of covariance (ANCOVA) was performed to compare slopes and intercepts of the regressions obtained within each allometric relationship (Zar, 1996).

## RESULTS

A total of 399 specimens of *M. nodifrons* was captured during the study period. Their sizes ranged from 5.0 to 69.6 mm CW for males ( $N = 195$ ), and from 3.6 to 82.5 mm CW for females ( $N = 204$ ). The size at which 50% of the population was morphologically mature is shown in Fig. 1. The estimated sizes at onset of sexual maturity were 29.7 and 31.6 mm CW for males and females, respectively.

All of the relationships, numbers of specimens analyzed, and linearized allometric expressions followed by the power functions, the  $r^2$  values, and the allometric level of each relationship are given in Table I. The overall CL to CW ratio was 0.70. The CL vs. CW regression plotted for all individuals indicated isometric growth between these variables. Because the intercepts and slopes of a given allometric relationship did not differ between males and females, a single regression line could be fitted for both sexes combined (Table I). For the regression calculated for the male chelipeds, the only difference in relative growth was observed in the LPH vs. CW relationship, in which the juveniles showed isometric growth ( $b = 1.02$ ), and the adults showed positive allometric growth ( $b = 1.23$ ) (Fig. 2).




**Figure 1** - *Menippe nodifrons*. Logarithmic curve between the carapace width (CW) and percentage of adults, for males and females.

**Table 1** - Relationships between morphometric parameters and carapace width in *Menippe nodifrons* by means the allometric technique.

Variables	Group	N	Equation $Y = a \cdot x^b$	Linear transformation $\ln y = \ln a + b \cdot \ln x$	$r^2$	Allometry level*
CL	Total	399	$CL = 0.757 \cdot CW^{0.98}$	$\ln CL = -0.278 + 0.98 \ln CW$	0.99	=
	TM	195	$CL = 0.754 \cdot CW^{0.98}$	$\ln CL = -0.282 + 0.98 \ln CW$	0.99	=
	TF	204	$CL = 0.757 \cdot CW^{0.98}$	$\ln CL = -0.278 + 0.98 \ln CW$	0.99	=
RPL	TM	170	$RPL = 0.699 \cdot CW^{1.02}$	$\ln RPL = -0.358 + 1.02 \ln CW$	0.98	+
	JF	95	$RPL = 0.829 \cdot CW^{0.95}$	$\ln RPL = -0.188 + 0.95 \ln CW$	0.96	-
	AF	74	$RPL = 0.475 \cdot CW^{1.12}$	$\ln RPL = -0.744 + 1.12 \ln CW$	0.89	+
RPH	TM	173	$RPH = 0.377 \cdot CW^{0.99}$	$\ln RPH = -0.975 + 0.99 \ln CW$	0.97	=
	JF	97	$RPH = 0.424 \cdot CW^{0.95}$	$\ln RPH = -0.857 + 0.95 \ln CW$	0.95	-
	AF	69	$RPH = 0.183 \cdot CW^{1.18}$	$\ln RPH = -1.7 + 1.18 \ln CW$	0.87	+
LPL	TM	154	$LPL = 0.569 \cdot CW^{1.07}$	$\ln LPL = -0.564 + 1.07 \ln CW$	0.98	+
	TF	162	$LPL = 0.639 \cdot CW^{1.03}$	$\ln LPL = -0.448 + 1.03 \ln CW$	0.99	+
LPH	JM	120	$LPH = 0.313 \cdot CW^{1.02}$	$\ln LPH = -1.162 + 1.02 \ln CW$	0.96	=
	AM	27	$LPH = 0.152 \cdot CW^{1.23}$	$\ln LPH = -1.880 + 1.23 \ln CW$	0.88	+
	TF	166	$LPH = 0.319 \cdot CW^{1.02}$	$\ln LPH = -1.142 + 1.02 \ln CW$	0.98	+
GL	JM	86	$GL = 0.039 \cdot CW^{1.63}$	$\ln GL = -3.244 + 1.63 \ln CW$	0.92	+
	AM	42	$GL = 0.397 \cdot CW^{0.94}$	$\ln GL = -0.924 + 0.94 \ln CW$	0.93	=
AW	JF	117	$AW = 0.231 \cdot CW^{0.95}$	$\ln AW = -1.465 + 0.95 \ln CW$	0.93	-
	AF	84	$AW = 0.047 \cdot CW^{1.47}$	$\ln AW = -3.048 + 1.47 \ln CW$	0.92	+

\* *t* test;  $p < 0.05$

(N = number of individuals; TM = total males; JM = juvenile males; AM = adult males; TF = total females; JF = juvenile females; AF = adult females;  $r^2$  = determination coefficient; CL = carapace length; RPL = right cheliped propodus length; RPH = right chelar propodus height; LPL = left cheliped propodus length; LPH = left cheliped propodus height; GL = gonopod length; AW = abdomen width).




**Figure 2** - *Menippe nodifrons*. Scatter plot between the carapace width (CW) and left cheliped propodus height (LPH) in males.


For females, the regression calculated for the right propodus (RPL vs. CW and RPH vs. CW) indicated two statistically different regression lines between the maturation phases ( $p < 0.05$ ). There was negative allometric growth in the early phase, and positive allometric growth in the adult phase (Figs. 3, 4).

The relationships which best indicated the change in the allometric coefficient between juveniles and adults were AW vs. CW for females (Fig. 5) and


GL vs. CW for males (Fig. 6). These regressions were statistically different between the maturation phases. Males showed positive allometric growth during the juvenile phase, and isometry during the adult phase. Females showed negative allometric growth during the juvenile phase, and positive allometric growth after the puberty molt.




**Figure 3** - *Menippe nodifrons*. Scatter plot between the carapace width (CW) and right cheliped propodus length (RPL) in females.



**Figure 4** - *Menippe nodifrons*. Scatter plot between the carapace width (CW) and right cheliped propodus height (RPH) in females.



**Figure 5** - *Menippe nodifrons*. Scatter plot between the carapace width (CW) and abdomen width (AW) in females.



**Figure 6** - *Menippe nodifrons*. Scatter plot between the carapace width (CW) and gonopod length (GL) in males.

## DISCUSSION

The size at the onset of sexual maturity and the number of instars following the puberty molt provide important information on the reproductive biology of brachyuran crabs (Pinheiro and Fransozo, 1998). This knowledge is essential for managing commercially exploited populations (Campbell and Eagles, 1983). In general, when two dimensions of the carapace of brachyurans are correlated, growth tends to be isometric during ontogeny, and does not change much with the molt to puberty. However, Fransozo et al. (1988) found that in *M. nodifrons* there were variations in the length-width proportions of the carapace up to juvenile stage V, which coincides with the appearance of secondary sex characters, and only after this instar these proportions stabilize with the length about 70% of the width. This variation in the CL/CW proportion also occurred in the juvenile stages of other xanthoids, as observed by Fransozo and Negreiros-Fransozo (1987) in *Eriphia gonagra* and *Eurypanopeus abbreviatus* (Stimpson, 1860). Savage and Sullivan (1978) in Florida and Caldwell (1992) in South Carolina also observed that the carapace length was proportionally longer than the carapace width in *M. mercenaria*.

In the males of *M. nodifrons*, the relationships of the chelipeds were not good indicators of the molt to puberty. This was also found for other xanthoids by Góes and Fransozo (1997) with *Eriphia gonagra*, and by Guimarães and Negreiros-Fransozo (2002) with *Eurytium limosum*. However, most of the relationships showed positive allometric growth, indicating that these structures had an important role in agonistic behaviors. As described by Porter (1960), the males of *M. mercenaria* used their chelipeds to protect the entrances of holes occupied by pre-molt females, prior to copulation. Savage and Sullivan (1978) also found that the propodus length of *M. mercenaria* showed positive allometric growth, so that the propodi of adult males were proportionately larger than those of females. Caldwell (1992) found similar results for this species, and noted that because of this difference, the males could be harvested commercially at a smaller carapace width than the females.

Among xanthoids, this sexual dimorphism may be greatly accentuated, as occurs in *Eriphia smithi*, in

which the presence of a powerful chela is related to the conquest and defense of suitable burrows. The number of these burrows, especially copulatory burrows, may be a limiting factor on the rocky shores where this species lives (Vaninni, 1987; Vaninni and Gherardi, 1988).

The regression analysis CW vs. GL for *M. nodifrons* indicated positive allometric growth during the juvenile phase. With the passage to the adult phase, growth was isometric. This result is similar to observations by Hartnoll (1965) for *Aratus pisonii* (H. Milne Edwards, 1837) and by Góes and Fransozo (1997) for *Eriphia gonagra*. In other species, the positive allometric relationship only decreased slightly, as observed by Vannini and Gherardi (1988) for *Eriphia smithi*, by Cobo and Fransozo (1998) for *Goniopsis cruentata* (Latreille, 1803), and by Fransozo et al. (2002) for *Ocypode quadrata* (Fabricius, 1787). Guimarães and Negreiros-Fransozo (2002) established that gonopod growth in *Eurytium limosum* passed from positively allometric in the juvenile phase, to negative in the adults.

This differential growth of the male gonopod is related to the ability of these individuals to copulate successfully (Hartnoll, 1974). The decrease in growth at the puberty molt may be adaptive, allowing males to copulate with females of a wide range of sizes. However, in this species, no standardization (*sensu* Hartnoll, 1965, 1972) was observed. This suggested that the remarkable positive allometry in the juvenile phase of *M. nodifrons* could confer an advantage, in that males could reach sexual maturity at a smaller size than could females, allowing them to compete successfully with other males to copulate with the mature females.

The ontogenic changes in the abdomen width of females of *M. nodifrons* are related to the capacity of the female to incubate the egg mass, following the common pattern of most brachyuran species. This allometric relationship is often used to detect the molt to puberty in the females (Hartnoll, 1965; Finney and Abele, 1981; Pinheiro and Fransozo, 1998; Fransozo et al., 2002).

The estimated value (31.6 mm CW) for the beginning of morphological sexual maturity of the females of *M. nodifrons* was lower than that obtained by Oshiro (1999). Oshiro estimated that the mean size at the first gonadal maturation of this species was 37.8 mm CW, and the smallest ovigerous female caught in the area of her study

(Sepetiba Bay, Rio de Janeiro) was 38 mm CW. This suggested that these two events did not occur in synchrony in this species.

In another xanthoid, *Eriphia gonagra*, studied by Góes (2000), morphological maturity occurred practically in synchrony with physiological maturity in both sexes. Nevertheless, these events commonly occurred asynchronously in brachyurans, as for instance in the portunids *Liocarcinus puber* (Linnaeus, 1767), studied by Choy (1988); *Callinectes ornatus* Ordway, 1863, by Haefner (1990); and *Ovalipes stephensoni* Williams, 1976, by Haefner (1985). In all these cases, the females reached morphological maturity before they reached physiological maturity.

*Menippe nodifrons* appears to mature precociously in relation to other species of the genus *Menippe*. *M. mercenaria* reached sexual maturity at 60 mm CW, according to the work of Sullivan (1979) who reported the smallest ovigerous female caught in southwest Florida, and to the analysis of Wilber (1992) of the presence of spermatozooids in the seminal receptacle in individuals in the northeastern Gulf of Mexico. In South Carolina, Caldwell (1992) reported that the females of this species initiated maturation at 55 mm and the males at 60 mm CW; the largest individuals caught were 135 mm CW. This later maturation in *M. mercenaria* than in *M. nodifrons* was probably related to the larger size reached by the former species.

Although *M. nodifrons* is smaller than *M. mercenaria* and *M. adina*, raising this commercially species could be interesting. Oshiro et al. (1998) estimated that the average meat yield from *M. nodifrons* was higher than from other species of crabs which were commercially important in Brazil, such as *Ucides cordatus* (Linnaeus, 1763) and *Cardisoma guanhumi* Latreille, 1825.

## ACKNOWLEDGEMENTS

The authors are grateful for financial support during collections (# 94/4878-8; 97/12108-6; 97/12107-0; 98/3134-6). We thank many colleagues from the NEBECC group who helped with sampling and laboratory analysis, to Dr. Maria Lucia Negreiros Fransozo for her helpful comments on the early draft of this manuscript and Instituto Brasileiro do Meio Ambiente e dos

Recursos Naturais Renováveis (IBAMA) for permission during the samplings.

## RESUMO

O objetivo do presente estudo foi determinar o crescimento relativo e a maturidade sexual morfológica de *Menippe nodifrons*. As coletas foram realizadas na Praia Grande e Tenório, Ubatuba, São Paulo. Os caranguejos foram separados quanto ao sexo e mensurados na região da largura (LC) e comprimento da carapaça, comprimento e altura dos própodos quelares direito e esquerdo, largura do abdome (LA) nas fêmeas e comprimento do gonopódio (CG) nos machos. Obteve-se 399 indivíduos, sendo 195 machos e 204 fêmeas. Os machos atingiram a maturidade sexual com 29,7mm LC e as fêmeas com 31,6mm LC. Para as fêmeas a melhor relação que indicou a muda da puberdade foi LA vs. LC, sendo que o crescimento foi alométrico negativo na fase jovem e alométrico positivo após a muda da puberdade. Para os machos foi CG vs. LC evidenciando crescimento alométrico positivo na fase jovem e isométrico na fase adulta.

## REFERENCES

- Caldwell, M. A. (1992), Aspects of the biology of the stone crab, *Menippe mercenaria* (Say), from South Carolina, with comments on the South Carolina stone crab fishery. *Flo. Mar. Res. Publ.*, **50**: 90-106.
- Campbell, A., and Eagles, M. D. (1983), Size at maturity and fecundity of rock crabs, *Cancer irroratus*, from the Bay of Fundy and southwestern Nova Scotia. *Fish. Bul.*, **81**(2): 357-362.
- Choy, S. C. (1988), Reproductive biology of *Liocarcinus puber* and *L. holsatus* (Decapoda, Brachyura, Portunidae) from the Gower Peninsula, South Wales. *Mar. Ecol.*, **9**(3): 227-241.
- Cobo, V. J., and Fransozo, A. (1998), Relative growth of *Goniopsis cruentata* (Crustacea, Brachyura, Grapsidae), on the Ubatuba region, São Paulo, Brazil. *Iheringia, Sér. Zool.*, **(84)**: 21-28.
- Ehrhardt, N. M. (1990), Mortality and catchability estimates for the stone crab (*Menippe mercenaria*) in Everglades National Park. *Bull. Mar. Sci.*, **46**(2): 324-334.

- Finney, W. C., and Abele, L. G. (1981), Allometric variation and sexual maturity in the obligate coral commensal *Trapezia ferruginea* Latreille (Decapoda, Xanthidae). *Crustaceana*, **41**(2): 113-130.
- Fransozo, A., and Negreiros-Fransozo, M. L. (1987), Morfologia dos primeiros estágios juvenis de *Eriphia gonagra* (Fabricius, 1781) e *Eurypanopeus abbreviatus* (Stimpson, 1860) (Crustacea, Decapoda, Xanthidae) obtidos em laboratório. *Papéis Avulsos de Zool.*, **36**(22): 257-277.
- Fransozo, A., Bertini, G., and Correa, M. O. D. (1999), Population biology and habitat utilization of the stone crab *Menippe nodifrons* in the Ubatuba region, São Paulo, Brazil. *Crust. Issues*, **12**: 275-281.
- Fransozo, A., Negreiros-Fransozo, M. L., and Hiyodo, C. M. (1988), Développement juvénile de *Menippe nodifrons* Stimpson, 1859 (Crustacea, Decapoda, Xanthidae) au laboratoire. *Rev. Hydrobiol. Trop.*, **21**(4):297-308.
- Fransozo, A., Negreiros-Fransozo, M. L., and Bertini, G. (2002), Morphometric study of the ghost crab *Ocypode quadrata* (Fabricius, 1787) (Brachyura, Ocypodidae) from Ubatuba, São Paulo, Brazil. In: Escobar-Briones, E. and F. Alvarez (eds.) *Modern Approaches to the Study of Crustacea*, pp.189-195.
- Góes, J. M. (2000), *Biologia do caranguejo Eriphia gonagra (Fabricius, 1781) (Crustacea, Brachyura, Xanthidae) na região de Ubatuba, São Paulo*. Universidade Estadual Paulista (UNESP), Instituto de Biociências, Botucatu, SP, 175p. [Doctoral Thesis].
- Góes, J. M., and Fransozo, A. (1997), Relative growth of *Eriphia gonagra* (Fabricius, 1781) (Crustacea, Decapoda, Xanthidae) in Ubatuba, State of São Paulo, Brazil. *Nauplius*, **5**(2): 85-98.
- Guerrero-Ocampo, C. M., Negreiros-Fransozo, M. L., and Costa, T. M. (1998), Comparação do peso dos quelípodos e crescimento em duas espécies de "siris" do gênero *Callinectes* (Brachyura, Portunidae). *Braz. Arch. Biol. Technol.*, **41**(4): 483-488.
- Guimarães, F. J., and Negreiros-Fransozo, M. L. (2002), Sexual maturity of *Eurytium limosum* (Say, 1818) from a subtropical mangrove in Brazil. In: Escobar-Briones, E. and F. Alvarez (eds.) *Modern Approaches to the Study of Crustacea*, pp. 157-161.
- Haefner, P. A. Jr. (1985), Morphometry, reproduction, diet and epizotes of *Ovalipes stephensi* Williams, 1976 (Decapoda, Brachyura). *J. Crustacean Biol.*, **5**(4): 658-672.
- Haefner, P. A. Jr. (1990), Morphometry and size at maturity of *Callinectes ornatus* (Brachyura, Portunidae) in Bermuda. *Bull. Mar. Sci.*, **46**(2): 274-286.
- Hartnoll, R. G. (1965), Notes on the marine grapsid crabs of Jamaica. *Proc. Linn. Soc. Lond.*, **176**: 113-147.
- Hartnoll, R. G. (1972), The biology of the burrowing crab, *Corystes cassvelaunus*. *Bijdr. Dierk.*, **42**(2): 139-155.
- Hartnoll, R. G. (1974), Variation in growth pattern between some secondary sexual characters in crabs (Decapoda, Brachyura). *Crustaceana*, **27**: 131-136.
- Hartnoll, R. G. (1982), Growth. In: D. E. Bliss (ed.) *The Biology of Crustacea: Embryology, Morphology and Genetics*, Vol. **3**, pp. 111-196.
- Huber, M. E. (1985), Allometric growth of the carapace in *Trapezia* (Brachyura, Xanthidae). *J. Crustacean Biol.*, **5**(1): 79-83.
- Landry, A. M. Jr. (1992), Characterization and fishery development potential of Galveston Bay, Texas, stone crab (*Menippe adina*) stocks. *Flo. Mar. Res. Publ.*, **50**: 67-73.
- Oshiro, L. M. Y. (1999), Aspectos reprodutivos do caranguejo guaia, *Menippe nodifrons* Stimpson (Crustacea, Decapoda, Xanthidae) da Baía de Sepetiba, Rio de Janeiro, Brasil. *Revta. Brasil. Zool.*, **16**(3): 827-834.
- Oshiro, L. M. Y., Silva, R., Souza, E. P., and Silveira, C. M. (1998), Rendimento de carne em três caranguejos braquiúros (Crustacea, Decapoda) da Baía de Sepetiba, RJ. Paper presented at XXII Congresso Brasileiro de Zoologia, Recife (PE) Brasil. p 77.
- Pinheiro, M. A. A., and Fransozo, A. (1998), Sexual maturity of the speckled swimming crab *Arenaeus cribrarius* (Lamarck, 1818) (Decapoda, Brachyura, Portunidae) in the Ubatuba littoral, São Paulo State, Brazil. *Crustaceana*, **71**(6): 444 - 452.
- Porter, H. J. (1960), Zoal stages of the stone crab, *Menippe mercenaria* Say. *Chesapeake Sci.*, **1**: 168-177.
- Savage, T., and Sullivan, J. R. (1978), Growth and claw regeneration of the stone crab, *Menippe mercenaria*. *Flo. Mar. Res. Publ.*, number 32. 23 pp.
- Scotto, L. E. (1979), Larval development of the Cuban stone crab, *Menippe nodifrons* (Brachyura, Xanthidae), under laboratory conditions with notes on the status of the family Menippidae. *Fish. Bull.*, **77**(2): 359-386.
- Simonson, J. L., and Steele, P. (1981), Cheliped asymmetry in the stone crab, *Menippe mercenaria*, with notes on claw reversal and regeneration. *Northeast Gulf Sci.*, **5**(1): 21-30.



- Stuck, K. C., and Perry, H. M. (1992), Life history characteristics of *Menippe adina* in Mississippi coastal waters. *Flo. Mar. Res. Publ.*, **50**: 82-98.
- Sullivan, J. R. (1979), The stone crab, *Menippe mercenaria*, in the southwest Florida fishery. *Flo. Mar. Res. Publ.*, number **36**. 37 pp.
- Vannini, M. (1987), Notes on the ecology and behaviour of the pebble crab *Eriphia smithi* MacLeay (Decapoda, Brachyura). *Monit. Zool. Ital. Suppl.*, **22**: 383-410.
- Vannini, M., and Gherardi, F. (1988), Studies on the pebble crab, *Eriphia smithi* MacLeay, 1838 (Xanthoidea, Menippidae): patterns of relative growth and population structure. *Trop. Zool.*, **1**: 203-216.
- Vazzoler, A. E. A. M. (1996), *Biologia da reprodução de peixes teleósteo: teorias e prática*. Maringá: EDUEM. 169p.
- Wilber, D. H. (1989), Reproductive biology and distribution of stone crabs (Xanthidae, *Menippe*) in the hybrid zone in the northeastern Gulf of Mexico. *Mar. Ecol. Prog. Ser.*, **52**: 235-244.
- Wilber, D. H. (1992), Observations on the distribution and mating patterns of adult stone crabs (Genus *Menippe*) on the northern Gulf Coast of Florida. *Flo. Mar. Res. Publ.*, **50**: 10-16.
- Zar, J.H.(1996), *Biostatistical analysis*. Prentice-Hall, Upper Saddle River.

Received: June 08, 2005;  
Revised: February 15, 2006;  
Accepted: October 25, 2006.

PÁGINA  
EM  
BRANCO