



# The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



## Case report

# Vertebral osteomyelitis and epidural abscesses caused by *Prevotella oralis*: a case report

Hemant Goyal<sup>a,\*</sup>, Shitij Arora<sup>a</sup>, Sneha Mishra<sup>a</sup>, Syed Jamil<sup>b</sup>, Uday Shah<sup>a</sup>

<sup>a</sup> Division of Internal Medicine, Wyckoff Heights Medical Center, Brooklyn, NY, USA

<sup>b</sup> Department of Infectious Diseases, Wyckoff Heights Medical Center, Brooklyn, NY, USA

### ARTICLE INFO

#### Article history:

Received 30 March 2012

Accepted 11 July 2012

Available online 8 November 2012

#### Keywords:

Anaerobes

*Prevotella*

Vertebral osteomyelitis

### ABSTRACT

An interesting case of pyogenic vertebral osteomyelitis with multiple epidural abscesses caused by non-pigmented *Prevotella oralis* is reported. The patient was a 68-year-old female who presented to the emergency room (ER) with severe pain and tenderness in her lower back with fever. She had recently undergone esophagogastroduodenoscopy (EGD) for complaints of esophageal reflux, which showed submucosal cyst in the esophagus. Magnetic resonance imaging (MRI) of the thoracic spine revealed multiple spinal epidural abscesses with signal enhancement at the level of T6 and T7, suggestive of vertebral osteomyelitis. Two blood cultures drawn one hour apart grew *Prevotella oralis*. The body fluid aspirated from the abscesses was also positive for the anaerobic commensal *P. oralis*. Necrosis associated with the submucosal cyst was implicated as the cause of sepsis and osteomyelitis due to this organism.

© 2012 Elsevier Editora Ltda. All rights reserved.

## Introduction

Vertebral osteomyelitis is a rare cause of back pain in adults. The usual implicated organisms are *Staphylococcus aureus*, beta hemolytic *Streptococcus*, and Viridans *Streptococci*.<sup>1</sup> Anaerobic Gram-negative bacilli (AGNB) are a rare cause of vertebral osteomyelitis and amongst the AGNB, *Prevotella* is an extremely rare etiological agent. The *Prevotella* genus is further classified into pigmented and non-pigmented species, based on production of pigmented colonies on blood agar.<sup>2</sup> Here, an interesting case of pyogenic vertebral osteomyelitis with multiple epidural abscesses caused by non-pigmented *Prevotella oralis* is reported. To the authors' knowledge, this is the first reported case of pyogenic vertebral osteomyelitis caused by non-pigmented *P. oralis*.

## Case presentation


A 68-year-old female was referred by her primary physician to the ER for evaluation and management of severe pain in her back along the rib cage, which radiated towards the anterior abdominal wall. She reported that the pain was 10/10 in intensity, sharp in quality, continuous, and increased with movements. She also complained of mild fever with lethargy. She denied weakness of the limbs and fecal or urinary incontinences. The patient had past medical history of hypertension and gastroesophageal reflux disease (GERD). She had undergone esophagogastroduodenoscopy (EGD) two weeks prior, which was positive for submucosal cyst. She had diffuse tenderness over her back, which was more significant over the lower thoracic vertebrae. Laboratory data showed a WBC count

\* Corresponding author at: Wyckoff Heights Medical Center, Department of Internal Medicine, Room # C-406, 374, Stockholm Street, Brooklyn, NY, 11237, USA.

E-mail address: doc.hemant@yahoo.com (H. Goyal).

1413-8670/\$ – see front matter © 2012 Elsevier Editora Ltda. All rights reserved.

<http://dx.doi.org/10.1016/j.bjid.2012.10.008>



**Fig. 1 – (A) Non-contrast T1 magnetic resonance imaging (MRI) of thoracic spine revealing signal abnormalities of the bodies of T6 and T7 vertebrae consistent with osteomyelitis. (B) Contrast-enhanced T1 MRI showing multiple abscesses.**

of 18,000, with neutrophilic predominance of 91.4%. Erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) level were 120 mm/hour and 207.5 mg/L respectively. Magnetic resonance imaging (MRI) of the thoracic spine (Fig. 1A and B) was performed, which revealed an abscess on the posterior aspect of the spinal cord displacing the spinal cord anteriorly at the level of T5 and T6. Another abscess anterior to the spinal cord, which was displacing the spinal cord posteriorly at the level of T6 and T7 extending to T8, was also observed. There was signal abnormality of the bodies of T6 and T7 vertebrae consistent with osteomyelitis. The patient was started empirically on intravenous vancomycin and doripenem, with pending blood cultures. Two blood cultures isolated from the patient one hour apart at the time of admission grew Gram-negative rods positive for anaerobic bacteria, *P. oralis*. The body fluid aspirated from the abscess was also positive for

*P. oralis*. The API® 20A anaerobic system (bioMérieux Inc.) was used in identification of bacteria. Antibiotics were changed to ampicillin-sulbactam and metronidazole after the availability of culture reports. The patient was later transferred to a skilled nursing facility with a long-term intravenous access, and later responded well to the treatment after antibiotic therapy for eight weeks.

**Discussion**

*P. oralis* was first described as *Bacteroides oralis* as a new species isolated from the oral cavity of a man which was distinguished in many ways from *Bacteroides fragilis* by Loesche et al.<sup>3</sup> It was later reclassified and the new genus *Prevotella* was suggested by Shah and Collins.<sup>2</sup> In most studies of osteomyelitis

**Table 1 – Clinical characteristics of the patients with vertebral osteomyelitis caused by *Prevotella* spp. previously described in the literature.**

Authors	Patient characteristics					
	Age	Gender	<i>Prevotella</i> species	Site of osteomyelitis	Risk factors	Treatment outcome
Surbled et al. <sup>4</sup>	44	M	<i>P. melaninogenica</i>	L5-S1	None	Favorable
Salavert et al. <sup>5</sup>	27	F	<i>P. melaninogenica</i>	L2-L3	IV drug use	Favorable
Fukuoka et al. <sup>6</sup>	60	M	<i>P. intermedia</i>	T7-T8	Age	Favorable
Schöber et al. <sup>7</sup>	45	F	<i>P. intermedia</i>	L1-L2	None	Favorable
Mukhopadhyay et al. <sup>8</sup>	35	F	<i>P. melaninogenica</i>	S1	Dental cleaning	Favorable
Salliot et al. <sup>9</sup>	62	M	<i>Prevotella</i> spp. (the species could not be identified)	L5-S1	Age, steroids, spodyloarthropathy	Favorable
Purushothaman et al. <sup>10</sup>	74	F	<i>Prevotella</i> spp. (the species could not be identified)	L3-L4	Age, ovarian malignancy	Favorable
Huang et al. <sup>11</sup>	73	F	<i>P. melaninogenica</i>	T12-L3	Age, diabetes	Patient died

caused by anaerobic bacteria, the common implicated organisms are *Bacteroides* spp., *Fusobacterium* spp., and *Clostridium* spp. On review of the PubMed database, with the search words 'Prevotella and osteomyelitis', 'Prevotella and vertebral', and 'Prevotella and spondylodiscitis', eight cases of human vertebral osteomyelitis were retrieved that were caused by different species of *Prevotella*; however, none of them was caused by *P. oralis*.<sup>4-11</sup> Clinical characteristics of these patients are summarized in Table 1.

*P. oralis* as an isolated microbiologic agent in vertebral osteomyelitis is an unknown entity. *P. oralis* is a major commensal on the mucosal surfaces of the body such as the oral cavity, gastrointestinal tract, and vagina. Previously, it has been reported to cause periodontal infections, urinary tract infections, and rarely intra-abdominal abscesses. The source of sepsis in this patient could have been the breach in mucocutaneous barrier during the EGD, which displaced the oral flora into the blood stream. The possibility of necrosis associated with the submucosal cyst could also be implicated in the septicemia caused by *P. oralis*, which ultimately led to seeding of the organism in the bones and subsequent development of osteomyelitis. The present case highlights the importance of therapy directed against anaerobic pathogens in appropriate clinical settings.

---

### Conflict of interest

All authors declare to have no conflict of interest.

---

### Acknowledgements

The authors sincerely thank Monica Chiang, MD, for translation of reference No. 5 from Spanish to English.

---

### REFERENCES

1. Kumar A, Sandoe J, Kumar N. Three cases of vertebral osteomyelitis caused by *Streptococcus dysgalactiae* subsp. J Med Microbiol. 2005;54:1103-5.
2. Shah HN, Collins DM. *Prevotella*, a new genus to include *Bacteroides melaninogenicus* and related species formerly classified in the genus *Bacteroides*. Int J Syst Bacteriol. 1990;40:205-8.
3. Loesche WJ, Socransky SS, Gibbons RJ. *Bacteroides oralis*, proposed new species isolated from the oral cavity of man. J Bacteriol. 1964;88:1329-37.
4. Surbled M, Perrier-Creach C, Rabouille Y, et al. Spondylodiscitis caused by *Bacteroides melaninogenicus*. Presse Med. 1992;21:1870-1.
5. Salavert M, Navarro V, Roig P, et al. Vertebral osteomyelitis by *Prevotella melaninogenica*, *Candida albicans*, and *Mycobacterium tuberculosis* in a intravenous drug addict. Enferm Infecc Microbiol Clin. 1997;15:117-9.
6. Fukuoka M, Aita K, Aoki Y, et al. Pyogenic vertebral osteomyelitis caused by *Prevotella intermedia*. J Infect Chemother. 2002;8:182-4.
7. Schöber W, Horger M, Niehues D. One case of Gram-negative anaerobic spondylodiscitis with *Prevotella intermedia*. Arch Orthop Trauma Surg. 2003;123:436-8.
8. Mukhopadhyay S, Rose F, Frechette V. Vertebral osteomyelitis caused by *Prevotella* (*Bacteroides*) *melaninogenicus*. South Med J. 2005;98:226-8.
9. Salliot C, Lavie F, Azria A, et al. Retroperitoneal fibrosis secondary to spondylodiscitis after infection with *Prevotella*. J Rheumatol. 2005;32:957-8.
10. Purushothaman B, Lakshmanan P, Gatehouse S, Fender D. Spondylodiscitis due to *Prevotella* associated with ovarian mass - a rare case report and review of literature. World Neurosurg. 2010;73:119-22.
11. Huang CR, Lu CH, Chuang YC, et al. Clinical characteristics and therapeutic outcome of Gram-negative bacterial spinal epidural abscess in adults. J Clin Neurosci. 2011;18:213-7.