



The Brazilian Journal of  
**INFECTIOUS DISEASES**

[www.elsevier.com/locate/bjid](http://www.elsevier.com/locate/bjid)



**Clinical image**

**Labial ulcer: oral manifestation of syphilis**



Cristian Vera-Kellet<sup>a</sup>, Isidora Harz-Fresno<sup>b</sup>, Juan Manriquez<sup>a,\*</sup>

<sup>a</sup> Department of Dermatology, Pontificia Universidad Católica de Chile, Santiago, Chile

<sup>b</sup> Hospital Santiago Oriente Dr. Luis Tisné Brousse, Santiago, Chile

ARTICLE INFO

Article history:

Received 25 March 2014

Accepted 28 March 2014

Available online 14 May 2014

A 39-year-old woman presented with a 6-week history of an ulcerated and indurated lesion in the upper lip (Fig. 1). The histological study showed an ulcer with dense lymphoplasmacytic infiltrate and the immunohistochemistry (IHC) was positive for spirochetes. Both, the rapid plasma reagin test (RPR) and the microhemagglutination of the *Treponema pallidum* (MHA-TP) test were reactive with a RPR of 1:32.

Due to physical examination, biopsy, IHC and laboratory tests (MHA-TP and RPR) the patient was diagnosed with

primary syphilis. Infective syphilis is caused by *T. pallidum*, an anaerobic filamentous spirochete.<sup>1</sup> The primary stage is known as syphilis chancre.<sup>2</sup> The mouth is a rare place of presentation of syphilis chancre, with 10–20% of cases intrarectal, perianal or oral lesions.<sup>2</sup> The treatment of primary syphilis



Fig. 1



Fig. 2

\* Corresponding author at: Department of Dermatology, Pontificia Universidad Católica de Chile, Marcoleta 350. Santiago, Chile.

E-mail address: [jjmanriq@uc.cl](mailto:jjmanriq@uc.cl) (J. Manriquez).

<http://dx.doi.org/10.1016/j.bjid.2014.03.006>

1413-8670/© 2014 Elsevier Editora Ltda. All rights reserved.

is intramuscular benzathine penicillin.<sup>3</sup> The patient and his partner were treated with 2,400,000, intramuscular benzathine penicillin with a complete resolution of the lesion (Fig. 2).

---

**Conflicts of interest**

The authors declare no conflicts of interest.

**REFERENCES**

---

1. Leão JC, Gueiros LA, Porter SR. Oral manifestations of syphilis. *Clin São Paulo Braz*. 2006;61:161–6.
2. Ficarra G, Carlos R. Syphilis: the renaissance of an old disease with oral implications. *Head Neck Pathol*. 2009;3:195–206.
3. Dylewski J, Duong M. The rash of secondary syphilis. *CMAJ Can Med Assoc J*. 2007;176:33–5.