

Antidiabetic activity evaluation of *Onobrychis* species on alloxan-induced diabetic mice

Hanefi Ozbek¹, Ozlem Bahadir Acikara², Burcin Ergene Oz², Serkan Ozbilgin², Neriman Ipek Kirmizi^{3*}, Bade Cevriye Ozrenk³, Mehmet Tekin⁴, Gulcin Saltan²

¹Istanbul Medipol University, Faculty of Medicine, Department of Pharmacology, Istanbul, Turkey, ²Ankara University, Faculty of Pharmacy, Department of Pharmacognosy, Ankara, Turkey, ³Istanbul Medipol University, IMU Vocational School, Department of Occupational Health and Safety, Istanbul, Turkey, ⁴Trakya University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Sivas, Turkey

In the current study, four *Onobrychis* species, *O. albiflora* Hub.-Mor., *O. argyrea* Boiss. subsp. *argyrea* Boiss., *O. galegifolia* Boiss., and *O. tournefortii* (Willd.) Desv. were collected from Anatolia to be evaluated for their antidiabetic activities. Methanol water extracts of the aerial parts were used for experiments. An alloxan-induced diabetic mice test model was used. Phytochemical analysis of the tested extracts was investigated using the HPLC method. The highest activity was observed with treatment of *O. albiflora* aerial part extract. Significant decrements were detected in the blood glucose levels as follows: 180.83±47.48 and 252.83±50.47mg/dL at 100 mg/kg and 200 mg/kg doses of *O. albiflora*, respectively, when compared to the isotonic saline solution control group, eliciting a blood glucose level of 494.20±27.32. Among the tested standard compounds, rutin and isoquercetin were detected in the examined species. The highest amount of rutin (1.1981±0.0017%) and isoquercetin (0.7318±0.0197%) were found in *O. albiflora* and *O. argyrea* subsp. *argyrea*, respectively. Antidiabetic activities of the tested *Onobrychis* species seem to indicate a possible correlation with their rutin and isoquercetin contents. Therefore, rutin and isoquercetin may be the antidiabetic compounds that contribute to the antidiabetic activity of the tested *Onobrychis* species.

Keywords: *Onobrychis*. Antidiabetic activity. Alloxan. Rutin. Isoquercetin. Phytochemical analysis.

INTRODUCTION

Onobrychis Miller belongs to the Fabaceae family and is widely distributed in Southwest Asia, Mediterranean countries, and temperate regions of Europe; with around 170 species (Karamian, Asadbegyan 2016). *Onobrychis viciifolia* Scop., known as sainfoin, is used as fodder legume in many countries and has many beneficial effects for ruminants, including providing high protein and energy value, exhibiting antihelmintic activity in cattle and sheep with preventive activities against bloat; which are attributed to the tannin content of the plant (Regos, Urbanella, Treutter, 2009). From a chemical point of view, no sufficient studies have been conducted on the *Onobrychis* species. *Onobrychis viciifolia*, the best known species of this genus, contains

proanthocyanidin polymers (Koupai-Abyazani *et al.*, 1993), cinnamic acid derivatives, flavonoid glycosides (Lu *et al.*, 2000), isoflavonoids, and acylated flavonoids (Veitch *et al.*, 2011). Biological activity studies also revealed that *Onobrychis* species have not been yet investigated properly. Antibacterial activity and moderate antitumor activity by aerial parts of *O. oxydonta* Boiss. have also been reported after the experiments using the potato disc method (Usta, Yildirim, Turker, 2014). *O. armena* Boiss & Huet flower and roots were reported to display antioxidant activity as well as antimicrobial activity against some bacteria, fungus, and fish pathogens, but no cytotoxic activity was reported on kidney fibroblast (BHK 21) cells (Karakoca *et al.*, 2015). The antioxidant activity of *O. hypargyrea* Boiss. in different test models was examined and moderate antioxidant properties were determined in ethyl acetate, methanol, and water extracts in all test systems when compared to trolox and EDTA as positive controls. Methanol and aqueous extracts of *O.*

*Correspondence: N. I. Kirmizi. Istanbul Medipol University, IMU Vocational School, Department of Occupational Health and Safety, Istanbul, Turkey. E-mail: nikirmizi@medipol.edu.tr

hyparygrea also inhibited α -glucosidase with 3.89 ± 0.097 and 5.86 ± 0.050 mg/mL IC_{50} values, respectively (Zengin *et al.*, 2015). *O. viciifolia*, *O. sosnovskyi* Grossh, and *O. melanotricha* Boiss. aerial parts were reported to have strong antioxidant activities in DPPH radical scavenging, metal chelating, and β -carotene bleaching methods (Karamian, Asadbegy, 2016).

In the current study, the aerial parts of four different *Onobrychis* species namely, *O. albiflora* Hub.-Mor., *O. argyrea* Boiss. subsp. *argyrea* Boiss., *O. galegifolia* Boiss., and *O. tournefortii* (Willd.) Desv. were evaluated for their potential antidiabetic activities in alloxan-induced diabetic mice. Furthermore, the phytochemical content of the aerial parts was analyzed using the HPLC method.

MATERIAL AND METHODS

Plant material

Onobrychis species were collected from different regions of Anatolia (Table I). Taxonomic identification of the plants was confirmed by Assistant Professor Mehmet Tekin (Cumhuriyet University, Faculty of Pharmacy, Department of Pharmaceutical Botany). Voucher specimens were kept in the Herbarium of Cumhuriyet University, Faculty of Science (CUFH).

Extraction of plant materials for biological activity tests

Dried and powdered aerial parts of each plant (30 g) were extracted with a 300 mL methanol water solution (80:20) at room temperature during 8 h for 3 days by continuous stirring. Each extract was filtered and concentrated to dryness under reduced pressure and low temperature (40–50°C) on a rotary evaporator to yield crude extracts.

HPLC analysis

HPLC analyses were carried out using an Agilent LC 1200 model chromatograph (Agilent Technologies, California, USA). The diode array detector (DAD) was set at wavelengths of 210 nm, 254 nm and 330 nm, and peak areas were integrated automatically using Agilent computer software. The chromatograms were plotted and processed using the above mentioned software. Separation was carried out using an ACE 5 C18 (250 mm \times 4.6 mm; 5 μ m) column. The mobile phase was composed of 0.2% phosphoric acid in water (A), acetonitrile (B), and in gradient elution: initially 0 min, A–B (90:10, v/v); then 0–10 min, linear change from A–B (90:10, v/v) to A–B (80:20); then 10–20 min, linear change to A–B (80:20); then 20–35 min, linear change to A–B (70:30); then 35–45 min, linear change to A–B (30:70); then 45–50 min, linear change to A–B (0:100), and 50–55 min, isocratic elution was from A–B (0:100). The flow rate was 0.5 mL/min and column temperature was maintained at 20°C. The sample injection volume was 10 μ L.

Extraction procedure, preparation of standard and sample solutions

Aerial parts of the four different *Onobrychis* species were used in this experiment. A methanol water solution (80:20) was used for the extraction of plant samples. For HPLC analysis, 10 mg of methanol water extracts of four species were weighed in 10 mL volumetric flasks, dissolved in methanol water solutions and adjusted to the final volume separately. Each extract was filtered through a 0.45 μ m membrane filter after adjusting to a final volume of 10 mL with the same solvent. Triplicate 10 μ L injections were performed for the plant samples.

Standard stock solutions were prepared as follows: 0.1 mg/mL for caffeic acid, ferulic acid, luteolin-7-

TABLE I - List of *Onobrychis* species collected for the study

Plant Species	Location	Collector and herbarium number
<i>O. albiflora</i>	Sivas: Sincan to Kangal, 3. km roadside 1220 m, 39° 28' 01,9" N; 37° 50' 34,5" E	M. Tekin, 1291
<i>O. argyrea</i> subsp. <i>argyrea</i>	Sivas: Zara - Divrigi road, 55 km from Divrigi 1691 m, 39° 36' 58,1" N; 37° 44' 28,4" E	M. Tekin, 1293
<i>O. galegifolia</i>	Sivas: Divrigi to Sincan, 2-3 km from Sincan, roadside, 1312 m, 39° 30' 21,7" N; 37° 49' 45,4" E	M. Tekin, 1294
<i>O. tournefortii</i>	Sivas: Sivas to Hafik, Emre village vicinity, 1317 m, 39° 49' 37,5" N; 37° 17' 05,1" E	M. Tekin 1290

glucoside, hyperoside, isoquercetin, rutin, orientin, myricetin, and apigenin. All compounds were weighed in a 10 mL volumetric flask, dissolved in a methanol water solution (80:20) and adjusted to the final volume separately. Different concentration levels (between 0.0005 mg/mL - 0.05 mg/mL) were prepared by diluting the stock solution. Triplicate 10 µL injections were performed for each standard solution. The peak area of each solution was plotted against the concentration to obtain the calibration curves.

Validation procedure - Limit of detection and quantification

Limit of detection (LOD) and limit of quantification (LOQ) were established at signal to noise (S/N) ratios of 3 and 10, respectively. LOD and LOQ concentrations were experimentally verified by 9 injections of standard compounds.

Animals

The study protocol [30/09/2015-69] was approved by the Ethics Committee of Istanbul Medipol University. Balb/C strain mice [22-30 g] obtained from Istanbul Medipol University Regenerative and Restorative Medical Center (REMER), were used for this experiment. The animals were housed in standard cages [48 cm × 35 cm × 22 cm] at room temperature [22±2 °C], with artificial light from 7.00 am to 7.00 pm, and provided with pelleted food and water ad libitum.

Chemicals

Alloxan was obtained from Sigma [Steinheim-Germany]. *Onobrychis* extracts were prepared using water [w/v] before being given to the mice.

Antidiabetic activity assay

An alloxan-induced test model was used to evaluate antidiabetic activity (Kumar *et al.*, 2008; Tripathi, Verma, 2014; Ozbek *et al.*, 2017). Mice were kept without food for 18 hours before the alloxan treatment. Alloxan was applied in an isotonic saline solution by i.p. [intraperitoneal] administration [150 mg/kg] three times over a 48-hour period. Mice were kept without food following the last alloxan treatment, and blood glucose levels were measured. Mice with blood glucose levels of 200 mg/dL and higher were subjected to further studies as diabetic animals. Diabetic mice were divided into nine

groups in order to determine the antidiabetic activities of *Onobrychis* aerial part extracts (all extracts were dissolved in an isotonic saline solution). Animals were assigned into following groups (n=6): Group 1, the control group, received isotonic saline solution [ISS] 0.1 mL; Group 2 and Group 3 received *O. tournefortii* extract 100 mg/kg and 200 mg/kg i.p., respectively; Group 4 and Group 5 received *O. argyrea* subsp. *argyrea* extract 100 mg/kg and 200 mg/kg i.p., respectively; Group 6 and Group 7 received *O. galegifolia* extract 100 mg/kg and 200 mg/kg i.p., respectively; and finally Group 8 and Group 9 received *O. albiflora* extract 100 mg/kg and 200 mg/kg i.p, respectively.

Following the administration of the tested extracts, blood samples collected from the tail in the 1st, 2nd and 4th hours were measured for their glucose concentrations using an Accu-Check® sugar strip according to the glucose-oxidase-peroxidase method.

Statistical analyses

Statistical analysis was carried out using SPSS 18.0 software. The results were reported as mean±SEM [standard error of mean]. A one-way analysis of variance [post-hoc Tukey's HSD, Dunnett-t test] was used for statistical analyses. Probability levels of less than 0.05 [P<0.05] were considered as significant.

RESULTS

HPLC analysis

O. albiflora, *O. argyrea* subsp. *argyrea*, *O. galegifolia*, and *O. tournefortii* were analyzed for their phenolic contents using the HPLC method. The HPLC method was developed by comparing the chromatographic profile and data obtained from the standards and samples, considering the following parameters: retention time, analyzing time for samples, separation, peak shapes, and maximum UV absorption of the standards. Figure 1 displays the chromatogram of the aerial part extract of *O. albiflora*, and Figure 2 and Figure 3 show the standard compounds. Peaks were identified by comparison of the retention times and UV absorption spectra with the acquired standards.

For the quantification of the phenolic compounds, standard stock solutions were prepared as described in the experimental section, and diluted to six different concentrations to establish the calibration curves. All calibration curves showed good linearity responses with high correlation coefficients. The contents of the




FIGURE 1 - HPLC chromatogram of *O. albiflora*.




FIGURE 2 - HPLC chromatogram of rutin.




FIGURE 3 - HPLC chromatogram of isoquercetin

extracts are shown in Table II and the results are given as percentages (%) in plant material. Among the tested compounds, rutin was identified in all *Onobrychis* species tested, and the highest concentration was detected in aerial parts of *O. albiflora* ($1.1981 \pm 0.0017\%$). Isoquercetin was also found in all *Onobrychis* species, except *O. galegifolia*.

O. argyrea subsp. *argyrea* contained the highest percentage of isoquercetin ($0.7318 \pm 0.0197\%$), (Table II).

LOD and LOQ concentrations were determined as 0.103 and 0.344 $\mu\text{g/mL}$ for rutin, whereas these were 0.0944 and 0.315 $\mu\text{g/mL}$ for isoquercetin, respectively.

TABLE II - Rutin and isoquercetin content of *Onobrychis* species

Plant Species	Rutin (%)	Isoquercetin (%)
<i>O. albiflora</i>	1.1981±0.0017	0.1288±0.0003
<i>O. argyrea</i> subsp. <i>argyrea</i>	0.3719±0.0012	0.7318±0.0197
<i>O. galegifolia</i>	0.1462±0.0003	-
<i>O. tournefortii</i>	0.2135±0.0004	0.4077±0.0000

TABLE III - Blood sugar levels of alloxan-induced diabetic mice

Groups	Blood sugar levels (mg/dL)			
	Before <i>Onobrychis</i> treatment	After <i>Onobrychis</i> treatment		
		1 h	2 h	4 h
Control (ISS)	470,60±23,62	471,00±13,78	493,80±15,55	494,20±27,32
OT 100 mg/kg	390,50±36,55	322,33±30,25	294,83±32,71*	333,50±43,67
OT 200 mg/kg	374,00±34,22	364,17±26,24	352,17±43,85	268,50±41,61*
OAA 100 mg/kg	378,67±52,78	399,00±27,51	365,67±45,75	286,33±44,58*
OAA 200 mg/kg	352,50±50,06	415,83±40,47	374,00±47,23	332,33±52,78
OG 100 mg/kg	357,17±49,07	358,50±61,88	338,83±55,95	340,83±40,93
OG 200 mg/kg	430,33±16,84	442,50±17,63	427,33±20,82	373,33±34,07
OA100 mg/kg	374,17±27,79	292,83±56,01	235,17±45,85*	180,83±47,48*
OA 200 mg/kg	419,50±20,67	375,16±14,48	349,43±60,90	252,83±50,47*
F/p-value	1,000 / 0,450	1,838 / 0,095	2,613 / 0,020	3,744 / 0,002

OA: *O. albiflora*, OAA: *O. argyrea* subsp. *argyrea*, OG: *O. galegifolia*, OT: *O. tournefortii*

Antidiabetic activity

Aerial part extracts of *O. albiflora*, *O. argyrea* subsp. *argyrea*, *O. galegifolia*, and *O. tournefortii* were tested for their potential antidiabetic activities using alloxan-induced diabetic animals in current study. Blood glucose levels were measured at four different times; before *Onobrychis* extract treatment and after 1, 2 and 4 hours of treatment. All measured levels of blood glucose are presented in Table III. *O. tournefortii* and *O. albiflora* were associated with significantly decreased blood glucose levels at 100 mg/kg dose after 2 hours of treatment when compared to isotonic saline group. *O. tournefortii* 200 mg/kg, *O. argyrea* subsp. *argyrea* 100 mg/kg, and *O. albiflora* 100 mg/kg and 200 mg/kg caused notable decrease in blood glucose level after 4 hours of treatment as shown in Table III. However, *O. albiflora* aerial part extract displayed the highest activity at 100 mg/dose and after 4 hours of treatment.

DISCUSSION

Our study showed that *O. albiflora* treatment resulted in the highest antidiabetic activity in alloxan-

induced diabetic animals at 100 mg/kg dose. The highest level of activity was observed after 4 hours of treatment. However, increasing the dosage to 200 mg/kg did not reduce blood sugar levels in animals even after 4 hours of the *O. albiflora* treatment. *O. tournefortii* and *O. argyrea* subsp. *argyrea* were determined to be effective at 200 mg/kg and 100 mg/kg doses, respectively, after 4 hours of treatment. Both extracts lead to slight reductions in blood glucose levels when compared to *O. albiflora*. *O. galegifolia* did not significantly reduce elevated blood glucose levels, but displayed a tendency to lower the blood glucose at 1, 2 and 4 hours when compared with the control group. These results demonstrated that treatment with *O. albiflora* could significantly improve blood glucose levels in alloxan-induced diabetic animals.

We used the alloxan-induced diabetic model, a well-known method of evaluating especially the type I DM. Alloxan selectively destroys insulin-producing beta cells found in the pancreas. A sudden increase in insulin secretion induced by alloxan treatment, in the presence or absence of glucose, is associated with a complete suppression of beta islet response to glucose, even when high glucose concentrations are used (Rohilla, Ali, 2012;

Tripathi, Verma, 2014). Alloxan inhibits glucose-induced insulin secretion by inhibiting the enzyme glucokinase, a glucose sensor in beta cells, by reacting with two –SH groups in the sugar binding site. Alloxan and dialuric acid, a reduction product of alloxan, induces a redox cycle resulting in the formation of reactive oxygen species (ROS) and superoxide radicals. Ferric ions from ferritin are released by superoxide radicals and transformed to ferrous and ferric ions. Additionally, superoxide radicals undergo dismutation to hydrogen peroxide in the presence of superoxide dismutase, and highly active hydroxyl radicals are formed. The destruction of beta cells occurs rapidly due to ROS with simultaneous excessive increases in cytosolic calcium concentration (Rohilla, Ali, 2012; Tripathi, Verma, 2014). Alloxan-induced toxicity can be prevented by enzymatic antioxidants such as catalase and dismutase, as well as non-enzymatic hydroxyl radical scavengers (Etuk, 2010).

Experimental and clinical evidence has revealed that ROS are elevated in both types of diabetes, and play an important role in the development and progression of DM. In normal healthy cells, the presence of the protective activity of antioxidant enzymes and the non-enzymatic antioxidant levels of ROS can be controlled. However, excessive cellular levels of ROS induced by hyperglycemia in diabetics cause major complications (Matough *et al.*, 2012). Multiple factors induce increased oxidative stress in diabetics. ROS generation can be elevated directly by hyperglycemia. Glucose autooxidation is one of the main factors among these resulting $\cdot\text{OH}$ radicals. Moreover, glucose reacts with proteins which promote the formation of advanced glycation end-products (Johansen *et al.*, 2005; Jebur, Mokhamer, El-Demerdash, 2016). Additionally, increased production of $\cdot\text{O}_2^-$ occurs in hyperglycemic subjects due to the enhanced metabolism of glucose. NOS, NAD(P)H oxidase, and xanthine oxidase are the enzymatic sources of reactive species in diabetics. $\cdot\text{O}_2^-$ production is increased by enzymatic sources and predominantly mediated by NAD(P)H oxidase. Furthermore, a mitochondrial respiratory chain precipitates the hyperglycemic induced generation of $\cdot\text{O}_2^-$ at the mitochondrial level in diabetics (Johansen *et al.*, 2005). The long-term complications in diabetic patients include damage, dysfunction, and failure of different organs; resulting in retinopathy, nephropathy, neuropathy, atherosclerosis, coronary artery diseases, cerebrovascular, and peripheral vascular diseases (Matough *et al.*, 2012). Oxidative stress caused by hyperglycemia induces free radicals to play an important role in the development and progression of diabetes mellitus (Johansen *et al.*, 2005; Matough *et al.*, 2012). Treatment with antioxidants

could be an effective strategy for improving diabetic complications (Johansen *et al.*, 2005). Naturally occurring antioxidants such as vitamin C, E and α -lipoic acid treatments were reported to reduce the oxidative stress experimentally in diabetic animals and clinically in human (Johansen *et al.*, 2005; Jebur *et al.*, 2016). For this reason, recently there has been an increasing interest in natural antioxidants derived from plant materials. Natural antioxidants occur in all parts of plants as phytochemicals such as monoterpenes, diterpenes, triterpenes, and phenolic compounds, flavonoids, coumarins, lignans, phenylpropanoids, and tannins (Jebur, Mokhamer, El-Demerdash, 2016). It has been reported that species of *Onobrychis* also have important antioxidant activities. *O. armena* Boiss & Huet flowers and roots have antioxidant capacities as shown by the phosphomolybdenum method, cupric ion reducing antioxidant capacity, DPPH radical scavenging activity, and ferric ion reducing capacity (Karakoca *et al.*, 2015). Total antioxidant capacity, scavenging activities against DPPH and ABTS radicals, metal chelating activity of ferrous ions, β -carotene/linoleic acid bleaching activity, ferric reducing antioxidant power and cupric ion reducing antioxidant capacity were measured to determine the antioxidant activity of *O. hypargyrea* Boiss. Moderate antioxidant capacities were observed in ethyl acetate, methanol, and water extracts in all test systems when compared to trolox and EDTA as positive controls (Zengin *et al.*, 2015). *O. viciifolia*, *O. sosnovskyi* Grossh and *O. melanotricha* Boiss. aerial parts were also reported to have strong antioxidant activities in DPPH radical scavenging, metal chelating, and β -carotene bleaching methods (Karamian and Asadbegy 2016). The total antioxidant capacity of *O. altissima* Grossh was determined to be an ABTS radical scavenger, and the plant's ability to reduce glutathione levels was identified (Erbil *et al.*, 2015). The current study did not report on the biological activities of *O. albiflora*, *O. argyrea* subsp. *argyrea*, *O. galegifolia*, and *O. tournefortii* in terms of their antidiabetic activities. However, previous studies of the *Onobrychis* genus could be used to suggest that potential antioxidant activities may contribute to their antidiabetic effects.

In order to clarify the active compound(s) of the tested *Onobrychis* species, HPLC analysis was performed by using some standard compounds. Among the tested phenolic compounds, rutin and isoquercetin were determined. The presence of rutin was detected in all tested species while isoquercetin was detected in *O. albiflora*, *O. argyrea* subsp. *argyrea* and *O. tournefortii*. The highest content of rutin and isoquercetin was found in *O. albiflora* and *O. argyrea* subsp. *argyrea*, respectively.

O. albiflora displayed the highest antidiabetic activity while *O. galegifolia* exhibited non-significant activity due to having the lowest rutin content. Antidiabetic activities of the tested *Onobrychis* species seem to be correlated with rutin content. In fact, antidiabetic activities of rutin and isoquercetin were reported previously. Rutin treatment was associated with decreased levels of plasma glucose, glycosylated hemoglobin, thiobarbituric acid reactive substances, and lipid hydroperoxides, and elevated levels of insulin, C-peptide, and non-enzymatic antioxidants such as glutathione, vitamin C, E, and ceruloplasmin in streptozotocin-induced diabetic rats. On the other hand, in normal rats, no parameters were altered by rutin treatment. It was suggested that rutin acted as a free radical scavenger and inhibited lipid peroxidation resulting in the prevention of oxidative stress and the protection of beta cells (Kamalakkannan, Stanley, Prince, 2006). Rutin significantly lowered plasma glucose levels and increased insulin levels. Besides, the restoration of glycogen content and activities of carbohydrate metabolic enzymes was observed in diabetic mice treated with a 100 mg/kg dose. Glucose uptake by rutin treatment has been shown in rat soleus muscle through PI3K, a typical protein kinase C and mitogen-activated protein kinase pathways (Vinayagam, Xu, 2015). Jadhav and Puchchakayala reported that rutin reduced blood glucose levels after 2h and 4h in streptozotocin/nicotinamide-induced diabetic rats. The mechanism of rutin was thought to be increased peripheral utilization of glucose and inhibition of glucose transport (Jadhav, Puchchakayala, 2012). It was also reported that in diabetic rats induced by high fat diet + streptozotocin, rutin treatment (50 mg/kg and 100 mg/kg) for three weeks improved the metabolic status of rats. Plasma glucose levels and glycosylated hemoglobin, as well as proinflammatory cytokines (IL-6 and TNF-alpha) were reduced. Liver antioxidant status and serum lipid profile were also improved (Niture, Ansari, Naik, 2014). Furthermore, protective and ameliorative effects of rutin in myocardial dysfunction, oxidative stress, apoptosis, and inflammation were observed in the hearts of diabetic rats (Vinayagam, Xu, 2015). Isoquercetin was also identified as the compound responsible for the antidiabetic activity of *Vaccinium vitis idaea*. Isoquercetin, as well as its aglycone quercetin, stimulated the AMPK pathway at 25-100 mM concentrations. Both isoquercetin and quercetin compounds enhanced glucose uptake by 38-59% and 37%, respectively. However, inhibition of ATP synthase in isolated mitochondria was achieved only by quercetin at 25 and 100 mM (34% and 79%), respectively (Eid *et al.*, 2010). Furthermore, hepatic glucose-6-phosphatase was inhibited by isoquercetin *in vitro* (Hung *et al.*, 2012).

The short period of time used for the study procedure could be regarded as a limitation of current study. Therefore, further studies should be performed to observe for the potential long-term antidiabetic effects of the *Onobrychis* species.

In conclusion, *O. albiflora* that contains rutin in relatively high amounts was determined to be most effective species for alloxan-induced diabetic mice test models. This was followed by *O. tournefortii* at 200 mg/kg dose, and *O. argyrea* subsp *argyrea* at 100 mg/kg dose. *O. galegifolia*, which exhibited the least activity, contained no isoquercetin and contained rutin in very low levels when compared to other species. It could be suggested that rutin could be the main compound responsible for antidiabetic activities of the tested *Onobrychis* species. Additionally, isoquercetin, which has been proven to be an antidiabetic compound in earlier studies, could contribute to the *Onobrychis*' antidiabetic activities. Nevertheless, further studies are warranted to reveal responsible compounds and to identify their mechanisms.

REFERENCES

- Eid HM, Martineau LC, Saleem A, Muhammad A. Stimulation of AMP-activated protein kinase and enhancement of basal glucose uptake in muscle cells by quercetin and quercetin glycosides, active principles of the antidiabetic medicinal plant *Vaccinium vitis-idaea*. *Mol Nutr Res.* 2010;54(7):991-1003.
- Erbil N, Duzguner V, Durmuskahya C, Alan Y. Antimicrobial and Antioxidant Effects of Some Turkish Fodder Plants Belongs to Fabaceae Family (*Vicia villosa*, *Trifolium ochroleucum* and *Onobrychis altissima*). *Orient J Chem.* 2015;31(3):53-8.
- Etuk EU. Animals models for studying diabetes mellitus. *Agric Biol J N Am.* 2010;1(2):130-4.
- Hung H, Qian K, Morris-Natschke SL, Hsu C, Lee K. Recent discovery of plant-derived anti-diabetic natural products. *Nat Prod Rep.* 2012;29(5):580-606.
- Jadhav R, Puchchakayala G. Hypoglycemic and antidiabetic activity of flavonoids: boswellic acid, ellagic acid, quercetin, rutin on streptozotocin-nicotinamide induced type 2 diabetic rats. *Int J Pharm Pharm Sci.* 2012;4(2):251-7.
- Jebur AB, Mokhamer MH, El-Demerdash FM. A Review on oxidative stress and role of antioxidants in diabetes mellitus. *Austin Endocrinol Diabetes Case Rep.* 2016;1(1):id1006.

- Johansen JS, Harris AK, Rychly DJ, Ergul A. Oxidative stress and the use of antioxidants in diabetes: Linking basic science to clinical practice. *Cardiovasc Diabetol*. 2005;4(5):1-11.
- Kamalakkannan N, Stanely P, Prince M. Antihyperglycaemic and antioxidant effect of rutin, a polyphenolic flavonoid, in streptozotocin-induced diabetic Wistar rats. *Basic Clin Pharmacol Toxicol*. 2006;98(1):97-103.
- Karakoca K, Asan-Ozusaglam M, Cakmak YS, Teksen M. Phenolic compounds, biological and antioxidant activities of *Onobrychis armena* Boiss. & huet flower and root extracts. *Chiang Mai J Sci*. 2015;42(2):376-92.
- Karamian R, Asadbegy M. Antioxidant activity, total phenolic and flavonoid contents of three onobrychis species from Iran. *Pharm Sci*. 2016;22(2):112-9.
- Koupai-Abyazani MR, Muir AD, Bohm BA, Towers GHN, Gruber MY. The proanthocyanidin polymers in some species of *Onobrychis*. *Phytochemistry*. 1993;34(1):113-7.
- Kumar S, Kumar D, Deshmukh RR, Lokhande PD, More SN, Rangari VD. Antidiabetic potential of *Phyllanthus reticulatus* in alloxan-induced diabetic mice. *Fitoterapia*. 2008;79(1):21-3.
- Lu Y, Sun Y, Foo LY, McNabb WC, Molan AL. Phenolic glycosides of forage legume *Onobrychis viciifolia*. *Phytochemistry*. 2000;55(1):67-75.
- Matough FA, Budin SB, Hamid ZA, Alwahaibi N, Mohamed J. The role of oxidative stress and antioxidants in diabetic complications. *SQU Med J*. 2012;12(1):5-18.
- Niture NT, Ansari AA, Naik SR. Anti-hyperglycemic activity of Rutin in streptozotocin-induced diabetic rats: An effect mediated through cytokines, antioxidants and lipid biomarkers. *Indian J Exp Biol*. 2014;52(7):720-7.
- Ozbek H, Bahadir O, Keskin I, Kirmizi NI, Ozbilgin S, Ergene Oz B, et al. Evaluation of hepatoprotective and antidiabetic activity of *Alchemilla mollis*. *Biomed Pharmacother*. 2017;86:172-6.
- Regos I, Urbanella A, Treutter D. Identification and quantification of phenolic compounds from the forage legume sainfoin (*Onobrychis viciifolia*). *J Agric Food Chem*. 2009;57(13):5843-52.
- Rohilla A, Ali S. Alloxan induced diabetes: mechanisms and effects. *Int J Research Pharm Biomed Sci*. 2012;3(2):819-23.
- Tripathi V, Verma J. Different models used to induced diabetes: a comprehensive review. *Int J Pharm Pharm Sci*. 2014;6(6):29-32.
- Usta C, Yildirim AB, Turker AU. Antibacterial and antitumour activities of some plants grown in Turkey. *Biotechnol Biotechnol Equip* 2014;28(2):306-15.
- Veitch NC, Regos I, Kite GC, Treutter D. Phytochemistry Acylated flavonol glycosides from the forage legume, *Onobrychis viciifolia* (sainfoin). *Phytochemistry*. 2011;72(4-5):423-9.
- Vinayagam R, Xu B. Antidiabetic properties of dietary flavonoids: a cellular mechanism review. *Nutr Metab*. 2015;12(60):1-20.
- Zengin G, Guler GO, Aktumsek A, Ceylan R, Marie C, Picot N, et al. Enzyme inhibitory properties, antioxidant activities, and phytochemical profile of three medicinal plants from Turkey. *Adv Pharmacol Sci*. 2015;2015:410675.

Received for publication on 15th March 2018

Accepted for publication on 27th April 2018