

Weak solutions for the electrophoretic motion of charged particles

LUCIANO BEDIN¹ and MARK THOMPSON²

¹Department of Mathematics, UFSC, Trindade, 88040-900 Florianópolis, SC.

²Department of Pure and Applied Mathematics, IM, UFRGS
Av. Bento Gonçalves 9500, 91509-900 Porto Alegre, RS.

E-mails: luciano@mtm.ufsc.br / thompson@mat.ufrgs.br

Abstract. We introduce a weak formulation for a system of electrostatic and hydrodynamic equations modelling the electrophoretic motion of charged particles in ionized fluids. We obtain a local in time existence theorem, using the results established in [11] and properties of the solutions of the Poisson-Boltzmann equation. These properties follows from singular integral operators techniques.

Mathematical subject classification: 35J60, 92C05, 35J50, 42B20.

Key words: Poisson-Boltzmann equation, singular integral operators, Stokes problem, electrophoresis.

1 Introduction

In this paper we establish results on existence and uniqueness of the weak solutions for a system modelling the motion of a charged particle driven by the action of an external electrical field. This phenomena is known as electrophoresis and is important in many technical applications (a vast literature is available: see for example [29], [24], [27], [2], [1], [3], [13], [30], [14]).

We are considering a particle (a charged polymer, for example) immersed in an ionized solution (a viscous incompressible fluid). On the boundary of the enclosure an external electric field induces the electrical potential inside the enclosure which is determined by the Poisson-Boltzmann equation (see [19],

[10]). The hydrodynamic behavior of the system is governed by the Navier-Stokes equation.

We are not considering the usual approximation for the effect of the electrical field on the particle based on Prandtl boundary layer theory well known in colloid science, the so called slip-velocity condition (for details see [4], [22], [27]); their derivation requires better regularity properties of the boundary of the particle (see [28] and the discussion in the introduction of [27]).

As remarked in [24] the theoretical analysis of the electrophoretic motions is quite difficult, as it combines specific features of polymer physics with the intrinsic complexity of electrokinetic phenomena. From a mathematical standpoint the difficulties reside in the treatment of the electrical-hydrodynamic couple and on the low regularity of the data. We have established elsewhere [6] the existence of a H^1 -variational solution for the electrostatic potential, for a general class of domains. In the case of Lipschitz regions we have established a $H^{3/2}$ -regularity result by means of the singular integral operators theory; this regularity is optimal, even in C^1 -domains (see the comments and negative results in [20]). For $C^{1,\alpha}$ -domains, $0 < \alpha \leq 1$, and suitable charge distribution this theory can be applied in the classical sense ([25], [18], [7]) in order to obtain more regularity for the potential [5].

Recently, the motion of rigid bodies in a bounded domain filled with a viscous flow has been treated rigorously ([11], [17]). Special techniques (from the transport theory [23]) have been used in order to obtain existence and properties of the suitable weak solutions for these systems. In particular in [11] a global weak formulation is introduced and existence of solutions local in time is established when a L^2 body force and $C^{1,1}$ -domains are considered. Evidently, in the study of the electrophoretic motion we can not use directly these results because we have an external electrical field interacting with the ionized solution. However, we obtain a similar result of local existence (see Theorem 3.1); this is obtained as a consequence of uniform bounds and convergence properties involving the electrical force term \mathbf{F} (Corollary 4.1, Theorem 4.2). Following the discussion in [6] we choose to prove these properties restricted to the case which the surface charge distribution of the particle and the fixed charge distribution (in the particle and in the solution) are L^2 and L^∞ functions, respectively; in this case we need only consider the Lipschitz regularity on the boundary of the particle. The

existence of weak solutions obtained in Theorem 3.1 follows from the additional hypothesis that the particle domain is of class $C^{1,1}$. More properties on \mathbf{F} can be obtained in the $C^{1,\alpha}$ context if we assume stronger hypotheses [5] on the charge distributions but we do not consider this situation in this paper.

2 The governing equations

Consider a rigid, charged particle immersed in a electrolyte solution under the action of an external electrical field. We suppose that the solution (a viscous fluid) occupies a region $D \subset \mathbb{R}^3$ and at the initial moment of time, the particle (a rigid body) occupies a compact region $\overline{K}_0 \subset D$ such that its center of mass is located at the origin $\mathbf{y} = \mathbf{0}$ of a Cartesian coordinates \mathbf{y} .

Let us define $K(t)$ as the domain occupied by the particle in the time t and $\phi(\mathbf{x}, t)$ as a real valued function which represents the electrical potential in $(\mathbf{x}, t) \in D \times [0, T]$, where $\forall t \in [0, T]$, $\text{dist}(\partial D; \partial K(t)) > d$, where d is a fixed constant. We set $\psi(\mathbf{x}, t) = \frac{\phi(\mathbf{x}, t)e}{\Lambda}$, where Λ is the temperature (constant) of the system and e is the electron charge. The governing equations and boundary conditions to ψ are (see [6], [19])

$$\begin{aligned} \nabla \cdot (\mathbf{k}(\mathbf{x}, t) \nabla \psi(\mathbf{x}, t)) - b(\mathbf{x}, \psi(\mathbf{x}, t)) &= \rho(\mathbf{x}, t), \quad \mathbf{x} \in D, \\ \psi_2(\mathbf{x}, t) &= \psi_1(\mathbf{x}, t), \quad \mathbf{x} \in \partial K(t), \\ \psi_2(\mathbf{x}, t) &= \Psi(\mathbf{x}), \quad \mathbf{x} \in \partial D, \\ k_1 \frac{\partial \psi_1}{\partial n}(\mathbf{x}, t) - k_2 \frac{\partial \psi_2}{\partial n}(\mathbf{x}, t) &= \frac{4\pi e}{\Lambda} \sigma(\mathbf{x}, t), \quad \mathbf{x} \in \partial K(t). \end{aligned} \tag{2.1}$$

Here

- $\mathbf{k} : D \times [0, T] \rightarrow \mathbf{L}(\mathbb{R}^3, \mathbb{R}^3)$ is defined as $k_{ij}(\mathbf{x}, t) = \delta_{ij}k_1$ if $\mathbf{x} \in K(t)$, $k_{ij}(\mathbf{x}, t) = \delta_{ij}k_2$ if $\mathbf{x} \in D \setminus \overline{K(t)}$, where k_1, k_2 are the dielectric constants in $K(t)$ and $D \setminus \overline{K(t)}$ respectively.
- $b : D \times \mathbb{R} \rightarrow \mathbb{R}$, $b(\mathbf{x}, \psi(\mathbf{x}, t)) = k_2 r_D^{-2} \sinh \psi(\mathbf{x}, t)$ if $\mathbf{x} \in D \setminus \overline{K(t)}$, $b(\mathbf{x}, \psi(\mathbf{x}, t)) = 0$ if $\mathbf{x} \in K(t)$, r_D^{-2} is the Debye radius [10].
- $\Psi(x) = \frac{e\Phi(x)}{\Lambda}$, $\psi_1(\mathbf{x}, t) = \psi(\mathbf{x}, t)|_{K(t)}$ and $\psi_2(\mathbf{x}, t) = \psi(\mathbf{x}, t)|_{D \setminus \overline{K(t)}}$.

- σ is a superficial charge distribution and $\rho(\mathbf{x}, t) = (\rho_1(\mathbf{x}, t), \rho_2(\mathbf{x}, t))$, where

$$\begin{aligned}\rho_1(\mathbf{x}, t) &= -\frac{4\pi e}{\Lambda} \rho_1^0(\mathbf{x}), & \rho_2(\mathbf{x}, t) &= -\frac{4\pi e}{\Lambda} \rho_2^0(\mathbf{x}, t), \\ \rho_1^0 &= \rho^0|_{K(t)}, & \rho_2^0 &= \rho^0|_{D \setminus \overline{K(t)}}\end{aligned}$$

are the charge distribution in $K(t)$ and $D \setminus \overline{K(t)}$ respectively.

The action of the electrical field on the particle produce its motion. The motion of fluid is described by the velocity field $\mathbf{v}^f(\mathbf{x}, t)$ (velocity of the fluid material point which has Cartesian coordinates \mathbf{x} at time t) and satisfies the Navier-Stokes equation

$$\begin{aligned}\bar{\nu}_f (\partial_t \mathbf{v}^f + \operatorname{div}(\mathbf{v}^f \otimes \mathbf{v}^f)) - \eta \Delta \mathbf{v}^f + \nabla p &= \bar{\nu}_f \mathbf{F}, & \text{in } \mathcal{D}'(\Omega_T)^3 \\ \operatorname{div} \mathbf{v}^f &= 0, & \text{in } \Omega_T \\ \mathbf{v}^f &= 0, & \text{in } \partial D \\ \mathbf{v}^f|_{t=0} &= \mathbf{v}_0^f & \text{in } D \setminus \overline{K_0}\end{aligned}\tag{2.2}$$

for all $t \in (0, T)$. Here $\eta > 0$ is the viscosity of the fluid, $\bar{\nu}_f > 0$ is the homogeneous fluid density (of the mass) and $\Omega_T = \{(t, \mathbf{x})/t \in (0, T), \mathbf{x} \in D \setminus \overline{K(t)}\}$; denoting ρ^{ions} as the ion density of the solution, we have $\mathbf{F} = -(\rho_2^0 + \rho^{ions})\nabla\phi_2$ as the electrical force on the fluid domain (see [30]). Then $\mathbf{F} = \frac{\Lambda}{4\pi e} (\rho_2 + r_D^{-2} k_2 \sinh(\psi_2)) (\nabla\psi_2) I_{D \setminus \overline{K(t)}}$, using the Boltzmann distribution for ρ^{ions} (see [16]).

Let us set $\mathbf{x}_c(t)$ as the center of mass of the particle; $\mathbf{w}(t)$ the rotation vector; $\mathbf{R}(t)$ the translational velocity; A the symmetric inertial matrix; \mathbf{v}^p the velocity of the particle. We observe that if $\bar{\nu}_p > 0$ is the density (of the mass) of the particle,

$$\mathbf{y}^T A \mathbf{y} = \bar{\nu}_p \int_{K_0} |\mathbf{y} \times (\mathbf{x} - \mathbf{x}_c(0))|^2 d\mathbf{x},$$

for all $\mathbf{y} \in \mathbb{R}^3$. We have also $\mathbf{v}^p(\mathbf{x}, t) = \mathbf{R}(t) + \mathbf{w}(t) \times (\mathbf{x} - \mathbf{x}_c(t))$ for $\mathbf{x} \in K(t)$. It is important to observe the implicit dependence of \mathbf{F} on \mathbf{v}^p .

From the Newtonian mechanics for rigid bodies and the stress tensor in fluid dynamics, if M is the mass of the particle, the evolution law for the motion is given by

$$M \frac{d\mathbf{R}(t)}{dt} = \int_{\partial K(t)} \sigma^H(\mathbf{x}, t) \cdot \mathbf{n}(\mathbf{x}, t) ds + \int_{\partial K(t)} \sigma^E(\mathbf{x}, t) \cdot \mathbf{n}(\mathbf{x}, t) ds$$

and

$$A \frac{d\mathbf{w}(t)}{dt} = \int_{\partial K(t)} (\mathbf{x} - \mathbf{x}_c(t)) \times (\sigma^H(\mathbf{x}, t) \cdot \mathbf{n}(\mathbf{x}, t)) ds + \mathbf{w}(t) \times (A \cdot \mathbf{w})(t) \\ + \int_{\partial K(t)} (\mathbf{x} - \mathbf{x}_c(t)) \times (\sigma^E(\mathbf{x}, t) \cdot \mathbf{n}(\mathbf{x}, t)) ds.$$

If we set $D(\mathbf{v}^f) = \frac{1}{2}(\nabla \mathbf{v}^f + (\nabla \mathbf{v}^f)^T)$ then $\sigma^H(\mathbf{x}, t) = 2\eta D(\mathbf{v}^f(\mathbf{x}, t)) - p(\mathbf{x}, t)\mathbf{I}$ is the stress tensor of the fluid;

$$\sigma^E(\mathbf{x}, t) = \sigma_{ij}^E(\mathbf{x}, t) = \frac{\Lambda k_2}{4\pi e} \left(\frac{\partial \psi_2}{\partial x_i} \frac{\partial \psi_2}{\partial x_j} - \frac{1}{2} \delta_{ij} (\nabla \psi_2)^2 \right)$$

is the electrostatic tensor (see [30]).

We assume the following hypotheses on the data

- (i) K_0 is a Lipschitz domain.
- (ii) D is a C^2 -domain, $\Psi \in H^1(\partial D) \cap C(\partial D)$, $\sigma(\cdot, t) \in L^2(\partial K(t))$, $\rho(\cdot, t) \in L^\infty(D)$, $\forall t \in [0, T]$.

Remark 2.1. As remarked in the introduction of this paper, under the above hypotheses we have established elsewhere [6] that

$$\psi(\cdot, t) = (\psi_1, \psi_2)(\cdot, t) \in H^1(D) \cap (H^{3/2}(K(t)), H^{3/2}(D \setminus \overline{K(t)})),$$

for all $t \in [0, T]$. This regularity result and Trudinger's inequality (see discussion in [6]) give us that

$$\|\sinh(\psi_2)(\cdot, t)\|_{0,p,D \setminus \overline{K(t)}} < +\infty, \quad \forall 1 \leq p < +\infty.$$

Recalling the Sobolev embedding $H^{1/2}(D) \subset L^3(D)$, the Hölder's inequality gives us

$$\int_{D \setminus \overline{K(t)}} \sinh^2(\psi_2(\mathbf{x}, t)) |\nabla \psi_2(\mathbf{x}, t)|^2 d\mathbf{x} \\ \leq \|\sinh(\psi_2(\cdot, t))\|_{0,6,D \setminus \overline{K(t)}}^2 \|\psi_2(\cdot, t)\|_{0,3,D \setminus \overline{K(t)}}^2 < +\infty.$$

Then $\mathbf{F}(\cdot, t) \in L^2(D)^3$, $\forall t \in [0, T]$ (evidently this implies $\mathbf{F} \in L^2((0, T) \times D)^3$). A similar calculation shows us that $\sigma_{ij}^E(\cdot, t) \in L^{3/2}(D \setminus \overline{K(t)})$, $\forall t \in [0, T]$.

As can be seen in the following section an existence result local in time of the suitable weak solutions for (2.2) coupled with (2.1) is available if we assume that K_0 is a $C^{1,1}$ -domain.

3 The notion of weak solution

Following [11], let us define the Eulerian densities $v_p(\mathbf{x}, t) = \bar{v}_p I_{K(t)}(\mathbf{x})$, $v_f(\mathbf{x}, t) = \bar{v}_f I_{D \setminus \overline{K(t)}}(\mathbf{x})$ and the global density $v = v_p + v_f$. We also define the global velocity in D as

$$\mathbf{u}(\mathbf{x}, t) = \begin{cases} \mathbf{v}^f(\mathbf{x}, t) & \text{if } \mathbf{x} \in D \setminus \overline{K(t)}, \\ \mathbf{v}^p(\mathbf{x}, t) & \text{if } \mathbf{x} \in K(t). \end{cases}$$

In view of the conservation of mass, v satisfies the linear transport equation in D

$$\partial_t v + \operatorname{div}(v\mathbf{u}) = 0.$$

We require that $\mathbf{v}^p \cdot \mathbf{n} = \mathbf{v}^f \cdot \mathbf{n}$ and $\sigma^H \cdot \mathbf{n} = \mathbf{T}$ in $\partial K(t)$, where $-\mathbf{T}$ is the force applied by the particle on the fluid. We can write $\mathbf{T} = \Sigma \cdot \mathbf{n}$, where Σ is the Cauchy stress tensor in the body.

On the walls, we enforce homogeneous Dirichlet boundary conditions $\mathbf{u}|_{\partial D} = 0$. Moreover, the incompressibility of the fluid, the rigidity of the structure and $\mathbf{v}^p \cdot \mathbf{n} = \mathbf{v}^f \cdot \mathbf{n}$ imply that $\operatorname{div} \mathbf{u} = 0$.

The evolution laws of the momentum for the fluid and for the particle are given by

$$\begin{aligned} \partial_t(v_f \mathbf{u}) + \operatorname{div}(v_f \mathbf{u} \otimes \mathbf{u}) &= \frac{1}{\bar{v}_f} \operatorname{div}(v_f(2\eta D(\mathbf{u}) - pI)) + \frac{1}{\bar{v}_p} \Sigma \cdot \nabla v_p + \bar{v}_f \mathbf{F} \\ \partial_t(v_p \mathbf{u}) + \operatorname{div}(v_p \mathbf{u} \otimes \mathbf{u}) &= \frac{1}{\bar{v}_p} \operatorname{div}(v_p \Sigma) - \frac{1}{\bar{v}_p} \sigma^H \cdot \nabla v_p - \frac{1}{\bar{v}_p} \sigma^E \cdot \nabla v_p, \end{aligned}$$

respectively. Here $D(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T)$ is the global rate-of-deformation tensor.

Introducing the global stress tensor

$$\mathcal{T} = \frac{v_f \sigma^H}{\bar{v}_f} + \frac{v_p \Sigma}{\bar{v}_p},$$

we obtain the global system in $\mathcal{D}'((0, T) \times D)^3$,

$$\begin{aligned} \partial_t(v\mathbf{u}) + \operatorname{div}(v\mathbf{u} \otimes \mathbf{u}) &= \operatorname{div} \mathcal{T} + v\mathbf{F}, \\ \operatorname{div} \mathbf{u} = 0, \quad \partial_t v + \operatorname{div}(v\mathbf{u}) &= 0, \quad v_p D(\mathbf{u}) = 0, \end{aligned} \tag{3.1}$$

$$\mathbf{u}(\mathbf{x}, 0) = \begin{cases} \mathbf{v}_0^f(\mathbf{x}), & \mathbf{x} \in D \setminus \overline{K_0} \\ \mathbf{v}^p(\mathbf{x}, 0) = \mathbf{w}(0) \times (\mathbf{x} - \mathbf{x}_c(0)) + \mathbf{R}(0), & \mathbf{x} \in K_0, \end{cases} \quad (3.2)$$

$$\begin{aligned} v_p(\mathbf{x}, 0) &= \bar{v}_p I_{K_0}(\mathbf{x}), \quad v_f(\mathbf{x}, 0) = \bar{v}_f I_{D \setminus \overline{K_0}}(\mathbf{x}) \\ v(\mathbf{x}, 0) &= v_0(\mathbf{x}) = v_p(\mathbf{x}, 0) + v_f(\mathbf{x}, 0). \end{aligned} \quad (3.3)$$

We reproduce here, following the paper [11], the notion of the weak solution of the above system

Definition 3.1. (v, \mathbf{u}) is a weak solution of (3.1)–(3.3) in $(0, T)$ if it satisfies the a priori energy bounds

$$v \in L^\infty((0, T) \times D), \quad \mathbf{u} \in L^\infty(0, T; L^2(D))^3 \cap L^2(0, T; H_0^1(D))^3,$$

and if for all $\phi \in \mathcal{V}$ and for almost every $t \in (0, T)$,

$$\begin{aligned} & \int_0^t \int_D (v \mathbf{u} \cdot \partial_t \phi + v \mathbf{u} \otimes \mathbf{u} : D(\phi) - \eta D(\mathbf{u}) : D(\phi) + v \mathbf{F} \cdot \phi) \, d\mathbf{x} d\tau \\ & + \int_D v_0 \mathbf{u}_0 \cdot \phi(0) \, d\mathbf{x} = \left(\int_D v \mathbf{u} \cdot \phi \, d\mathbf{x} \right) (t), \\ & \partial_t v + \operatorname{div}(v \mathbf{u}) = 0, \quad \operatorname{div} \mathbf{u} = 0, \\ & v_p D(\mathbf{u}) = 0, \quad \mathbf{u}|_{\partial D} = 0, \quad \text{in } \mathcal{D}'((0, T) \times D)^3, \\ & v_0 \in L^\infty(D), \quad \mathbf{u}(\cdot, 0) \in L^2(D)^3, \end{aligned} \quad (3.4)$$

where \mathcal{V} is defined by

$$\mathcal{V} = \{ \varphi \in H^1((0, T) \times D)^3 / \varphi(t) \in V(t), \quad \forall t \in (0, T) \},$$

and

$$V(t) = \{ \varphi \in H_0^1(D)^3 / \operatorname{div} \varphi = 0, \quad v_p D(\varphi) = 0 \}.$$

The following existence theorem for the above weak solutions is available [11]

Theorem 3.1. *Under the hypothesis (ii)–(v) (see Section 4) and the additional assumptions that K_0 is a $C^{1,1}$ -domain, $\mathbf{u}_0 \in H_0^1(D)^3$, $\operatorname{div} \mathbf{u}_0 = 0$, $v_p D(\mathbf{u}_0) = 0$ and $\delta(0) > d$, there exist $T^* \in (0, +\infty]$ and a solution (v, \mathbf{u}) of (3.4) such that*

- (i) $\beta(v) \in C([0, T]; L^p(\Omega)) \cap L^\infty((0, \infty) \times D)$ for all $T < T^*$, $p < \infty$ and $\beta \in C^1(\mathbb{R}; \mathbb{R})$.

(ii) $\mathbf{u} \in L^\infty(0, T; H_0^1(D))^3$ and $\partial_t \mathbf{u} \in L^2((0, T) \times D)^3$ for all $T < T^*$.

In [11] the hypothesis $\mathbf{F} \in L^2((0, T) \times D)^3$ (a body force) is assumed (see Remark 2.1) and the proof of the existence theorem as Theorem 3.1 is based on the solution of an approximated system (obtained by regularization techniques). The existence of the approximated solutions is obtained by the Schauder fixed-point theorem (see [5]), via a solution of an appropriate inhomogeneous (linear) Stokes equation. Using the $C^{1,1}$ -regularity of the domains and the smoothness of the coefficients, this linear problem has a solution with the necessary regularity. The solution (\mathbf{u}, ν) is built as a limit of these approximations; the existence of this limit is derived from the compactness properties of the linear transport equation [12]. This is made possible if we can obtain elliptic estimates and a priori bounds for \mathbf{u} as well as energy bounds for ν (see Section 4 in [11]). However as \mathbf{F} depends on \mathbf{u} we need to take some care in this regard. More precisely let us define for each $m \in \mathbb{N}$, $(\mathbf{u}^{(m)}, \nu^{(m)}, \mathbf{F}^{(m)})$ such that $\nu^{(m)}$ is bounded in $L^\infty((0, T) \times D)$ uniformly in m , $\mathbf{u}^{(m)}$ is bounded in $L^2(0, T; H_0^1(D) \cap W^{1,4}(D))^3$, $\partial_t \mathbf{u}^{(m)}$ is bounded in $L^2((0, T) \times D)^3$ uniformly in m , $\nu_0^{(m)}$ converges to ν_0 in $L^2(D)$, $\mathbf{u}_0^{(m)}$ converges to \mathbf{u}_0 in $L^2(D)^3$ and (3.4) is valid, for all $\varphi^{(m)} \in \mathcal{V}^{(m)}$. We set $K^{(m)}(t) = M^{(m)}(t)K_0$, where $M^{(m)}(t)$ is an invertible affine transformation, and we suppose that $\delta = \inf\{\delta^{(m)}(t), t \in [0, T], m \geq 0\} > d$, where $\delta^{(m)}(t) = \text{dist}(\partial D, \partial K^{(m)}(t))$; $\mathbf{F}^{(m)}$ is defined by the calculation of $\psi^{(m)}$, the solution of (2.1) considering $(K^{(m)}(t), \rho^{(m)}, \sigma^{(m)})$. We have to show that $\int_0^T \|\mathbf{F}^{(m)}(\cdot, \tau)\|_{0,2,D}^2 d\tau \leq C$, where C does not depends on m .

Admitting this uniform bound on $\mathbf{F}^{(m)}$, the stability results [12] for linear transport equation can be used as in [11]: there exist (ν, \mathbf{u}) such that up to the extraction of a subsequence, $\beta(\nu^{(m)})$ converges to $\beta(\nu)$ weak* in $L^\infty((0, T) \times D)$ and in $C([0, T]; L^p(D))$ for all $p < +\infty$ and all $\beta \in C^1(\mathbb{R})$, and $\mathbf{u}^{(m)}$ converges to \mathbf{u} in $C([0, T]; H_0^s(D))^3$ for all $s < 1$. However, if \mathbf{F} is related with ψ , where ψ is the solution of (2.1) considering $(K(t), \rho, \sigma)$, it is not obvious that (3.4) holds for $(\mathbf{u}, \nu, \mathbf{F})$ for all given $\varphi \in \mathcal{V}$. This is established by the special argument in Section 4 of [11] if we can show that

$$\int_0^t \int_D \mathbf{F}^m \cdot \varphi d\mathbf{x} d\tau \rightarrow \int_0^t \int_D \mathbf{F} \cdot \varphi d\mathbf{x} d\tau, m \rightarrow \infty, \quad \forall \varphi \in \mathcal{V}, \quad \forall t \in [0, T].$$

As we shall see in the next section, in order to establish these results for $\mathbf{F}^{(m)}$, \mathbf{F} , we need to study the properties of the solution ψ of (2.1). Additional hypotheses

on ρ and σ are also necessary and as remarked in the Section 2 we shall prove the results in a more general framework: we assume hypothesis (i), i.e., K_0 is a Lipschitz domain.

4 Bounds and convergence for the potentials

As remarked in [6], the determination of the charge densities in bio-molecular systems is a non-trivial question which is treated in some computational studies using Hartree-Fock approximation techniques (see [9], [26]). Here we assume explicitly the following hypothesis

- (iii) Let $\rho^{(m)}$ as in the Section 3. Then $\|\rho^{(m)}\|_{L^\infty((0,T)\times D)} \leq C$, where C does not depend on m ;
- (iv) For all $t \in [0, T]$, $\sigma^{(m)}(M^{(m)}(t)\mathbf{x}, t) = \sigma(M^{(m)}(t)\mathbf{x}, t) = \sigma(M(t)\mathbf{x}, t) = \sigma(\mathbf{x}, 0)$, $\forall \mathbf{x} \in \partial K_0$.
- (v) $\|\rho^{(m)} - \rho\|_{L^\infty((0,T)\times D)} \rightarrow 0$, $m \rightarrow +\infty$.

Recalling that $\Psi \in H^1(\partial D)$ and following [6] we consider the problem

$$\begin{aligned} \nabla \cdot (\mathbf{k}(\mathbf{x}, t) \nabla \widehat{\Psi}(\mathbf{x}, t)) &= 0, \quad \mathbf{x} \in D \\ \widehat{\Psi}(\mathbf{x}, t) &= \Psi(\mathbf{x}), \quad \mathbf{x} \in \partial D, \end{aligned}$$

which has a solution $\widehat{\Psi} \in H^{3/2}(D)$ such that $\frac{\partial \widehat{\Psi}_1}{\partial n}, \frac{\partial \widehat{\Psi}_2}{\partial n} \in L^2(\partial K(t))$, $\forall t \in [0, T]$ and

$$\left\| \frac{\partial \widehat{\Psi}}{\partial n}(\cdot, t) \right\|_{0,2,\partial K(t)} \leq C \|\widehat{\Psi}(\cdot, t)\|_{3/2,2,D}, \quad (4.1)$$

where C depends only on the Lipschitz nature of ∂K_0 (see the papers [8], [31] for details). We observe also that $\widehat{\Psi}(\cdot, t) \in L^\infty(D)$ (see [15] or [21]).

Introducing $\psi = \widehat{\psi} + \widehat{\Psi}$, we see that (2.1) may be reformulated as

$$\begin{aligned} \nabla \cdot (\mathbf{k}(\mathbf{x}, t) \nabla \widehat{\psi}(\mathbf{x}, t)) - b(\mathbf{x}, \widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x})) &= \rho(\mathbf{x}, t), \quad \mathbf{x} \in D, \\ \widehat{\psi}_2(\mathbf{x}, t) &= \widehat{\psi}_1(\mathbf{x}, t), \quad \mathbf{x} \in \partial K(t), \\ \widehat{\psi}_2(\mathbf{x}, t) &= 0, \quad \mathbf{x} \in \partial D, \\ k_1 \frac{\partial \widehat{\psi}_1}{\partial n}(\mathbf{x}, t) - k_2 \frac{\partial \widehat{\psi}_2}{\partial n}(\mathbf{x}, t) &= \frac{4\pi e}{T} \sigma(\mathbf{x}, t) - \left(k_1 \frac{\partial \widehat{\Psi}_1}{\partial n}(\mathbf{x}, t) - k_2 \frac{\partial \widehat{\Psi}_2}{\partial n}(\mathbf{x}, t) \right) \\ &= \widehat{\sigma}(\mathbf{x}, t), \quad \text{on } \partial K(t). \end{aligned} \quad (4.2)$$

Let us consider $t \in [0, T]$ fixed and recall the weak formulation for (4.2).

Find $u \in V = H_0^1(D)$ such that $b(\mathbf{x}, u + \widehat{\Psi}) \in L^2(D)$ and satisfying

$$a(u, v) + (N(u), v) = L(v), \quad \forall v \in H_0^1(D), \quad (4.3)$$

where

$$\begin{aligned} a(u, v) &= \int_D k(\mathbf{x}, t) \nabla u \nabla v d\mathbf{x}, \quad (N(u), v) \\ &= \int_D b(\mathbf{x}, u + \widehat{\Psi}) v d\mathbf{x}, \quad L(v) \\ &= \int_{\partial K(t)} \widehat{\sigma} \gamma_0 v ds - \int_D \rho v d\mathbf{x}, \quad \gamma_0 \end{aligned}$$

is the usual trace operator. Here $k(\mathbf{x}, t) = k_1$ if $\mathbf{x} \in K(t)$, $k(\mathbf{x}, t) = k_2$ if $\mathbf{x} \in D \setminus \overline{K(t)}$.

As can be seen in [6] (see also [19]), this problem is equivalent to find the minimum in $H_0^1(D)$ of the functional

$$F(u) = \frac{1}{2} \int_D k(\mathbf{x}, t) |\nabla u|^2 d\mathbf{x} - L(u) + J(u) \quad (4.4)$$

where $J(\cdot)$ is defined as

$$\begin{aligned} J(u) &= k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \{ \cosh(u + \widehat{\Psi}) - \cosh(\widehat{\Psi}) \} d\mathbf{x}, \\ &\text{if } \int_{D \setminus \overline{K(t)}} | \cosh(u + \widehat{\Psi}) - \cosh(\widehat{\Psi}) |^2 d\mathbf{x} < \infty, \\ J(u) &= +\infty \text{ if the square integral is } +\infty. \end{aligned}$$

Below we establish the first bound on ψ derived from (4.3)

Lemma 4.1. *Let us assume the hypothesis (i)–(iv). Then the solution $\psi \in H^1(D)$ of (2.1) belongs to $L^\infty(0, T; H^1(D))$ and $\|\psi\|_{L^\infty(0, T; H^1(D))} \leq M$, where M depends only on $\|\rho\|_{L^\infty((0, T) \times D)}$, $\|\sigma(\cdot, t)\|_{0, 2, \partial K(t)}$, k_1 , k_2 , r_D^{-2} , D , ∂K_0 and Ψ .*

Proof. By $\psi = \widehat{\psi} + \widehat{\Psi}$ we only need to prove the lemma for $\widehat{\psi}$. Using (4.3) we have, for all $t \in [0, T]$,

$$\begin{aligned} & -k_1 \int_{K(t)} |\nabla \widehat{\psi}(\mathbf{x}, t)|^2 d\mathbf{x} - k_2 \int_{D \setminus \overline{K(t)}} |\nabla \widehat{\psi}(\mathbf{x}, t)|^2 d\mathbf{x} + \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \widehat{\psi})(\mathbf{x}, t) ds \\ & = \int_D (\rho \widehat{\psi})(\mathbf{x}, t) d\mathbf{x} + k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) \widehat{\psi}(\mathbf{x}, t) d\mathbf{x} \end{aligned}$$

or

$$\begin{aligned} & \min(k_1, k_2) \int_D |\nabla \widehat{\psi}(\mathbf{x}, t)|^2 d\mathbf{x} \\ & \leq \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \widehat{\psi})(\mathbf{x}, t) ds - \int_D (\rho \widehat{\psi})(\mathbf{x}, t) d\mathbf{x} + \\ & \quad - k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) (\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) d\mathbf{x} \\ & \quad + k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) \widehat{\Psi}(\mathbf{x}, t) d\mathbf{x} \\ & \leq \frac{\|\rho(\cdot, t)\|_{0,2,D}^2}{2\epsilon_1} + \frac{\epsilon_1 \|\widehat{\psi}(\cdot, t)\|_{0,2,D}^2}{2} + \frac{\|\widehat{\sigma}(\cdot, t)\|_{0,2,\partial K(t)}^2}{2\epsilon_2} + \frac{\epsilon_2 \|(\gamma_0 \widehat{\psi})(\cdot, t)\|_{0,2,\partial K(t)}^2}{2} \\ & \quad + k_2 r_D^{-2} \|\widehat{\Psi}(\cdot, t)\|_{\infty,D} \int_{D \setminus \overline{K(t)}} |\sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t))| d\mathbf{x}, \end{aligned}$$

where we have used inequalities of Schwarz and Young. Now, recalling that $\widehat{\psi}$ is the minimum of the functional $F(\cdot)$ defined in (4.4), we have $F(\widehat{\psi}) \leq F(0) = 0$ so that

$$\begin{aligned} & k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} |\sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t))| d\mathbf{x} \\ & \leq k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \cosh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) d\mathbf{x} \\ & \leq k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \cosh(\widehat{\Psi}(\mathbf{x}, t)) d\mathbf{x} \\ & \quad + \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \widehat{\psi})(\mathbf{x}, t) ds - \int_D (\rho \widehat{\psi})(\mathbf{x}, t) d\mathbf{x}, \end{aligned}$$

and

$$\begin{aligned}
& \min(k_1, k_2) \int_D |\nabla \widehat{\psi}(\mathbf{x}, t)|^2 d\mathbf{x} \\
& \leq \frac{\|\rho(\cdot, t)\|_{0,2,D}^2}{2\epsilon_1} + \frac{\epsilon_1 \|\widehat{\psi}(\cdot, t)\|_{0,2,D}^2}{2} + \frac{\|\widehat{\sigma}(\cdot, t)\|_{0,2,\partial K(t)}^2}{2\epsilon_2} \\
& \quad + \frac{\epsilon_2 \|(\gamma_0 \widehat{\psi})(\cdot, t)\|_{0,2,\partial K(t)}^2}{2} + k_2 r_D^{-2} \|\widehat{\Psi}\|_{\infty,D} \int_{D \setminus \overline{K(t)}} \cosh(\widehat{\Psi}(\mathbf{x}, t)) d\mathbf{x} \\
& \quad + \frac{\|\widehat{\Psi}(\cdot, t)\|_{\infty,D}^2 \|\widehat{\sigma}(\cdot, t)\|_{0,2,\partial K(t)}^2}{2\epsilon_3} + \frac{\epsilon_3 \|(\gamma_0 \widehat{\psi})(\cdot, t)\|_{0,2,\partial K(t)}^2}{2} \\
& \quad + \frac{\|\widehat{\Psi}(\cdot, t)\|_{\infty,D} \|\rho(\cdot, t)\|_{0,2,D}^2}{2\epsilon_4} + \frac{\epsilon_4 \|\widehat{\psi}(\cdot, t)\|_{0,2,D}^2}{2}.
\end{aligned}$$

By the trace theorem and Poincaré's inequality there exist constants $\lambda_1 = \lambda_1(\partial K_0, D) > 0$ and $\lambda_2 = \lambda_2(D) > 0$ such that $\|\gamma_0 \widehat{\psi}(\cdot, t)\|_{0,2,\partial K(t)} \leq \lambda_1 \|\widehat{\psi}(\cdot, t)\|_{1,2,D}$ and $\|\widehat{\psi}(\cdot, t)\|_{1,2,D} \leq \lambda_2 \|\nabla \widehat{\psi}(\cdot, t)\|_{0,2,D}$. If we choose $0 < \epsilon < \min(k_1, k_2)/(2\lambda_1^2 \lambda_2^2)$, $\epsilon_1 = \epsilon_4 = \lambda_1^2 \epsilon$, $\epsilon_2 = \epsilon_3 = \epsilon > 0$ we have, for $C_1 = C_1(k_1, k_2, \lambda_1, \lambda_2)$, $C_2 = C_2(k_1, k_2, \lambda_1, \lambda_2)$,

$$\begin{aligned}
& \int_D |\nabla \widehat{\psi}(\mathbf{x}, t)|^2 d\mathbf{x} \\
& \leq C_1 (\|\rho(\cdot, t)\|_{0,2,D}^2 + \|\widehat{\sigma}(\cdot, t)\|_{0,2,\partial K(t)}^2) (1 + \|\widehat{\Psi}(\cdot, t)\|_{\infty,D}^2) \\
& \quad + C_2 k_2 r_D^{-2} \|\widehat{\Psi}(\cdot, t)\|_{\infty,D} \int_{D \setminus \overline{K(t)}} \cosh(\widehat{\Psi}(\mathbf{x}, t)) d\mathbf{x}.
\end{aligned}$$

Finally, Poincaré's inequality gives us

$$\begin{aligned}
\|\widehat{\psi}\|_{L^\infty(0,T;H_0^1(D))} & \leq C^* (|D|^{1/2} \|\rho\|_{L^\infty(0,T;L^\infty(D))} + \|\sigma\|_{0,2,\partial K_0} + \|\widehat{\Psi}\|_{L^\infty(0,T;H^{3/2}(D))}) \\
& (1 + \|\widehat{\Psi}\|_{L^\infty(D,T;L^\infty(D))}) + C^* \|\widehat{\Psi}\|_{L^\infty(D,T;L^\infty(D))} \|\cosh \widehat{\Psi}\|_{L^\infty(D,T;L^1(D))}^{1/2},
\end{aligned}$$

where

$$C^* = \max \left(\lambda_1 C_1^{1/2}, \lambda_1 C_2^{1/2} k_2^{1/2} r_D^{-1}, \frac{4\pi e}{\Lambda}, C \max\{k_1, k_2\} \right).$$

Then

$$\|\psi\|_{L^\infty(0,T;H_0^1(D))} \leq M,$$

where $M = M(C^*, \Psi, \|\rho\|_{L^\infty(0,T;L^\infty(D))}, \|\sigma\|_{0,2,\partial K(t)})$.

The following theorem is central in order to establish an uniform bound for $\mathbf{F}^{(m)}$.

Theorem 4.1. *Let us assume hypotheses (i)–(iv) then the solution $\psi^{(m)}$ of (2.1) related with $(K^{(m)}(t), \rho^{(m)}, \sigma^{(m)})$ satisfies, for each $t \in [0, T]$,*

$$\max \left(\|\psi_1^{(m)}(\cdot, t)\|_{3/2, 2, K^{(m)}(t)}, \|\psi_2^{(m)}(\cdot, t)\|_{3/2, 2, D \setminus \overline{K^{(m)}(t)}} \right) \leq C,$$

where C does not depend on m .

Proof. We only need to obtain the bounds for $\widehat{\psi}^{(m)} = (\widehat{\psi}_1^{(m)}, \widehat{\psi}_2^{(m)})$. Let us consider the problems

$$\begin{aligned} \nabla \cdot (\mathbf{k}(\mathbf{x}, t) \nabla v^{(m)}(\mathbf{x}, t)) &= 0, & \mathbf{x} \in D, \\ v_2^{(m)}(\mathbf{x}, t) &= v_1^{(m)}(\mathbf{x}, t), & \mathbf{x} \in \partial K^{(m)}(t), \\ v_2^{(m)}(\mathbf{x}, t) &= 0, & \mathbf{x} \in \partial D, \\ k_2 \frac{\partial v_2^{(m)}}{\partial n}(\mathbf{x}, t) - k_1 \frac{\partial v_1^{(m)}}{\partial n}(\mathbf{x}, t) &= -\widehat{\sigma}(\mathbf{x}, t), & \mathbf{x} \in \partial K^{(m)}(t), \end{aligned} \quad (4.5)$$

and

$$\begin{aligned} \nabla \cdot (\mathbf{k}(\mathbf{x}, t) \nabla f^{(m)}(\mathbf{x}, t)) - b(\mathbf{x}, f^{(m)}(\mathbf{x}, t) + v^{(m)}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x})) \\ = \rho^{(m)}(\mathbf{x}, t), & \mathbf{x} \in D, \\ f_2^{(m)}(\mathbf{x}, t) &= f_1^{(m)}(\mathbf{x}, t), & \mathbf{x} \in \partial K^{(m)}(t), \\ f_2^{(m)}(\mathbf{x}, t) &= 0, & \mathbf{x} \in \partial D, \\ k_1 \frac{\partial f_1^{(m)}}{\partial n}(\mathbf{x}, t) - k_2 \frac{\partial f_2^{(m)}}{\partial n}(\mathbf{x}, t) &= 0, & \mathbf{x} \in \partial K^{(m)}(t). \end{aligned} \quad (4.6)$$

Using a variational formulation analogous to that in (4.3) we obtain solutions (weak) $v^{(m)}, f^{(m)} \in H_0^1(D)$ of the problems (4.5) and (4.6), respectively. We observe that $f^{(m)} + v^{(m)}$ satisfies (2.1) (in the weak sense), from the uniqueness of the variational solution for this problem, we have $\widehat{\psi}^{(m)} = f^{(m)} + v^{(m)}$.

The Theorem follows from the lemmas below.

Lemma 4.2. *Under hypotheses (i)–(iv) the solution $v^{(m)} \in H_0^1(D)$ of (4.5) has the additional regularity*

$$v_1^{(m)}(\cdot, t) \in H^{3/2}(K^{(m)}(t)), \quad v_2^{(m)}(\cdot, t) \in H^{3/2}(D \setminus \overline{K^{(m)}(t)}),$$

for each $m \geq 0$ and $t \in [0, T]$. Furthermore

$$\max \left(\|v_1^{(m)}\|_{L^\infty(0,T;H^{3/2}(K^{(m)}(t)))}, \|v_2^{(m)}\|_{L^\infty(0,T;H^{3/2}(D \setminus \overline{K^{(m)}(t)}))} \right) \leq C,$$

where C does not depend on m .

Lemma 4.3. Under hypotheses (i)–(iv) the solution $f^{(m)} \in H_0^1(D)$ of (4.6) belongs to $C^0(\overline{D})$ and, for each $t \in [0, T]$,

$$\sup_m \sup_{x \in \overline{D}} |f^{(m)}(\mathbf{x}, t)| < +\infty.$$

Furthermore, $f_1^{(m)}(\cdot, t) \in H^{3/2}(K^{(m)}(t))$, $f_2^{(m)}(\cdot, t) \in H^{3/2}(D \setminus \overline{K^{(m)}(t)})$ and there exists $C > 0$ such that

$$\max \left(\|f_1^{(m)}\|_{L^\infty(0,T;H^{3/2}(K^{(m)}(t)))}, \|f_2^{(m)}\|_{L^\infty(0,T;H^{3/2}(D \setminus \overline{K^{(m)}(t)}))} \right) \leq C,$$

where C does not depend on m . □

Proof of Lemma 4.2. Let us consider $m \geq 0$ and $t \in [0, T]$ fixed and define $\widehat{v}_1^{(m)} = k_1 v_1^{(m)}$, $\widehat{v}_2^{(m)} = k_2 v_2^{(m)}$, then $\widehat{v} = (\widehat{v}_1, \widehat{v}_2)$ satisfies (in the weak sense)

$$\begin{aligned} \Delta \widehat{v}^{(m)}(\mathbf{x}, t) &= 0, & \mathbf{x} \in D, \\ \mu_2 \widehat{v}_2^{(m)}(\mathbf{x}, t) &= \mu_1 \widehat{v}_1^{(m)}(\mathbf{x}, t), & \mathbf{x} \in \partial K^{(m)}(t), \\ \widehat{v}_2^{(m)}(\mathbf{x}, t) &= 0, & \mathbf{x} \in \partial D, \\ \frac{\partial \widehat{v}_2^{(m)}}{\partial n}(\mathbf{x}, t) - \frac{\partial \widehat{v}_1^{(m)}}{\partial n}(\mathbf{x}, t) &= -\widehat{\sigma}(\mathbf{x}, t), & \mathbf{x} \in \partial K^{(m)}(t), \end{aligned} \tag{4.7}$$

where $\mu_2 = k_2^{-1}$ and $\mu_1 = k_1^{-1}$. Following [31], we seek a solution $\widehat{v} = (\widehat{v}_1, \widehat{v}_2)$ in the form

$$\begin{aligned} \widehat{v}_1^{(m)} &= D^{(m)} \zeta^{(m)} + \mu_1 S^{(m)} \varphi^{(m)} \\ \widehat{v}_2^{(m)} &= D^{(m)} \zeta^{(m)} + \mu_2 S^{(m)} \varphi^{(m)} + D_0^{(m)} \chi^{(m)}, \end{aligned} \tag{4.8}$$

for $\zeta^{(m)}(\cdot, t) \in H^1(\partial K^{(m)}(t))$, $\varphi^{(m)}(\cdot, t) \in L^2(\partial K^{(m)}(t))$ and $\chi^{(m)}(\cdot, t) \in H^1(\partial D)$. Here

$$\begin{aligned} (S^{(m)}\varphi^{(m)})(\mathbf{x}, t) &= \int_{\partial K^{(m)}(t)} G(\mathbf{x} - \mathbf{y})\varphi^{(m)}(\mathbf{y}, t)ds(\mathbf{y}), \\ (D^{(m)}\zeta^{(m)})(\mathbf{x}, t) &= \int_{\partial K^{(m)}(t)} \frac{\partial G(\mathbf{x} - \mathbf{y})}{\partial n(\mathbf{y})}\zeta^{(m)}(\mathbf{y}, t)ds(\mathbf{y}), \\ (D_0^{(m)}\chi^{(m)})(\mathbf{x}, t) &= \int_{\partial D} \frac{\partial G(\mathbf{x} - \mathbf{y})}{\partial n(\mathbf{y})}\chi^{(m)}(\mathbf{y}, t)ds(\mathbf{y}). \end{aligned} \quad (4.9)$$

where $G(\mathbf{x}) = \frac{-1}{4\pi|\mathbf{x}|}$. The boundary conditions in (4.7) give us (see [6] for details)

$$\begin{bmatrix} 0 \\ -\widehat{\sigma} \\ 0 \end{bmatrix} = \mathcal{A}^{(m)} \begin{bmatrix} \zeta^{(m)} \\ \varphi^{(m)} \\ \chi^{(m)} \end{bmatrix},$$

where

$$\mathcal{A}^{(m)} = \begin{bmatrix} \mu_2(-\frac{1}{2}I + D_1^{(m)}) - \mu_1(\frac{1}{2}I + D_1^{(m)}) & (\mu_2^2 - \mu_1^2)s^{(m)} & \mu_2(\gamma_e^{(m)}(t)D_0^{(m)}) \\ 0 & \mu_2(\frac{1}{2}I + D_1^{(m)*}) - \mu_1(-\frac{1}{2}I + D_1^{(m)*}) & \gamma_e^{(m)}(t)\left(\frac{\partial D_0^{(m)}}{\partial n}\right) \\ (\gamma_0^{(m)}D^{(m)}) & \mu_2(\gamma_0^{(m)}s^{(m)}) & (\frac{1}{2}I + D_{0,1}^{(m)}) \end{bmatrix}$$

and

$$\begin{aligned} (D_1^{(m)}\zeta^{(m)})(\mathbf{x}, t) &= (p.v. D^{(m)}\zeta^{(m)})(\mathbf{x}, t), \quad x \in \partial K^{(m)}(t) \\ (D_{0,1}^{(m)}\chi^{(m)})(\mathbf{x}, t) &= (p.v. D_0\chi^{(m)})(\mathbf{x}, t), \quad x \in \partial D. \end{aligned}$$

For each $m \in \mathbb{N}$ and $t \in [0, T]$, the operators

$$\begin{aligned} S^{(m)} &: L^2(\partial K^{(m)}(t)) \rightarrow H^1(\partial K^{(m)}(t)) \\ \left(\frac{1}{2}I + D_1^{(m)}\right) &: H^1(\partial K^{(m)}(t)) \rightarrow H^1(\partial K^{(m)}(t)) \\ \left(-\frac{1}{2}I + D_1^{(m)}\right) &: H^1(\partial K^{(m)}(t)) \rightarrow H^1(\partial K^{(m)}(t)) \\ \left(\frac{1}{2}I + D_1^{(m)*}\right) &: H^1(\partial K^{(m)}(t)) \rightarrow L^2(\partial K^{(m)}(t)) \\ \left(-\frac{1}{2}I + D_1^{(m)*}\right) &: H^1(\partial K^{(m)}(t)) \rightarrow L^2(\partial K^{(m)}(t)) \end{aligned}$$

$$\left(\frac{1}{2}I + D_{0,1}^{(m)}\right) : H^1(\partial D) \rightarrow H^1(\partial D)$$

are continuous [8] with operator norms depending only on K_0 , D . Observing that

$$\|\nabla v(\cdot, t)\|_{0,2,\partial K^{(m)}(t)} + \|(\mathbf{n} \cdot \nabla v)(\cdot, t) \cdot \mathbf{n}\|_{0,2,\partial K^{(m)}(t)}$$

is an equivalent norm to $\|\cdot\|_{1,2,\partial D}$ (see [32], Definition 1.9) and using the hypothesis $|\mathbf{x} - \mathbf{y}| > d$, $\forall \mathbf{x} \in \partial K^{(m)}(t)$, $\forall \mathbf{y} \in \partial D$, a direct calculation gives us that the following trace operators

$$\begin{aligned} \gamma_e^{(m)}(t)D_0^{(m)} &: H^1(\partial D) \rightarrow H^1(\partial K^{(m)}(t)), \\ \gamma_0^{(m)}(t)D^{(m)} &: H^1(\partial K^{(m)}(t)) \rightarrow H^1(\partial D) \\ \gamma_0^{(m)}(t)S^{(m)} &: L^2(\partial K^{(m)}(t)) \rightarrow H^1(\partial D) \\ \gamma_e^{(m)}(t) \left(\frac{\partial D_0^{(m)}}{\partial n}\right) &: H^1(\partial D) \rightarrow L^2(\partial K^{(m)}(t)). \end{aligned}$$

are bounded and compact, with operator norms depending only on K_0 , D and d . The compactness follows from an analogous argument as in [7] (Theorems 1.6, 1.7 and 1.10) if we observe that the kernels are continuous.

Now it follows as in the paper by Torres and Welland [31] that $\mathcal{A}^{(m)-1}$ exist and is a bounded operator on $X^{(m)} = H^1(\partial K^{(m)}(t)) \times L^2(\partial K^{(m)}(t)) \times H^1(\partial D)$ to $X^{(m)}$. The operator norm $\|\mathcal{A}^{(m)-1}\|$ depends only on d , K_0 , D . Then

$$\begin{aligned} \left\| \begin{array}{l} \zeta^{(m)}(\cdot, t) \\ \varphi^{(m)}(\cdot, t) \\ \chi^{(m)}(\cdot, t) \end{array} \right\|_{X^{(m)}} &\leq \|\mathcal{A}^{(m)-1}\| \|\widehat{\sigma}(\cdot, t)\|_{0,2,\partial K^{(m)}(t)} \\ &\leq C \|\mathcal{A}^{(m)-1}\| (\|\widehat{\sigma}(\cdot, t)\|_{0,2,\partial K^{(m)}(t)} + \|\widehat{\Psi}(\cdot, t)\|_{3/2,2,D}), \end{aligned}$$

where $X^{(m)}$ is equipped with the product norm.

As observed in [31], the operator norms of the operators in (4.9) and its appropriate inverses depend only on the Lipschitz character of the domain, so that

there exist $L_1 = L_1(\partial K_0) > 0$, $L_2 = L_2(\partial D, \partial K_0) > 0$, such that, $\forall t \in [0, T]$,

$$\begin{aligned} & \|\widehat{v}_1^{(m)}(\cdot, t)\|_{3/2,2,K(t)} \\ & \leq L_1 \max \left(\|\zeta^{(m)}(\cdot, t)\|_{1,2,\partial K^{(m)}(t)}, \|\varphi^{(m)}(\cdot, t)\|_{0,2,\partial K^{(m)}(t)} \right) \\ & \|\widehat{v}_2^{(m)}(\cdot, t)\|_{3/2,2,D \setminus \overline{K^{(m)}(t)}} \\ & \leq L_2 \max \left(\|\zeta^{(m)}(\cdot, t)\|_{1,2,\partial K^{(m)}(t)}, \|\varphi^{(m)}(\cdot, t)\|_{0,2,\partial K^{(m)}(t)}, \|\chi^{(m)}(\cdot, t)\|_{1,2,\partial D} \right). \end{aligned}$$

Hence, using hypothesis (iv), we have established that

$$\max \left(\|v_1^{(m)}\|_{L^\infty(0,T;H^{3/2}(K(t)))}, \|v_2^{(m)}\|_{L^\infty(0,T;H^{3/2}(D \setminus \overline{K(t)})} \right) \leq C,$$

where C does not depends on m . □

Proof of Lemma 4.3. Let us consider $m \geq 0$ and $t \in [0, T]$ fixed. The variational solution $f^{(m)}$ of (4.6) satisfies

$$\begin{aligned} & -k_1 \int_{K^{(m)}(t)} (\nabla \widehat{\psi}^{(m)} \cdot \nabla f^{(m)})(\mathbf{x}, t) d\mathbf{x} - k_2 \int_{D \setminus \overline{K^{(m)}(t)}} (\nabla \widehat{\psi}^{(m)} \cdot \nabla f^{(m)})(\mathbf{x}, t) d\mathbf{x} \\ & = \int_D (\rho^{(m)} \widehat{\psi}^{(m)})(\mathbf{x}, t) d\mathbf{x} + k_2 r_D^{-2} \int_{D \setminus \overline{K^{(m)}(t)}} \sinh(\widehat{\psi}^{(m)}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x})) \widehat{\psi}^{(m)}(\mathbf{x}, t) d\mathbf{x}. \end{aligned}$$

Then the Young's inequality gives us that

$$\begin{aligned} & \frac{k_1}{2} \int_{K^{(m)}(t)} |\nabla f^{(m)}(\mathbf{x}, t)|^2 d\mathbf{x} + \frac{k_2}{2} \int_{D \setminus \overline{K^{(m)}(t)}} |\nabla f^{(m)}(\mathbf{x}, t)|^2 d\mathbf{x} \\ & \leq - \int_D (\rho^{(m)} \widehat{\psi}^{(m)})(\mathbf{x}, t) d\mathbf{x} - k_2 r_D^{-2} \int_{D \setminus \overline{K^{(m)}(t)}} \sinh(\widehat{\psi}^{(m)}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x})) \widehat{\psi}^{(m)}(\mathbf{x}, t) d\mathbf{x}. \end{aligned}$$

Using a similar calculation as in Lemma 4.1, the bound established there for $\|\widehat{\psi}^{(m)}(\cdot, t)\|_{1,2,D}$ and hypothesis (iii), we have $\|f^{(m)}(\cdot, t)\|_{1,2,D} \leq C$, where C does not depends on m and t . Observing that D is a C^2 -domain, standard elliptic estimates (see Chapter 14, Theorem 2.1 in [21]) show us that $f^{(m)}(\cdot, t) \in C^0(\overline{D})$.

Let us define

$$R^{(m)}(t) = \sup_{\mathbf{x} \in \overline{D}} |f^{(m)}(\mathbf{x}, t)|, \quad \text{then} \quad \exists R(t) = \sup_m R^{(m)}(t) < +\infty.$$

In effect, let us suppose that $R(t) = +\infty$, then there exist a subsequence $f^{(m_k)}$ related with $(\mathbf{u}^{(m_k)}, v^{(m_k)})$, $\widehat{\psi}^{(m_k)}$, such that $R^{(m_k)}(t) \rightarrow +\infty$ with $k \rightarrow +\infty$.

Let us choose $\mathbf{x}^{(m_k)} \in \overline{D}$ such that $f^{(m_k)}(\mathbf{x}^{(m_k)}) = R^{(m_k)}(t)$, then there exists $\delta^{(m_k)} > 0$ such that

$$|f^{(m_k)}(\mathbf{x}^{(m_k)}, t) - f^{(m_k)}(\mathbf{x}, t)| < 2^{-m_k} \quad \text{if } \mathbf{x} \in B(\mathbf{x}^{(m_k)}, \delta^{(m_k)}) \cap \overline{D}.$$

Recalling that $f^{(m_k)}|_{\partial D} = 0$, if we take $k \rightarrow \infty$ we have, by the bound $\|f^{(m_k)}(\cdot, t)\|_{0,2,D} \leq C$, $\delta_{(m_k)} \rightarrow 0$. In this case $\|\nabla f^{(m_k)}(\cdot, t)\|_{0,2,D} \rightarrow +\infty$, which contradicts the bound $\|f^{(m_k)}(\cdot, t)\|_{1,2,D} \leq C$.

If we set $h^{(m)} = b(\mathbf{x}, f^{(m)} + v^{(m)} + \widehat{\Psi}) + \rho^{(m)}$ we have $h^{(m)}(\cdot, t) \in L^2(D)$, for all $t \in [0, T]$ and $m \geq 0$. Extending $h^{(m)}$ to be zero outside of D and setting $g^{(m)} = G * h^{(m)}$ we have $\Delta g^{(m)}(\cdot, t) = h^{(m)}(\cdot, t)$ a.e. in D and $g^{(m)}(\cdot, t) \in H^2(D)$; if

$$g_1^{(m)}(\cdot, t) = g^{(m)}(\cdot, t)|_{K^{(m)}(t)}, \quad g_2^{(m)}(\cdot, t) = g^{(m)}(\cdot, t)|_{D \setminus \overline{K^{(m)}(t)}},$$

we have

$$g_1^{(m)}(\cdot, t)|_{\partial K^{(m)}(t)} \in H^1(\partial K^{(m)}(t)), \quad g_2^{(m)}(\cdot, t)|_{\partial D} \in H^1(\partial D)$$

(see the proof of Theorem B in [20]), while $\frac{\partial g_1^{(m)}}{\partial n}(\cdot, t) - \frac{\partial g_2^{(m)}}{\partial n}(\cdot, t) = 0$ a.e. in $\partial K^{(m)}(t)$. Hence the solution $f^{(m)}$ of (4.6) can be written as (see [6])

$$\begin{pmatrix} \widehat{f}_1^{(m)} \\ \widehat{f}_2^{(m)} \end{pmatrix} = \begin{pmatrix} g_1^{(m)} \\ g_2^{(m)} \end{pmatrix} + \mathcal{H}^{(m)} \mathcal{A}^{(m)-1} \begin{bmatrix} g_1^{(m)}|_{\partial K^{(m)}(t)} \\ 0 \\ g_2^{(m)}|_{\partial D} \end{bmatrix},$$

where

$$\mathcal{H}^{(m)} = \begin{bmatrix} D^{(m)} & \mu_1 S^{(m)} & 0 \\ D^{(m)} & \mu_2 S & D_0^{(m)} \end{bmatrix}$$

and $\widehat{f}_1^{(m)} = k_1 f_1^{(m)}$, $\widehat{f}_2^{(m)} = k_2 f_2^{(m)}$. The operators in $\mathcal{H}^{(m)}$ were defined in (4.9) and $\mathcal{A}^{(m)}$ was defined in Lemma 4.2. The uniform estimate in the $H^{3/2}$ -norm follows in a similar way as in Lemma 4.2 if we get uniform $H^{3/2}$ -bounds for $g^{(m)}(\cdot, t)$ and H^1 -bounds for $g_1^{(m)}(\cdot, t)|_{\partial K^{(m)}(t)}$, $g_2^{(m)}(\cdot, t)|_{\partial D}$. From the continuous imbedding $H^2(D) \subset H^{3/2}(D)$ and by the boundedness of the operator $F: L^2(D) \rightarrow H^2(D)$, where $Fh^{(m)}(\cdot, t) = G * h^{(m)}(\cdot, t) = g^{(m)}(\cdot, t)$ (see the proof of Theorem 1 in [32]) we have the bounds

$$\begin{aligned} & \|g^{(m)}(\cdot, t)\|_{3/2,2,D} \\ & \leq \lambda(\|I_{D \setminus \overline{K^{(m)}(t)}}\| k_2 r_D^{-2} \sinh(f^{(m)} + v^{(m)} + \widehat{\Psi})(\cdot, t)\|_{0,2,D} + \|\rho^{(m)}(\cdot, t)\|_{0,2,D}). \end{aligned}$$

where λ depends only on D . A direct calculation and the above estimate give us

$$\begin{aligned} & \|g_1^{(m)}(\cdot, t)\|_{1,2,\partial K(t)} \leq \lambda' \|\rho^{(m)}(\cdot, t)\|_{0,2,D} \|g_2^{(m)}(\cdot, t)\|_{1,2,\partial D} \\ & \leq \lambda'' (\|I_{D \setminus \overline{K^{(m)}(t)}} k_2 r_D^{-2} \sinh(f^{(m)} + v^{(m)} + \widehat{\Psi})(\cdot, t)\|_{0,2,D} + \|\rho^{(m)}(\cdot, t)\|_{0,2,D}) \end{aligned}$$

where $\lambda' = \lambda'(K_0, \lambda)$, $\lambda'' = \lambda''(D, \lambda)$. Hence we need only obtain an uniform estimate for

$$\|I_{D \setminus \overline{K^{(m)}(t)}} k_2 r_D^{-2} \sinh(f^{(m)} + v^{(m)} + \widehat{\Psi})(\cdot, t)\|_{0,2,D \setminus \overline{K^{(m)}(t)}}.$$

We observe that

$$\begin{aligned} \sinh(f^{(m)} + v^{(m)} + \widehat{\Psi}) &= \sinh(f^{(m)} + v^{(m)}) \cosh(\widehat{\Psi}) \\ &+ \cosh(f^{(m)} + v^{(m)}) \sinh(\widehat{\Psi}). \end{aligned}$$

The result follows using the fact that $\Psi \in L^\infty(D)$, recalling that by Lemma 4.3 we have $\sup_m \|f^{(m)}(\cdot, t)\|_{C^0(\overline{D})} < +\infty$ and from the estimate

$$\begin{aligned} \|\cosh(v_2^{(m)})(\cdot, t)\|_{0,2,D \setminus \overline{K^{(m)}(t)}} &\leq C \exp\left(C \|v_2^{(m)}(\cdot, t)\|_{3/2,2,D \setminus \overline{K^{(m)}(t)}}\right) \\ &\leq C, \end{aligned} \quad (4.10)$$

(see [6] and Lemma 4.2). □

Corollary 4.1. $\int_0^T \|\mathbf{F}^{(m)}(\cdot, t)\|_{0,2,D}^2 dt \leq C$, where C does not depend on m .

As remarked in the preceding section the solution of (3.4) is constructed as a limit of a sequence of the appropriate approximation solutions and depends on the respective convergence of the force term $\mathbf{F}^{(m)}$ to \mathbf{F} in $L^2((0, T) \times D)^3$. Below we show in detailed manner how to do this. We begin with a technical lemma.

Lemma 4.4. *Let us assume hypothesis (i)–(v) and consider $\psi^{(m)}$, ψ the solutions of (2.1) related with $(K^{(m)}(t), \rho^{(m)}, \sigma^{(m)})$ and $(K(t), \rho, \sigma)$, respectively. Then, $\|\psi^{(m)}(\cdot, t) - \psi(\cdot, t)\|_{1,2,D} \rightarrow 0$, with $m \rightarrow +\infty$, for each $t \in [0, T]$.*

Proof. Let us set $\eta_m = \psi^{(m)} - \psi$. The variational formulation for $\psi, \psi^{(m)}$ gives us

$$\begin{aligned} & - k_1 \int_{K^{(m)}(t)} (\nabla \widehat{\psi}^{(m)} \cdot \nabla \eta_m)(\mathbf{x}, t) d\mathbf{x} \\ & - k_2 \int_{D \setminus \overline{K^{(m)}(t)}} (\nabla \widehat{\psi}^{(m)} \cdot \nabla \eta_m)(\mathbf{x}, t) d\mathbf{x} + \int_{\partial K^{(m)}(t)} (\widehat{\sigma} \gamma_0 \eta_m)(\mathbf{x}, t) ds \\ & = \int_D (\rho_m \eta_m)(\mathbf{x}, t) d\mathbf{x} + k_2 r_D^{-2} \int_{D \setminus \overline{K^{(m)}(t)}} \sinh(\widehat{\psi}^{(m)}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) \eta_m(\mathbf{x}, t) d\mathbf{x} \end{aligned}$$

and

$$\begin{aligned} & - k_1 \int_{K(t)} (\nabla \widehat{\psi} \cdot \nabla \eta_m)(\mathbf{x}, t) d\mathbf{x} \\ & - k_2 \int_{D \setminus \overline{K(t)}} (\nabla \widehat{\psi} \cdot \nabla \eta_m)(\mathbf{x}, t) d\mathbf{x} + \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \eta_m)(\mathbf{x}, t) ds \\ & = \int_D (\rho \eta_m)(\mathbf{x}, t) d\mathbf{x} + k_2 r_D^{-2} \int_{D \setminus \overline{K(t)}} \sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) \eta_m(\mathbf{x}, t) d\mathbf{x}. \end{aligned}$$

If we define $A^{(m)}(t) = K^{(m)}(t) \cap K(t)$, $B^{(m)}(t) = D \setminus \overline{(K(t) \cup K^{(m)}(t))}$ we have, after subtracting the above expressions,

$$\begin{aligned} & - k_1 \int_{A^{(m)}(t)} |\nabla \eta_m(\mathbf{x}, t)|^2 d\mathbf{x} - k_2 \int_{B^{(m)}(t)} |\nabla \eta_m(\mathbf{x}, t)|^2 d\mathbf{x} + \int_{\partial K^{(m)}(t)} (\widehat{\sigma} \gamma_0 \eta_m)(\mathbf{x}, t) ds \\ & - \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \eta_m)(\mathbf{x}, t) ds = \int_D (\rho^{(m)} - \rho)(\mathbf{x}, t) \eta_m(\mathbf{x}, t) d\mathbf{x} \\ & + k_2 r_D^{-2} \int_{B^{(m)}(t)} 2 \cosh\left(\frac{\eta_m}{2} + \widehat{\psi} + \widehat{\Psi}\right)(\mathbf{x}, t) \sinh\left(\frac{\eta_m(\mathbf{x}, t)}{2}\right) \eta_m(\mathbf{x}, t) d\mathbf{x} \\ & - k_1 \int_{K(t) \setminus A^{(m)}(t)} |\nabla \widehat{\psi}(\mathbf{x}, t)| |\nabla \eta_m(\mathbf{x}, t)| d\mathbf{x} \\ & - k_2 \int_{K^{(m)}(t) \setminus A^{(m)}(t)} |\nabla \widehat{\psi}(\mathbf{x}, t)| |\nabla \eta_m(\mathbf{x}, t)| d\mathbf{x} \\ & + k_1 \int_{K^{(m)}(t) \setminus A^{(m)}(t)} |\nabla \widehat{\psi}^{(m)}(\mathbf{x}, t)| |\nabla \eta_m(\mathbf{x}, t)| d\mathbf{x} \\ & + k_2 \int_{K(t) \setminus A^{(m)}(t)} |\nabla \widehat{\psi}^{(m)}(\mathbf{x}, t)| |\nabla \eta_m(\mathbf{x}, t)| d\mathbf{x} \\ & - k_2 r_D^{-2} \int_{K^{(m)}(t) \setminus A^{(m)}(t)} \sinh(\widehat{\psi}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) \eta_m(\mathbf{x}, t) d\mathbf{x} \\ & + k_2 r_D^{-2} \int_{K(t) \setminus A^{(m)}(t)} \sinh(\psi^{(m)}(\mathbf{x}, t) + \widehat{\Psi}(\mathbf{x}, t)) \eta_m(\mathbf{x}, t) d\mathbf{x}. \end{aligned}$$

Using Young's and Hölder's inequalities and positiveness of the second integral on the right side, we have

$$\begin{aligned}
& k_1 \int_{A^{(m)}(t)} |\nabla \eta_m(\mathbf{x}, t)|^2 d\mathbf{x} + k_2 \int_{B^{(m)}(t)} |\nabla \eta_m(\mathbf{x}, t)|^2 d\mathbf{x} \\
& \leq \left| \int_{\partial K^{(m)}(t)} (\widehat{\sigma}^{(m)} \gamma_0 \eta_m)(\mathbf{x}, t) ds - \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \eta_m)(\mathbf{x}, t) ds \right| \\
& \quad + |D|^{1/2} \|(\rho^{(m)} - \rho)(\cdot, t)\|_{0, \infty, D} \|\eta_m(\cdot, t)\|_{0, 2, D} \\
& \quad + k_1 |K(t) \setminus A^{(m)}(t)|^{1/6} \|(\nabla \widehat{\psi})(\cdot, t)\|_{0, 3, K(t)} \|\nabla \eta_m(\cdot, t)\|_{0, 2, K(t)} \\
& \quad + k_2 |K^{(m)}(t) \setminus A^{(m)}(t)|^{1/6} \|\nabla \widehat{\psi}(\cdot, t)\|_{0, 3, K^{(m)}(t)} \|\nabla \eta_m(\cdot, t)\|_{0, 2, K^{(m)}(t)} \\
& \quad + k_1 |K^{(m)}(t) \setminus A^{(m)}(t)|^{1/6} \|\nabla \widehat{\psi}^{(m)}(\cdot, t)\|_{0, 3, K^{(m)}(t)} \|\nabla \eta_m(\cdot, t)\|_{0, 2, K^{(m)}(t)} \\
& \quad + k_2 |K(t) \setminus A^{(m)}(t)|^{1/6} \|\nabla \widehat{\psi}^{(m)}(\cdot, t)\|_{0, 3, K(t)} \|\nabla \eta_m(\cdot, t)\|_{0, 2, K(t)} \\
& \quad + k_2 r_D^{-2} |K^{(m)}(t) \setminus A^{(m)}(t)|^{1/4} \|\sinh(\widehat{\psi} + \widehat{\Psi})(\cdot, t)\|_{0, 2, B^{(m)}(t)} \|\eta_m(\cdot, t)\|_{0, 4, B^{(m)}(t)} \\
& \quad + k_2 r_D^{-2} |K(t) \setminus A^{(m)}(t)|^{1/4} \|\sinh(\widehat{\psi}^{(m)} + \widehat{\Psi})(\cdot, t)\|_{0, 2, B_m(t)} \|\eta_m(\cdot, t)\|_{0, 4, B_m(t)}.
\end{aligned}$$

Now, following [11],

$$\begin{aligned}
& \sup_{t \in (0, T)} (|M^{(m)}(t) - M(t)| + |\dot{M}^{(m)}(t) - \dot{M}(t)|) \\
& \leq C_T |v_p^{(m)} \mathbf{u}^{(m)} - v_p \mathbf{u}|_{L^\infty(0, T; L^2(D))^n} \rightarrow 0,
\end{aligned} \tag{4.11}$$

with $m \rightarrow +\infty$, since $v_p^{(m)} \mathbf{u}^{(m)}$ converges to $v_p \mathbf{u}$ in $C([0, T]; L^2(D))^3$. Hence $|K^{(m)}(t) \setminus A^{(m)}(t)|$, $|K(t) \setminus A^{(m)}(t)| \rightarrow 0$, $m \rightarrow +\infty$. We observe that, by the Theorem 4.1 and the Sobolev imbedding $H^{1/2}(K^{(m)}(t)) \subset L^3(K^{(m)}(t))$,

$$\begin{aligned}
\|\nabla \psi_1^{(m)}(\cdot, t)\|_{0, 3, K^{(m)}(t)} & \leq C \|\nabla \psi_1^{(m)}(\cdot, t)\|_{1/2, 2, K^{(m)}(t)} \\
& \leq \|\psi_1^{(m)}(\cdot, t)\|_{3/2, 2, K^{(m)}(t)} \\
& \leq C,
\end{aligned}$$

where C does not depend on m . Similarly, using the Sobolev embedding $H^1(D) \subset L^4(D)$, Lemma 4.1 and hypothesis (iii) and (iv) we see that uniform bounds to $\|\eta_m\|_{0, 2, D}$, $\|\eta_m\|_{0, 4, D}$ are available. From hypothesis (v), $\|(\rho^{(m)} - \rho)(\cdot, t)\|_{0, \infty, D} \rightarrow 0$. Theorem 4.1 and similar argument as in (4.10) give us that

$$\begin{aligned}
\|\sinh(\psi^{(m)} + \widehat{\Psi})(\cdot, t)\|_{0, 2, B^{(m)}(t)} & \leq C \exp(C \|\psi^{(m)}(\cdot, t)\|_{3/2, 2, B^{(m)}(t)}) \\
& \leq C,
\end{aligned}$$

where C does not depend on m .

Using the rigidness of the particle we have

$$\begin{aligned} & \int_{\partial K^{(m)}(t)} (\widehat{\sigma}^{(m)} \gamma_0 \eta_m)(\mathbf{x}, t) ds(\mathbf{x}) - \int_{\partial K(t)} (\widehat{\sigma} \gamma_0 \eta_m)(\mathbf{x}, t) ds(\mathbf{x}) \\ &= \int_{\partial K_0} \widehat{\sigma}^{(m)}(M^{(m)}(t)\mathbf{y}, t) \gamma_0 (\eta_m(M^{(m)}(t)\mathbf{y}, t) - \eta_m(M(t)\mathbf{y}, t)) ds(\mathbf{y}) \quad (4.12) \\ &+ \int_{\partial K_0} \gamma_0 \eta_m(M(t)\mathbf{y}, t) (\widehat{\sigma}^{(m)}(M^{(m)}(t)\mathbf{y}, t) - \widehat{\sigma}(M(t)\mathbf{y}, t)) ds(\mathbf{y}). \end{aligned}$$

Now, we introduce, for each m , the usual C_0^∞ -regularization $\{\eta_m^{(h)}\}$ for η_m such that $\|(\eta_m^{(h)} - \eta_m)(\cdot, t)\|_{1,1,D} \rightarrow 0$ when $h \rightarrow 0$. Observing that

$$\|\gamma_0 \eta_m^{(h)}(M^{(m)}(t)\mathbf{y}, t) - \gamma_0 \eta_m^{(h)}(M(t)\mathbf{y}, t)\|_{0,2,\partial K_0} \rightarrow 0 \quad \text{with } m \rightarrow +\infty$$

the first integral in (4.12) tends to zero with $m \rightarrow +\infty$, as can be seen using the usual trace theorem. Analogously we have the same result for the second integral in (4.12), recalling that $\widehat{\sigma} = \frac{4\pi e}{T} \sigma + \left(k_1 \frac{\partial \widehat{\Psi}_1}{\partial n} - k_2 \frac{\partial \widehat{\Psi}_2}{\partial n}\right)$, the hypothesis (iv) and regularity $\widehat{\Psi}(\cdot, t) \in H^{3/2}(D)$. \square

Theorem 4.2. *Let us consider $\varphi^{(m)} \in H^1((0, T) \times D)^3$ such that $\varphi^{(m)} \rightarrow \varphi$ strongly in $C([0, T]; L^2(D))^3$ and $\varphi \in H^1((0, T) \times D)^3$. Then*

$$\lim_{m \rightarrow +\infty} \int_0^t \int_D (\mathbf{F}^{(m)} \cdot \varphi^{(m)})(\mathbf{x}, \tau) d\mathbf{x} d\tau = \int_0^t \int_D (\mathbf{F} \cdot \varphi)(\mathbf{x}, \tau) d\mathbf{x} d\tau,$$

for all $t \in (0, T)$.

Proof. Recalling that $\mathbf{F}^{(m)} = \frac{\Lambda}{4\pi e} (\rho_2^{(m)} + r_D^{-2} k_2 \sinh(\psi_2^{(m)})) (\nabla \psi_2^{(m)}) I_{D \setminus \overline{K^{(m)}(t)}}$, we can write

$$\begin{aligned} \mathbf{F}^{(m)} \cdot \varphi_m - \mathbf{F} \cdot \varphi &= C_1 I_{D \setminus \overline{K^{(m)}(t)}} (\varphi^{(m)} - \varphi) \cdot \nabla \psi^{(m)} \sinh(\psi^{(m)}) \\ &+ C_1 \varphi \cdot (I_{D \setminus \overline{K(t)}} \nabla \psi \sinh(\psi) \\ &- I_{D \setminus \overline{K^{(m)}(t)}} \nabla \psi^{(m)} \sinh(\psi^{(m)})) \\ &+ C_2 I_{D \setminus \overline{K^{(m)}(t)}} \rho^{(m)} (\varphi^{(m)} - \varphi) \cdot \nabla \psi^{(m)} \\ &+ C_2 \varphi \cdot \left(I_{D \setminus \overline{K(t)}} \rho \nabla \psi - I_{D \setminus \overline{K^{(m)}(t)}} \rho^{(m)} \nabla \psi^{(m)} \right), \end{aligned} \quad (4.13)$$

where $C_1 = \frac{k_2 r_D^{-2} \Lambda}{4\pi e}$ and $C_2 = \frac{\Lambda}{4\pi e}$.

In what follows we shall estimate the integral of each term above.

$$\begin{aligned} & \int_0^t \int_D I_{D \setminus \overline{K^{(m)}(\tau)}} ((\varphi_m - \varphi) \cdot \nabla \psi^{(m)} \sinh(\psi^{(m)}))(\mathbf{x}, \tau) d\mathbf{x} d\tau \\ & \leq \|\varphi^{(m)} - \varphi\|_{C([0, T]; L^2(D))}^3 \\ & \int_0^t \|\nabla \psi^{(m)}(\cdot, \tau)\|_{0, 3, D \setminus \overline{K^{(m)}(\tau)}} \|\sinh(\psi^{(m)})(\cdot, \tau)\|_{0, 6, D \setminus \overline{K^{(m)}(\tau)}} d\tau \\ & \leq C \|\varphi^{(m)} - \varphi\|_{C([0, T]; L^2(D))}^3 \rightarrow 0, \end{aligned}$$

with $m \rightarrow +\infty$. Here we have used the Hölder's inequality, the embeddings $H^{1/2}(D) \subset L^3(D)$ and $H^{1/2}(D) \subset H^{3/2}(D)$, Theorem 4.1 jointly with a similar estimate as in (4.10) and the convergence $\|\varphi^{(m)} - \varphi\|_{C([0, T]; L^2(D))}^3 \rightarrow 0$.

$$\begin{aligned} & \int_0^t \int_D \left(\varphi \cdot \left(I_{D \setminus \overline{K(\tau)}} \nabla \psi \sinh(\psi) - I_{D \setminus \overline{K^{(m)}(\tau)}} \nabla \psi^{(m)} \sinh(\psi^{(m)}) \right) \right) (\mathbf{x}, \tau) d\mathbf{x} d\tau \\ & \leq \int_0^t \left[\int_{B^{(m)}(\tau)} |\varphi(\mathbf{x}, \tau) \cdot (\nabla \psi \sinh(\psi) - \nabla \psi^{(m)} \sinh(\psi^{(m)}))(\mathbf{x}, \tau)| d\mathbf{x} \right. \\ & \quad + \int_{K^{(m)}(\tau) \setminus K(\tau)} |\varphi(\mathbf{x}, \tau) \cdot (\nabla \psi(\mathbf{x}, \tau) \sinh(\psi(\mathbf{x}, \tau)))| d\mathbf{x} \\ & \quad \left. + \int_{K(\tau) \setminus K^{(m)}(\tau)} |\varphi(\mathbf{x}, \tau) \cdot (\nabla \psi^{(m)}(\mathbf{x}, \tau) \sinh(\psi^{(m)}(\mathbf{x}, \tau)))| d\mathbf{x} \right] d\tau. \end{aligned} \tag{4.14}$$

Writing

$$\begin{aligned} & \nabla \psi \sinh(\psi) - \nabla \psi^{(m)} \sinh(\psi^{(m)}) \\ & = \sinh(\psi) (\nabla \psi - \nabla \psi^{(m)}) + \nabla \psi^{(m)} (\sinh(\psi) - \sinh(\psi^{(m)})) \end{aligned}$$

we have

$$\begin{aligned} & \int_0^t \int_{B^{(m)}(\tau)} \varphi(\mathbf{x}, \tau) \cdot (\nabla \psi \sinh(\psi) - \nabla \psi^{(m)} \sinh(\psi^{(m)}))(\mathbf{x}, \tau) d\mathbf{x} d\tau \\ & \leq \int_0^t \|\sinh(\psi(\cdot, \tau))\|_{0, 4, B^{(m)}(\tau)} \|\varphi(\cdot, \tau)\|_{0, 4, B^{(m)}(\tau)} \\ & \quad \|(\nabla \psi - \nabla \psi^{(m)})(\cdot, \tau)\|_{0, 2, B^{(m)}(\tau)} d\tau \\ & \quad + \int_0^t \|\varphi(\cdot, \tau)\|_{0, 6, B^{(m)}(\cdot, \tau)} \|\nabla \psi^{(m)}(\cdot, \tau)\|_{0, 3, B^{(m)}(\tau)} \\ & \quad \|(\sinh(\psi) - \sinh(\psi^{(m)}))(\cdot, \tau)\|_{0, 2, B^{(m)}(\tau)} d\tau. \end{aligned} \tag{4.15}$$

Now, observing that if

$$\begin{aligned} \psi^{(m)} - \psi \geq 0, \quad \cosh^2(\psi^{(m)} + \theta(\psi - \psi^{(m)})) &\leq \cosh^2(\psi) \quad \text{and if} \\ \psi^{(m)} - \psi < 0, \quad \cosh^2(\psi^{(m)} + \theta(\psi - \psi^{(m)})) &\leq \cosh^2(\psi - \psi^{(m)}), \end{aligned}$$

we have, by Schwarz's inequality

$$\begin{aligned} &\|(\sinh(\psi) - \sinh(\psi^{(m)}))(\cdot, t)\|_{0,2,B^{(m)}(t)}^2 \\ &= \int_{B^{(m)}(t)} \left(\int_0^1 (\psi - \psi^{(m)})(\mathbf{x}, t) \cosh(\psi^{(m)}(\mathbf{x}, t) + \theta(\psi - \psi^{(m)})(\mathbf{x}, t)) d\theta \right)^2 d\mathbf{x} \\ &\leq \int_{B^{(m)}(t)} \int_0^1 (\psi - \psi^{(m)})^2(\mathbf{x}, t) \cosh^2(\psi^{(m)}(\mathbf{x}, t) + \theta(\psi - \psi^{(m)})(\mathbf{x}, t)) d\theta d\mathbf{x} \\ &\leq C \|(\psi - \psi^{(m)})(\cdot, t)\|_{0,4,D}^2 \rightarrow 0, \end{aligned}$$

where we have used a similar estimate as in (4.10), Theorem 4.1 and Lemma 4.4. Hence the dominated convergence theorem, Theorem 4.1 and Lemma 4.4 gives us that the terms in (4.15) tends to zero with $m \rightarrow +\infty$. Passing the terms in (4.14) to the limit $m \rightarrow +\infty$, using (4.11) and Theorem 4.1 we obtain the desired result for the second term in (4.13). The convergence of the others terms in (4.13) to zero follows from a completely similar way, using hypothesis (iii), (iv) and (v), Theorem 4.1 and Lemma 4.4. \square

Acknowledgment. The first author was supported by a Doctoral Scholarship of the CNPq-Brasil.

REFERENCES

- [1] A. Ajdari, Electro-osmosis on inhomogeneous charged surfaces, *Physical Review Letters*, **75**(4) (1995), 755-758.
- [2] A. Ajdari and D. Long, Symmetry properties of the electrophoretic motion of patterned colloidal particles. *Physical Review Letters*, **81**(7) (1998), 1529-1533.
- [3] S. Allison, C. Cheng and D. Stigter, The Length Dependence of Translational Diffusion, Free solution Electrophoretic Motion and Electrophoretic Tether Force of Rigid Rod-Like Model Duplex DNA. *Biophys. J.*, **81**(5) (2001), 2558–2568.
- [4] J.L. Anderson, Colloidal transport by interfacial forces. *Ann. Re. Fluid Mech.*, **21** (1989), 61–99.

- [5] L. Bedin and M. Thompson, Motion of charged particles in ionized fluids via the Poisson-Boltzmann equation. To appear, *Math. Models and Methods Appl. Sci.*
- [6] L. Bedin, M. Thompson and M.T. Vilhena, On the Poisson-Boltzmann equation on non-smooth domains. *Int. J. Diff. Eq. App.*, **8**(4) (2003), 327–360.
- [7] D. Colton and R. Kress, *Integral Equations Methods in Scattering Theory*, John Wiley, New York, (1988).
- [8] M. Costabel, Boundary integral operators on Lipschitz domains: elementary results. *SIAM J. Math. Anal.*, **19** (1985), 613–626.
- [9] L.E. Dardenne, A.S. Werneck, M. Oliveira and P.M. Bisch, Reassociation of Fragments Using Multicentered Multipolar Expansions: Peptide Junction Treatment to Investigate Electrostatic Properties of Proteins. *J. Comp. Chem.*, **22** (2001), 689–701.
- [10] M. Daune, *Molecular Biophysics: Structures in Motion*. Oxford University Press, Oxford (1999).
- [11] B. Desjardins and M.J. Esteban, Existence of weak solutions for the motion of rigid bodies in a viscous fluid. *Arch. Rat. Mech. Anal.*, **146** (1999), 59–71.
- [12] R.J. DiPerna and P.L. Lions, Ordinary differential equations, transport theory and Sobolev spaces. *Inventiones mathematicae*, **98**(3) (1989), 511–547.
- [13] M.C. Fair and A. Ajdari, Electrophoresis of dumbbell-like colloidal particles. *Int. J. Multiphase Flow*. **16**(4) (1990), 663–679.
- [14] M. Ferreira, S.L.L. Verardi and J.M. Machado, Electroosmotic Pumping Rectangular Microchannels: a Numerical Treatment by the Finite Element Method. *Appl. Math.*, **2** (2002), 10–15.
- [15] D. Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, (1977).
- [16] A. Yu. Grosberg and A.R. Khokhlov, *Statistical Physics of Macromolecules*, AIP Press, New York, (1994).
- [17] M.D. Gunsburger, H.C. Lee and G.A. Seregin, Global existence of weak solutions for the viscous incompressible flows around a moving rigid body in three dimensions. *J. Math. Fluid Mech.*, **2** (2000), 219–266.
- [18] N.M. Gunther, *La Théorie du Potentiel et ses Applications Aux Problèmes Fondamentaux de la Physique Mathématique*. Gauthier-Villards, Paris, (1934).
- [19] M.J. Holst, *The Poisson-Boltzmann Equation: Analysis and multilevel numerical solution* (Monograph based on Ph.D.Thesis: Multilevel Methods for the Poisson-Boltzmann Equation). tech. rep., California Institute of Technology, (1994).
- [20] D. Jerison and C.E. Kenig, The inhomogeneous Dirichlet problem in Lipschitz domains. *J. Func. Anal.*, **130** (1995), 161–219.

- [21] O.A. Ladyzhenskaya and N.N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*. Academic Press, New York, (1968).
- [22] V.G. Levich, *Physicochemical Hydrodynamics*, Prentice Hall, Englewood Cliffs, New Jersey, (1962).
- [23] P. Lions, *Mathematical Topics in Fluid Mechanics: Incompressible Models*. Oxford Interscience, London, (1984).
- [24] D. Long, J.L. Viovy and A. Ajdari, Simultaneous action of electric fields and nonelectric forces on a polyelectrolyte: motion and deformation. *Physical Review Letters*, **76**(20) (1996), 3858–3861.
- [25] C. Miranda, *Partial Differential Equations of the Elliptic Type*. Springer-Verlag, Berlin, (1970).
- [26] H. Nakamura, Roles of electrostatic interactions in proteins. *Quart. J. Biophysics*, **29** (1996), 1–90.
- [27] A. Sellier, A note on the electrophoresis of a uniformly charged particle. *Q. J. Mech. Appl. Math.*, **55**(4) (2002), 561–572.
- [28] Y. Solomentsev and J.L. Anderson, Electrophoresis of slender particles. *J. Fluid Mech.*, **279** (1994), 197–215.
- [29] T.N. Swaminathan and H.H. Hu, Coiling of a long chain of particles undergoing electrophoresis, *Preprint-University of Pennsylvania*, (2001).
- [30] M. Teubner, The motion of charged particles in electrical fields. *J. Chem. Phys.*, **76**(11) (1982), 5564–5573.
- [31] R.H. Torres and G.V. Welland, The Helmholtz equation and transmission problems with Lipschitz interfaces. *Indiana J. Math.*, **42** (1993), 1457–1486.
- [32] G. Verchota, Layer Potentials and Regularity for the Dirichlet Problem for Laplace's Equation in Lipschitz Domains. *J. Func. Anal.*, **59** (1984), 572–611.