

Post-Traumatic Stress Disorder and severe maternal morbidity: is there an association?

Carina R. Angelini,¹ Rodolfo C. Pacagnella,¹ Mary A. Parpinelli,¹ Carla Silveira,¹ Carla B. Andreucci,^{1,II} Elton C. Ferreira,¹ Juliana P. Santos,¹ Dulce M. Zanardi,¹ Renato T. Souza,¹ Jose G. Cecatti^{1,*}

¹Departamento de Ginecologia e Obstetrícia, Faculdade de Ciências Médicas, Universidade Estadual de Campinas, Campinas, SP, BR. ^{II}Departamento de Medicina, Universidade Federal de São Carlos, São Carlos, SP, BR.

Angelini CR, Pacagnella RC, Parpinelli MA, Silveira C, Andreucci CB, Ferreira EC, et al. Post-Traumatic Stress Disorder and severe maternal morbidity: is there an association? Clinics. 2018;73:e309

*Corresponding author. E-mail: cecatti@unicamp.br

OBJECTIVE: To evaluate the occurrence of Post-Traumatic Stress Disorder among women experiencing a severe maternal morbidity event and associated factors in comparison with those without maternal morbidity.

METHODS: In a retrospective cohort study, 803 women with or without severe maternal morbidity were evaluated at 6 months to 5 years postpartum for the presence of Post-Traumatic Stress Disorder. Interviews were conducted by telephone and electronic data was stored. Data analysis was carried out by using χ^2 , Fisher's Exact test, and logistic regression analysis.

RESULTS: There was no significant change in the prevalence of Post-Traumatic Stress Disorder related to a previous severe maternal morbidity experience. There were also no differences in diagnostic criteria for severe maternal morbidity (hypertensive syndromes, hemorrhage, surgical intervention or intensive care unit admission required, among other management criteria). Low parity (2.5-fold risk) and increasing age were factors associated with Post-Traumatic Stress Disorder.

CONCLUSIONS: A severe maternal morbidity episode is not associated with Post-Traumatic Stress Disorder symptoms within five years of the severe maternal morbidity event and birth. However, a more advanced maternal age and primiparity increased the risk of Post-Traumatic Stress Disorder. This does not imply that women who had experienced a severe maternal morbidity event did not suffer or need differentiated care.

KEYWORDS: Post-Traumatic Stress Disorder; Maternal Morbidity; Maternal Near Miss.

INTRODUCTION

Severe maternal morbidity (SMM) and maternal near miss (MNM) have increasingly been referred to as important markers of maternal health and a resource to evaluate the quality of obstetric care (1,2). A continuum of morbidity exists between a healthy pregnancy and maternal death; survivors of the conditions that may lead to death are considered MNM cases (2).

Pregnancy and childbirth, even without complications, are potentially stressful situations and women may experience depression for prolonged periods after childbirth (3). Unplanned pregnancies, a negative subjective experience during childbirth, fear of maternal or infant death, limited support offered by the health team were associated with a higher prevalence of post-traumatic stress disorder (PTSD) (4,5).

Women experiencing obstetric complications have a higher risk of developing clinical and mental health problems (6,7).

Similar to their children, they are more physically and socially vulnerable, and have higher risk of death up to one year postpartum (8). Less severe but also important repercussions, including decrease in functioning (9), delay in resuming sexual intercourse (10), and deterioration in quality of life may occur in association with SMM (11). Feelings such as fear of death, hopelessness and loss of female identity have been identified in other studies (12,13).

Post-traumatic stress disorder (PTSD) is associated with higher levels of physical, social and work-related impairment, leading to higher health service use and increased medical costs (14). During pregnancy and the postpartum period, the disorder may have a major impact, since maternal mental health issues may negatively interfere in infant cognitive and language development (15) and may affect the partner and family dynamics (6).

PTSD is defined as a syndrome that can be triggered after exposure to a traumatic event involving death or a consistent threat of death or severe damage to the self or others (criterion A), which elicits a subjective response of intense fear, helplessness or impotence in the midst of the situation and horror. In response to such a traumatic event, a triad of symptoms characterizes the disorder: reliving the traumatic event (criterion B), avoiding stimuli associated with trauma (Criterion C), and having symptoms of increased psychic excitability (irritability, difficulty sleeping and concentrating) (14).

Copyright © 2018 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is properly cited.

No potential conflict of interest was reported.

Received for publication on July 30, 2017. **Accepted for publication on** October 27, 2017

DOI: 10.6061/clinics/2018/e309



Severe preeclampsia and SMM were recently identified as a risk factors for PTSD (6,16-18). Therefore, understanding which disorders affect women after a SMM episode and the repercussions on their mental health is a key-point. The purpose of this study was to identify the prevalence of PTSD after an SMM event and the factors associated with this diagnosis.

■ METHODS

This is an analysis of a retrospective cohort study investigating a SMM group and a control group without any maternal complication. Details on the method of this study have already been published elsewhere (19). Briefly, its general purpose was to evaluate several aspects of the woman's life following a SMM episode: general and reproductive health conditions, quality of life, sexual activity, capabilities and functioning, mental health, and growth and development of the child born in that pregnancy. It was conducted in a tertiary referral maternity hospital in Campinas, in the southeast of Brazil.

The SMM group was selected from women admitted to intensive care unit (ICU) during pregnancy that met the criteria for Potentially Life-Threatening Conditions (PLTC) or Maternal Near Miss (MNM) as recommended by the WHO (2), generally recognized as Severe Maternal Morbidity. The control group was composed of women with a full-term birth without any severe complications. Each control was randomly selected on the same year of birth as SMM cases to maintain a balance between groups regarding time after birth. All women who fulfilled the above requirements and had given birth from 2008 to 2012 were eligible for the study. The period between childbirth and the interview ranged from 6 months to 5 years. Data collection was conducted by chart review to identify criteria for PLTC and MNM, birth and hospital discharge conditions; and by telephone interview applying instruments to evaluate quality of life and PTSD. Information for PTSD assessment have already been collected and validated through telephone interviews (20).

From electronic clinical records, 1,157 women fulfilling selection criteria were identified. A team trained in teleresearch, using standardized CATI (computer-assisted telephone interview) procedures, searched for these women. Several phone calls were done at different times, and 840 eligible women were successfully traced and invited to participate in the study. Women agreeing to participate recorded their consent and were then interviewed by telephone, or were scheduled for interview at a more opportune date and hour. The interview was recorded and data immediately inserted into the electronic Lime Survey[®] database. For quality control, a tele-researcher supervisor randomly selected a sample fraction of about 5% of cases and listened to the recording, checking responses marked in the electronic database. Corrections were made when necessary, and any possible errors found were reported to the interviewer.

Sample size was determined by previous studies and included all the outcome variables and was considered to have enough power to assess differences in quality of life. Presuming that 50% of women had severe physical and emotional problems during the first year following delivery, an absolute difference of 11% between both groups, a Type I error of 5% and a Type II error of 10%, each group would require 337 women, totaling 674.

PTSD was evaluated by an instrument, the PTSD Checklist-Civilian Version (PCL-C) (21), already adapted and validated in Brazil for Portuguese language (22). The current study is not a validation of the content or psychometric properties of this instrument, but only its application in a sample population of women in the postpartum period. It was composed of 17 questions with a severity scale ranging from 1 to 5 (nothing to a lot). To make the diagnosis, 4 types of symptoms are required: Criterion A (triggering condition), Criterion B (questions 1 to 5), Criterion C (questions 6 to 12) and Criterion D (questions 13 to 17). Childbirth may be considered a stressful event in itself, including the experience of pregnancy and stress of labor.

For instrument analysis, three different criteria of analysis were used. For cluster analysis, PTSD diagnostic criteria were considered according to the DSM-IV: having a clinically significant symptom from group B, having three symptoms from group C and two from group D. To be considered clinically significant, a symptom received a score that should be equal to or greater than three. For remaining analyses, a total score was used and cutoff points were defined (40 and 50) (23) that refer to the sum of scores in each question of the instrument, with values ranging from a minimum of 17 to a maximum of 85 points. The higher the score, the worse was the response to PTSD.

Data was submitted to strict checking procedures for completion and internal consistency. Missing and/or incorrect data were confirmed in recordings, medical charts or alternatively by a repeat phone call to the woman, and then corrected in the database.

For statistical analysis, sociodemographic, pregnancy and childbirth characteristics were compared between groups. The prevalence of PTSD was then determined using the 3 different sets of criteria (cluster, cutoff ≥ 40 and ≥ 50) for both groups with differences assessed using Chi-square or Fisher's Exact tests. For women diagnosed with PTSD, the distribution between those with or without SMM and according to sociodemographics and pregnancy characteristics was provided for the three criteria. In addition, prevalences of PTSD between groups and according to the time since delivery were reported for all three sets of criteria, and according to diagnostic criteria used for SMM. Finally, multinomial regression analysis was performed to identify conditions independently associated with PTSD. For these procedures, the SPSS package was used (version 20; IBM, Armonk, NY, USA). The study was approved by the local Institutional Review Board (letter of approval CEP 233/2009). All participants recorded an audio consent form and/or signed a written version of the consent. The principles of the Helsinki Declaration of 1975, which was revised in 1983, were followed.

■ RESULTS

Initially 1,157 women were eligible for the study, 840 of them were traced by telephone and 803 responded to the PCL-C instrument for PTSD assessment. Among the 37 missing cases, 22 did not respond the questionnaire due to diverse personal reasons and 15 had already died in different periods after childbirth (9 from late maternal causes). The tracing rate was 72.6% and the response rate was 69.4%, although 13 women did not fully complete the questionnaire. Among 790 women with complete interviews, 381 had experienced




Figure 1 - Flow chart of women in the study inclusion. PTSD: post-traumatic stress disorder; SMM: severe maternal morbidity.

an SMM episode, while 409 had no severe complications and served as the control group (Figure 1).

The majority of women with SMM were older and more than 35% were over 35, as shown in Table 1. Groups did not show differences in parity, ethnicity, school education, socio-economic status, time since childbirth, having a partner and neonatal outcomes. The proportion of women with SMM undergoing cesarean section was very high (82%) in comparison with controls (48%). Briefly, women with SMM were proportionally older and had higher parity.

Table 2 shows the prevalence of PTSD using distinct criteria. By the method for cluster symptom analysis, 54.3% of women with SMM and 51.8% of those without SMM developed PTSD. In addition, no significant differences were found between groups when using the cut-off point ≥ 40 for subsyndromic PTSD, or a cutoff ≥ 50 .

Table 3 shows only women diagnosed with PTSD by the three different sets of criteria comparatively with and without SMM and according to sociodemographics and pregnancy characteristics. Among women with PTSD, maternal age ≥ 35 years was more common in the SMM group, i.e. older women had SMM more frequently. A significantly higher proportion of women with PTSD by the cutoff ≥ 50 had given birth within 2 years of the interview in the SMM group.

The proportion was significantly higher in more than 80% women with SMM and PTSD, than in women without SMM. For the remaining variables, there were no significant differences between groups.

Table 4 reports that there was no difference in the prevalence of PTSD in women with and without SMM at any time periods considered. The prevalence of PTSD showed no statistically significant differences according to the major determinants of SMM (hypertension, hemorrhage, surgical intervention or ICU admission). Therefore, the presence of a specific SMM condition did not modify the prevalence of PTSD in this sample (Data not shown).

In Table 5, the variables significantly associated with diagnosis of PTSD by a cutoff ≥ 50 were primiparity and maternal age above 40 years. Women with only one child had a 2.5 times higher risk of having PTSD than multiparous women. In contrast, in women aged 20 to 39 the risk of PTSD decreased by 50 to 60%. The presence of SMM was not independently associated with the risk of PTSD.

■ DISCUSSION

The current study did not find an association between SMM and PTSD in this population, at least within one to



Table 1 - Sociodemographics and pregnancy characteristics of women according to a previous experience of severe maternal morbidity (SMM).

Sociodemographics and pregnancy characteristics	With SMM		Without SMM		p-value*
	n	%	n	%	
Maternal age (years)					0.001
<20	16	4.2	24	5.9	
20 a 24	58	15.2	80	19.6	
25 a 29	86	22.6	110	26.9	
30 a 34	86	22.6	108	26.4	
35 a 39	82	21.5	54	13.2	
≥40	53	13.9	33	8.1	
Total	381		409		
Parity					0.244
1	98	31.8	109	34.0	
2	93	30.2	103	32.1	
3	54	17.5	63	19.6	
≥4	63	20.5	46	14.3	
Total	308		321		
Skin color/ethnicity					0.204
White	197	51.7	193	47.2	
Non white	184	48.3	216	52.8	
Total	381		409		
Literacy (years)					0.238
Up to 4	24	6.7	14	3.6	
5-8	103	28.9	102	26.4	
9-11	194	54.5	233	60.4	
≥12	35	9.8	37	9.6	
Total	356		386		
Socioeconomic classification					0.193
Class A+B	117	38.6	140	43.8	
Class C+D+E	186	61.4	180	56.3	
Total	303		320		
Time since delivery (y)					0.302
1 - 2	152	39.9	178	43.5	
3 - 5	229	60.1	231	56.5	
Total	381		409		
With a partner					0.926
Yes	255	82.2	265	83.1	
No	53	17.2	54	16.9	
Total	308		319		
Mode of delivery					<0.001
Vaginal	67	18.0	211	51.6	
Cesarean section	305	82.0	198	48.4	
Total	372		409		
Neonatal outcome					0.054
Alive	299	95.8	391	98.2	
Death	13	4.2	7	1.8	
Total	312		398		

* χ^2 test.

5 years of the SMM event. This does not mean that women, who had a SMM episode but no signs of PTSD, did not suffer any traumatic experience. The evaluation of PTSD at different times in the postpartum period showed that a higher proportion of women with a SMM episode were older and underwent cesarean section more frequently, which was probably related to the occurrence of maternal morbidity.

The association between advanced maternal age and SMM, as well as increased maternal age and higher likelihood of cesarean section, have already been described. Pregnancies at a later maternal age are more likely to have complications, mainly due to maternal premorbid conditions, as hypertension and diabetes (24), what may partly explain our results. However, even when indicated for maternal and fetal conditions, cesarean section is not a risk-free intervention. It can be very difficult to distinguish between cause

Table 2 - Prevalence of PTSD among women according to a previous experience of severe maternal morbidity (SMM) using three different criteria for analysis of the PCL-C instrument.

Criteria for PTSD	With SMM		Without SMM		p-value*
Criteria for Cluster	n	%	n	%	
Positive	207	54.3	212	51.8	0.482
Negative	174	45.7	197	48.2	
Cut-off ≥40	n	%	n	%	0.183
Positive	196	51.4	191	49.4	
Negative	185	48.6	218	54.1	
Cut-off ≥50	n	%	n	%	0.479
Positive	113	29.7	112	27.4	
Negative	268	70.3	297	72.6	
Total	381		409		

* χ^2 test.

and effect in the relationship between cesarean delivery and maternal morbidity (25).

The prevalence of PTSD was no different between morbidity groups using any criteria of analysis. Therefore, in this sample, the occurrence of PTSD was not influenced by SMM. These results differ from a study that described preeclampsia as an important factor predisposing to PTSD (17). Another cohort study evaluating British women at 6 to 8 weeks postpartum also identified a higher risk of PTSD symptoms among those experiencing severe maternal morbidity (18).

On the other hand, cesarean delivery appeared to be strongly associated with SMM, regardless the presence of PTSD with any criteria. This could probably be due to the impact of a potential threat to maternal or neonatal life during childbirth. It is not the birth event itself that may lead a woman to trauma, but her perception of this event. A birth may seem perfect and properly cared by the professional, but it is not necessarily felt the same way by the woman. In addition to performing a technically suitable procedure, it is vital to consider the issues that extend beyond traditional clinical parameters, towards a more personal care centered on the woman herself.

In women with an SMM episode, PTSD was higher with the cutoff ≥ 50 within 2 years of childbirth. This was the only positive result found for time since the occurrence of the morbid episode, suggesting that stress may decrease over time.

Several factors may contribute to PTSD in the postpartum period. Subjective suffering during childbirth and obstetric emergencies were the most important risk factors for PTSD found in a recent systematic review. Furthermore, emergency cesarian section, instrumental vaginal delivery and neonatal complications were also important predictors of PTSD (26).

A meta-analysis estimated that the prevalence of PTSD was 5-fold higher in women at risk (27). The perception of a negative subjective experience of childbirth and of limited support during childbirth were also factors strongly associated with PTSD after childbirth in another meta-analysis (28). They also found a strong association between difficulties in maternal coping and postpartum depression. In our study, we had no information on women's perception of maternal care received. However, we can infer that it may have been a more positive experience, or it was perceived no differently from women without complications, considering that SMM was not identified as an important factor for PTSD.

Estimates of PTSD after childbirth have varied enormously among studies. Many factors, including subjective elements,

**Table 3** - Women diagnosed with PTSD with or without severe maternal morbidity (SMM) using 3 different sets of criteria for the PCL-C instrument according to some sociodemographic and pregnancy characteristics.

Sociodemographic and pregnancy characteristics	Criteria for Cluster		Cut-off ≥ 40		Cut-off ≥ 50	
	With SMM %	Without SMM %	With SMM %	Without SMM %	With SMM %	Without SMM %
Maternal age (years)		n=419		n=387		n=225
<20	3.9	7.5	4.6	6.8	2.7	7.1
20 a 24	15.9	19.8	15.8	19.4	12.4	19.6
25 a 29	24.2	26.9	21.4	27.2	25.7	24.1
30 a 34	22.2	27.8	23.0	28.3	22.1	30.4
35 a 39	21.3	10.8	21.4	11.5	26.5	12.5
≥ 40	12.6	7.1	13.8	6.8	10.6	6.3
p-value		0.009		0.014		0.027
Parity		n=348		n=324		n=191
1	31.0	31.6	30.2	30.9	26.9	28.4
2	24.6	33.3	24.1	32.1	26.0	36.8
3	18.7	19.8	19.1	21.0	24.0	16.8
≥ 4	25.7	15.3	26.5	16.0	28.1	17.9
p-value		0.070		0.103		0.115
Skin color/ethnicity		n=419		n=387		n=225
White	46.9	45.3	48.0	46.1	46.0	46.4
Non white	53.1	54.7	52.0	53.9	54.0	53.6
p-value		0.746		0.710		0.951
Literacy (years)		n=402		n=373		n=217
Up to 4	5.6	3.9	5.9	4.3	7.5	3.6
5-8	32.1	26.7	33.5	26.1	31.8	30.0
9-11	55.1	62.6	53.5	61.7	52.3	58.2
≥ 12	7.1	6.8	7.0	8.0	8.4	8.2
p-value		0.452		0.326		0.606
Socioeconomic classification		n=303		n=322		n=190
Class A + B	34.9	40.7	33.8	40.1	32.6	38.9
Class C + D + E	65.1	59.3	66.3	59.9	67.4	61.1
p-value		0.269		0.236		0.364
Time since delivery (y)		n=381		n=387		n=225
1 - 2	58.5	54.7	58.7	55.0	53.3	46.4
3 - 5	41.5	45.3	41.3	45.0	46.7	53.6
p-value		0.440		0.463		0.039
With a partner		n=308		n=323		n=191
Yes	78.4	83.5	79.0	82.0	76.0	81.1
No	21.6	16.5	21.0	18.0	24.0	18.9
p-value		0.221		0.500		0.399
Mode of delivery		n=414		n=382		n=222
Vaginal	15.9	51.9	16.2	51.3	18.2	58.9
Cesarean section	84.2	48.1	83.8	48.7	81.8	41.1
p-value		0.000		0.000		0.000
Neonatal outcome		n=379		n=346		n=204
Alive	94.7	97.6	94.3	97.3	94.7	96.4
Death	5.3	2.4	5.7	2.7	5.3	3.6
p-value*		0.142		0.153		0.560

* χ^2 test.

childbirth care models and personal history of the woman have been identified as important predictors of PTSD. Therefore, it is difficult to point out a specific risk factor for PTSD or make a single estimate of its prevalence following a SMM event.

In the current study, it is noteworthy that women were evaluated within 5 years of the event. This may be a positive differential in comparison to other studies on PTSD after childbirth. Even after stratifying women by time between childbirth and the interview, in this study no association between SMM and PTSD was observed. However, the study has not enough power to identify difference between groups

according to time since the morbidity event, which could be a study limitation. Additional studies are recommended for the investigation of potential short-term and long-term effects of SMM on mental health of women and their families, preferably with multicenter studies with a wider population-based representativeness.

Our results indicate that a severe maternal morbidity experience is not more significantly associated with PTSD symptoms within five years of the SMM event. Findings did not imply that suffering did not occur or that differentiated and integrated care is not required in this group of women.



Table 4 - Prevalence of PTSD among women with or without severe maternal morbidity (SMM) according to criteria used for analysis of the PCL-C instrument stratified by time since delivery until interview.

Time since delivery	With SMM		Without SMM		p-values*
	n	%	n	%	
5 years (2008)					
Criteria for cluster	11	37.9	13	41.9	0.752
Cut-off ≥40	11	37.9	13	41.9	0.752
Cut-off ≥50	7	24.1	4	12.9	0.261
4 years (2009)					
Criteria for cluster	59	53.2	47	50.0	0.653
Cut-off ≥40 *	55	49.5	41	43.6	0.404
Cut-off ≥50 *	31	27.9	21	22.3	0.421
3 years (2010)					
Criteria for cluster	51	57.3	56	52.8	0.532
Cut-off ≥40	49	55.1	51	48.1	0.334
Cut-off ≥50	30	33.7	27	25.5	0.208
2 years (2011)					
Criteria for cluster	51	57.3	72	58.5	0.857
Cut-off ≥40	46	51.7	63	51.2	0.947
Cut-off ≥50	26	29.2	47	38.2	0.174
1 year (2012)					
Criteria for cluster	35	55.6%	24	43.6%	0.196
Cut-off ≥40	35	55.6%	23	41.8%	0.136
Cut-off ≥50	19	30.2%	13	23.6%	0.427

* χ^2 or Fisher's Exact test.

Table 5 - Factors independently associated with the diagnosis of PTSD using a cutoff point ≥50 as a criterion in a multivariate regression analysis.

Predictors	OR	95% CI		p-value
Maternal age				
<20	0.483	0.157	1.484	0.204
20 a 24	0.376	0.170	0.832	0.016
25 a 29	0.448	0.219	0.918	0.028
30 a 34	0.461	0.228	0.930	0.030
35 a 39	0.428	0.208	0.882	0.021
≥40	Ref.			
Parity				
1	2.553	1.408	4.629	0.002
2	1.636	0.956	2.799	0.072
3	1.335	0.754	2.365	0.322
≥4	Ref.			
Ethnicity				
White	1.085	0.758	1.555	0.655
Non white	Ref.			
Schooling (years)				
Up to 4	0.931	0.331	2.619	0.892
5-8	1.033	0.497	2.150	0.930
9-11	0.969	0.519	1.810	0.922
≥12	Ref.			
Socioeconomic classification				
Class A + B	1.215	0.809	1.825	0.348
Class C + D + E	Ref.			
With a partner				
Yes	1.490	0.945	2.349	0.086
No	Ref.			
Maternal morbidity				
Without SMM	1.094	0.766	1.564	0.621
With SMM	Ref.			

ACKNOWLEDGMENTS

This study was fully sponsored by the Brazilian National Research Council (CNPq, research grant 471142/2011-5). However, the organ played no role whatsoever in planning, implementing, analyzing, discussing the results or writing the current manuscript.

AUTHOR CONTRIBUTIONS

Angelini CR, Cecatti JG, Pacagnella RC, Parpinelli MA and Andreucci CB conceived and designed the experiments. Angelini CR, Andreucci CB, Pacagnella RC, Silveira C, Parpinelli MA, Ferreira EC, Santos JP, Zanardi DM and Cecatti JG performed the experiments. Angelini CR, Pacagnella RC and Cecatti JG analyzed the data. Angelini CR wrote the paper. Angelini CR, Andreucci CB, Pacagnella RC, Silveira C, Parpinelli MA, Ferreira EC, Zanardi DM, Santos JP, Souza RT and Cecatti JG reviewed, provided suggestions and agreed on the final version of the manuscript.

REFERENCES

- Firoz T, Chou D, von Dadelszen P, Agrawal P, Vanderkruik R, Tunçalp O, et al. Measuring maternal health: focus on maternal morbidity. *Bull World Health Organ.* 2013;91(10):794-6, <http://dx.doi.org/10.2471/BLT.13.117564>.
- Say L, Souza JP, Pattinson RC; WHO working group on Maternal Mortality and Morbidity classifications. Maternal near miss—towards a standard tool for monitoring quality of maternal health care. *Best Pract Res Clin Obstet Gynaecol.* 2009;23(3):287-96, <http://dx.doi.org/10.1016/j.bpobgyn.2009.01.007>.
- Brown S, Lumley J. Physical health problems after childbirth and maternal depression at six to seven months postpartum. *BJOG.* 2000;107(10):1194-201, <http://dx.doi.org/10.1111/j.1471-0528.2000.tb11607.x>.
- Adewuya AO, Ologun YA, Ibigbami OS. Post-traumatic stress disorder after childbirth in Nigerian women: prevalence and risk factors. *BJOG.* 2006;113(3):284-8, <http://dx.doi.org/10.1111/j.1471-0528.2006.00861.x>.
- Creeedy DK, Shochet IM, Horsfall J. Childbirth and the development of acute trauma symptoms: incidence and contributing factors. *Birth.* 2000; 27(2):104-11, <http://dx.doi.org/10.1046/j.1523-536x.2000.00104.x>.
- Hinton L, Locock L, Knight M. Support for mothers and their families after life-threatening illness in pregnancy and childbirth: a qualitative study in primary care. *Br J Gen Pract.* 2015;65(638):e563-9, <http://dx.doi.org/10.3399/bjgp15X686461>.
- Koblinsky M, Chowdhury ME, Moran A, Ronsmans C. Maternal morbidity and disability and their consequences: neglected agenda in maternal health. *J Health Popul Nutr.* 2012;30(2):124-30, <http://dx.doi.org/10.3329/jhpn.v30i2.11294>.
- Filippi V, Ganaba R, Baggaley RF, Marshall T, Storeng KT, Sombie I, et al. Health of women after severe obstetric complications in Burkina Faso: a longitudinal study. *Lancet.* 2007;370(9595):1329-37, [http://dx.doi.org/10.1016/S0140-6736\(07\)61574-8](http://dx.doi.org/10.1016/S0140-6736(07)61574-8).
- Silveira C, Parpinelli MA, Pacagnella RC, Andreucci CB, Ferreira EC, Angelini CR, et al. A cohort study of functioning and disability among women after severe maternal morbidity. *Int J Gynaecol Obstet.* 2016;134(1): 87-92, <http://dx.doi.org/10.1016/j.ijgo.2015.10.027>.
- Andreucci CB, Cecatti JG, Pacagnella RC, Silveira C, Parpinelli MA, Ferreira EC, et al. Does Severe Maternal Morbidity Affect Female Sexual Activity and Function? Evidence from a Brazilian Cohort Study. *PLoS One.* 2015;10(12):e0143581, <http://dx.doi.org/10.1371/journal.pone.0143581>.
- Waterstone M, Wolfe C, Hooper R, Bewley S. Postnatal morbidity after childbirth and severe obstetric morbidity. *BJOG.* 2003;110(2):128-33, <http://dx.doi.org/10.1046/j.1471-0528.2003.02151.x>.
- Tunçalp Ö, Hindin MJ, Adu-Bonsaffoh K, Adanu R. Listening to women's voices: the quality of care of women experiencing Severe Maternal Morbidity, in Accra, Ghana. *PLoS One.* 2012;7(8):e44536, <http://dx.doi.org/10.1371/journal.pone.0044536>.
- Souza JP, Cecatti JG, Parpinelli MA, Krupa F, Osis MJ. An emerging "maternal near-miss syndrome": narratives of women who almost died during pregnancy and childbirth. *Birth.* 2009;36(2):149-58, <http://dx.doi.org/10.1111/j.1523-536X.2009.00313.x>.
- APA – American Psychiatric Association. *Diagnostic and Statistic Manual of Mental Disorders, DSM-IV-TR (4th Ed. Text Revision)*. APA, 2000. 943p.
- Sohr-Preston SL, Scaramella LV. Implications of timing of maternal depressive symptoms for early cognitive and language development. *Clin Child Fam Psychol Rev.* 2006;9(1):65-83, <http://dx.doi.org/10.1007/s10567-006-0004-2>.
- Furuta M, Sandall J, Bick D. A systematic review of the relationship between severe maternal morbidity and post-traumatic stress disorder. *BMC Pregnancy Childbirth.* 2012;12:125, <http://dx.doi.org/10.1186/1471-2393-12-125>.
- Engelhard IM, van Rij M, Boullart I, Ekhart THA, Spaanderman ME, van den Hout MA, et al. Posttraumatic stress disorder after pre-eclampsia: an exploratory study. *Gen Hosp Psychiatry.* 2002;24(4):260-4, [http://dx.doi.org/10.1016/S0163-8343\(02\)00189-5](http://dx.doi.org/10.1016/S0163-8343(02)00189-5).
- Furuta M, Sandall J, Cooper D, Bick D. The relationship between severe maternal morbidity and psychological health symptoms at 6-8 weeks postpartum: a prospective cohort study in one English maternity unit. *BMC Pregnancy Childbirth.* 2014;14:133, <http://dx.doi.org/10.1186/1471-2393-14-133>.



19. Cecatti JG, Souza JP, Parpinelli MA, Haddad SM, Camargo RS, Pacagnella RC, et al. Brazilian network for the surveillance of maternal potentially life threatening morbidity and maternal near-miss and a multidimensional evaluation of their long term consequences. *Reprod Health*. 2009;6:15, <http://dx.doi.org/10.1186/1742-4755-6-15>.
20. Shlomi Polachek I, Dulitzky M, Margolis-Dorfman L, Simchen MJ. A simple model for prediction postpartum PTSD in high-risk pregnancies. *Arch Womens Ment Health*. 2016;19(3):483-90, <http://dx.doi.org/10.1007/s00737-015-0582-4>.
21. Weathers FW, Litz BT, Herman DS, Huska JA, Keane TM. The PTSD Checklist (PCL): Reliability, validity, and diagnostic utility. Paper currents at the 9th Annual Conference of the International Society for Traumatic Stress Studies, San Antonio, Tx, USA, 1993.
22. Berger W, Mendlowicz MV, Souza WF, Figueira I. [Semantic equivalence of the Portuguese version of the Post-Traumatic Stress Disorder Checklist - Civilian Version (PCL-C) for screening of post-traumatic stress disorder]. *Rev Psiquiatr Rio Gd Sul*. 2004;26(2):167-75, <http://dx.doi.org/10.1590/S0101-81082004000200006>.
23. Andrykowski MA, Cordova MJ, Studts JL, Miller TW. Posttraumatic stress disorder after treatment for breast cancer: prevalence of diagnosis and use of the PTSD Checklist-Civilian Version (PCL-C) as a screening instrument. *J Consult Clin Psychol*. 1998;66(3):586-90, <http://dx.doi.org/10.1037/0022-006X.66.3.586>.
24. Laopaiboon M, Lumbiganon P, Intarut N, Mori R, Ganchimeg T, Vogel JP, et al. Advanced maternal age and pregnancy outcomes: a multicountry assessment. *BJOG*. 2014;121 Suppl 1:49-56, <http://dx.doi.org/10.1111/1471-0528.12659>.
25. Souza JP, Gulmezoglu A, Lumbiganon P, Laopaiboon M, Carroli G, Fawole B, et al. Caesarean section without medical indications is associated with an increased risk of adverse short-term maternal outcomes: the 2004-2008 WHO Global Survey on Maternal and Perinatal Health. *BMC Med*. 2010;8:71, <http://dx.doi.org/10.1186/1741-7015-8-71>.
26. Andersen LB, Melvaer LB, Videbech P, Lamont RF, Joergensen JS. Risk factors for developing post-traumatic stress disorder following childbirth: a systematic review. *Acta Obstet Gynecol Scand*. 2012;91(11):1261-72, <http://dx.doi.org/10.1111/j.1600-0412.2012.01476.x>.
27. Grekin R, O'Hara MW. Prevalence and risk factors of postpartum post-traumatic stress disorder: a meta-analysis. *Clin Psychol Rev*. 2014;34(5):389-401, <http://dx.doi.org/10.1016/j.cpr.2014.05.003>.
28. Ayers S, Bond R, Bertullies S, Wijma K. The aetiology of post-traumatic stress following childbirth: a meta-analysis and theoretical framework. *Psychol Med*. 2016;46(6):1121-34, <http://dx.doi.org/10.1017/S0033291715002706>.