

Janaina Patricio de Lima<sup>1</sup>Irineu Tadeu Velasco<sup>1</sup>Denise Frediane Barbeiro<sup>1</sup>Eliane Schochat<sup>1</sup>**Keywords**

Audiometry, Evoked Response

Hearing

Brain ischemia

Sepsis

Brain Stem

**Descritores**

Audiometria de Resposta Evocada

Audição

Isquemia Encefálica

Sepse

Tronco Encefálico

**Auditory brainstem response in gerbils  
submitted to ischemia and sepsis*****Potencial evocado auditivo de tronco encefálico  
em gerbils submetidos à isquemia e sepse*****ABSTRACT**

**Introduction:** An ischemic stroke is a clinical condition that affects thousands of people worldwide. As a result of this injury, neuronal death can be observed, and in the natural course of recovery, the individual may develop sepsis. Sepsis is a systemic inflammatory response that can lead the patient to death. To assess the clinical condition of a patient with this condition, the Auditory Brainstem Response (ABR) can be useful, since it is not an invasive procedure, it is a fast technique and it can be done at the bedside. **Purpose:** To assess auditory brainstem response (ABR) latency values in gerbils subjected to ischemia and sepsis. **Methods:** ABR values were collected from 72 adult male gerbils, which were divided into six groups: control, sepsis, ischemia, sham, ischemia with sepsis, and sham with sepsis. For the induction of sepsis, lipopolysaccharide (LPS) was applied intraperitoneally in gerbils. The animals were anesthetized with a ketamine/xylazine combination before collection; their ABR were collected before any procedure (base collection), after ischemia, and 24 hours after the application of LPS. The absolute latency of wave IV was evaluated, and the values were compared between groups. **Results:** There were significant differences in the groups submitted to sepsis in the latency value of wave IV in relation with the other groups. **Conclusion:** ABR was sensitive to sepsis with the increase in latency of wave IV during the development of the disease in the experimental model used.

**RESUMO**

**Introdução:** O acidente vascular cerebral do tipo isquêmico é um quadro clínico que afeta milhares de pessoas no mundo. Como resultado dessa injúria observamos morte neuronal, e no decorrer natural de recuperação o indivíduo pode desenvolver a sepse. A sepse é uma resposta inflamatória sistêmica que pode levar o indivíduo à morte. Para avaliar as condições clínicas de um paciente com esse quadro, o potencial evocado auditivo de tronco encefálico (PEATE) pode ser útil, uma vez que ele não é um procedimento invasivo, sua técnica é rápida e pode ser feita à beira do leito. **Objetivo:** Verificar os valores de latência do PEATE em gerbils submetidos à isquemia e sepse. **Métodos:** Foram coletados os PEATEs de 72 gerbils machos adultos que foram distribuídos em 6 grupos: controle, sepse, isquemia, sham, isquemia com sepse e sham com sepse. Para a indução de sepse, foi aplicado lipopolissacarídeo (LPS) intraperitoneal nos gerbils. Os animais foram anestesiados antes das coletas com ketamina associada à xilazina, seus PEATEs foram coletados antes de qualquer procedimento (coleta base), após isquemia e 24 horas após a aplicação de LPS. Foi avaliada a latência absoluta da onda IV, e os valores foram comparados intergrupos. **Resultados:** Houve diferença significativa nos grupos submetidos à sepse no valor da latência da onda IV em relação aos demais grupos. **Conclusão:** O PEATE se mostrou sensível à sepse com aumento de latência da onda IV no desenvolver da doença no modelo experimental utilizado.

**Correspondence address:**

Janaina Patricio de Lima

Rua José Benedito Salinas, 68, apartamento 152B, Jardim Itapeva, São Paulo (SP), Brasil, CEP: 04674-200.

E-mail: janapatricio@yahoo.com.br

Received: 09/08/2014

Accepted: 01/11/2015

Study carried out at the Laboratory of Clinical Emergencies, School of Medicine, Universidade de São Paulo – USP – São Paulo (SP), Brazil.

(1) Universidade de São Paulo – USP – São Paulo (SP), Brazil.

**Conflict of interests:** nothing to declare.

## INTRODUCTION

Clinical use of the auditory brainstem response (ABR) in assessing the integrity of auditory pathways and in the investigation of the electrophysiological threshold is already routine<sup>(1,2)</sup>. However, using this potential as a prognostic index in various diseases is still object of study in the field of research. Studies aimed at characterizing the ABR in these situations are of great value, because, for clinical use in situations where the patient lies in bed, as in the case of patients who have suffered a stroke or were victims of sepsis, well-defined parameters are needed. With this approach, experimental studies play an important role, as it is possible to reduce variables in the laboratory, and thus better understand the mechanisms and the results<sup>(3)</sup>.

Ischemic stroke and sepsis are two diseases that affect a large number of patients. Ischemia can be classified as either global or focal, that is, when there is total or partial lack of oxygen delivery; and as either complete or incomplete, when there is no reperfusion, that is, the return of the vascular blood flow<sup>(4,5)</sup>. The lack of oxygen in neurons leads to neuronal death. In addition, during ischemia and reperfusion, there is excessive neuron excitation, a phenomenon known as excitotoxicity, which is caused by excess glutamate release by oxygen-deprived neurons. This increase in glutamate release is responsible for the death of postsynaptic neurons, because this substance in excessive concentration is toxic to neurons<sup>(6,7)</sup>.

Many patients are left with postischemic stroke sequelae. In addition, in the natural course of recovery, the individual may develop sepsis<sup>(8)</sup>, which is a systemic, exacerbated proinflammatory response that has a complex development, which, if not controlled, affects many organs and may even cause death<sup>(9)</sup>.

These patients are often kept in intensive care units, and transporting them for laboratory tests is difficult and risky. In this perspective, the ABR becomes a valuable tool, as it is a noninvasive, fast test and can be performed at the bedside<sup>(10)</sup>, which may indicate some level of prognosis for the patient's condition.

The objective of this study was to investigate the ABR latency values of wave IV in adult gerbils in pre- and postischemia, pre- and postsepsis conditions, compared with a control group and a sham group.

## METHODS

This study was approved by the Ethics Committee for Analysis of Research Projects of the Clinical Board of Hospital das Clínicas and of the School of Medicine of Universidade de São Paulo (Protocol No. 0456/09). The research was conducted at the Laboratory of Medical Research on Clinical Emergencies, Universidade de São Paulo.

A total of 72 male gerbils (*Meriones unguiculatus*), weighing between 56 and 79 g, aged 3 to 5 months, kept in controlled environmental conditions with light/dark cycle (12/12 hours), temperature in the range of 22 to 27°C, humidity of 45 to 65%, and sanitized environment, receiving common gerbil food and water *ad libitum*, from the vivarium of the Department of Clinical Medicine, School of Medicine, Universidade de São Paulo.

The use of these rodents is justified by the fact that they show a failure in the formation of the circle of Willis, which provides a single irrigation for 80% of the brain parenchyma. Thus, the surgical preparation of the experimental model is limited to the temporary carotid occlusion, eliminating unnecessary complications and producing far more consistent results.

The animals were divided into six experimental groups, and each group consisted of two animals, namely:

1. sham (surgery without carotid occlusion);
2. ischemia (surgery with carotid occlusion);
3. sepsis;
4. sham with sepsis;
5. ischemia with sepsis; and
6. control (without any type of procedure).

For the ischemia procedure, the gerbils were anesthetized with a 3% halothane in a gas mixture of 30% oxygen and 70% nitrous oxide and subjected to global cerebral ischemia for 7 minutes with bilateral carotid occlusion. During the ischemic period (7 minutes), followed by the reperfusion period (30 minutes), the animals were monitored and maintained at a controlled temperature of 36.0±0.2°C. Then, the animals were maintained on food and water *ad libitum*.

The ischemia and surgery procedure without occlusion of the carotid (sham group) was performed on the first day of the experiment, after the first ABR collection (base collection).

Regarding the groups with sepsis, during the second day of the experiment, that is, 24 hours after ischemia, they were injected with 1 mg/kg lipopolysaccharide (LPS) from *Escherichia coli* serotype 026: B6 (Sigma) intraperitoneally, thus inducing inflammation (sepsis) in the animal.

For the electrophysiological assessment, the equipment used was the Navigator model by Biologic. The stimuli were presented through 3A insert phones, positioned in the animals' left ear, with their right ear not occluded. The stimulation happened solely on the left ear.

Before all records were made, the animals were anesthetized with a peritoneal ketamine injection (Parke-Davis, São Paulo, Brazil) at a concentration of 100 mg/kg of body weight and Rompun, whose active principle is xylazine (Bayer HealthCare, Sao Paulo, Brazil) at a concentration of 4 mg/kg of body weight. The ABR was recorded by the use of three electrodes positioned in Cz, left mastoid M1, according to the international 10-20 system<sup>(11)</sup>, and a ground electrode. The left mastoid was used as reference, and the animal's paw was used for the ground electrode. The animals remained in a passive listening situation. To capture the ABR, 2,000 click-type stimuli were used with thin polarity and presentation rate of 13 stimuli per second, intensity of 80 dBnHL, and insert earphones coupled to the animal's auricular pavillion. Filters used were 100 Hz pass/low and 1,500 Hz pass/high filters. The recording window was of 10.66 ms after the stimulation. All responses were reproduced, thus ensuring the reproducibility of the acquired potential.

Before any procedure, an electrophysiological evaluation (ABR) was performed, so that this measurement could serve as a baseline for all the others that were performed (pregathering,

held on the first day of the experiment). The ABR was collected again on the second day in all animals after the ischemic procedure performed in those who belonged to the applicable group (collection A); and a third sample was collected in all animals 24 hours after the LPS injection in animals belonging to the infected group (B collection). Only wave IV of ABR was analyzed, owing to a visualization problem with the other waves<sup>(12)</sup>. The professional who examined the waves did not know to which group the animals belonged, neither at which step of the study they were.

The results were compared in pre- and postimplantation of the ischemic event and application of LPS.

For the statistical analysis, ANOVA and Tukey's tests<sup>(13)</sup> were used. Values lower than 0.05 ( $p < 0.05$ ) were considered significant.

## RESULTS

Table 1 shows the mean and standard deviation of the latency values of wave IV of ABR per group at all moments of the study. With the data, we can see a significant difference in the first collection (pregathering) and final collection (B) values (Table 2) in all groups that were induced to sepsis (sepsis, sham with sepsis, and ischemia with sepsis). The opposite is observed in groups in which sepsis was not induced (control, ischemia, and sham).

## DISCUSSION

The data in Tables 1 and 2 and Figure 1 show the values of wave IV of ABR in the different groups studied and their significant difference. One can see the difference between the induced sepsis groups (sepsis, ischemia/sepsis, and sepsis/sham) from those not induced to the inflammatory process (sham, ischemic, and control). On the other hand, in collection B, there was no difference between the ischemic and nonischemic groups at any point of the search. Thus, we have an increase in wave IV latency in the development of sepsis (collection B, 24 hours after the application of LPS), which is not observed in the other groups. This shows that the ABR was able to identify the animal with sepsis but not the animal with ischemia.

Sepsis is a disease that can lead to poor brain function, owing to the increase of cytokines that can cause apoptosis and alterations in synaptic transmission. It is a process that involves local and systemic alterations and is not yet fully understood. It is known, however, that there is an excess inflammation of

the organism, and, through positive feedback and synergistic responses of various systems, the inflammation is potentiated without control. Among the main events observed with excessive production of pro- and anti-inflammatory cytokines, there is the activation of the adhesion molecules, increased arachidonic acid, oxygen-free radicals, nitric oxide, and platelet activating factor. This lack of control of these substance causes damage to vital organs, and on the central nervous system, alterations in cytokine levels in sepsis can impair brain function, resulting in a condition known as septic encephalopathy<sup>(14,15)</sup>.

No studies were found involving ABR and sepsis, but only studies with other evoked potentials. In one study<sup>(16)</sup>, the authors verified the cortical (N20-N70) and subcortical (N13-N20) somatosensory potentials to assess the incidence of septic encephalopathy in patients with severe sepsis and septic shock. The conclusion of the group was that these potentials are sensitive to this type of evaluation, and their loss was associated with disease severity.

The groups subjected only to ischemia did not differ significantly from those without ischemia and without sepsis. The value of absolute latency of wave IV did not change between


**Table 2.** Statistical analysis comparing the latency of wave IV of the auditory brainstem response between the groups evaluated

	Precollection	Collection A	Collection B
Sepsis x Ischemia	NS	NS	0.007
Sepsis x ischemia/sepsis	NS	NS	NS
Sepsis x control	NS	NS	0.005
Sepsis x sham	NS	NS	0.022
Sepsis x sham/sepsis	NS	NS	NS
Ischemia x ischemia/sepsis	NS	NS	0.000
Ischemia x control	NS	NS	NS
Ischemia x sham	NS	NS	NS
Ischemia x sham/sepsis	NS	NS	0.002
Ischemia/sepsis x control	NS	NS	0.000
Ischemia/sepsis x sham	NS	NS	0.000
Ischemia/sepsis x sham/sepsis	NS	NS	NS
Control x sham	NS	NS	NS
Control x sham/sepsis	NS	NS	0.002
Sham x sham/sepsis	NS	NS	0.005

**Caption:** NS = nonsignificant  
Statistical significance ( $p \leq 0.05$ ).

**Table 1.** Descriptive analysis (mean and standard deviation) of latency (ms) of wave IV of the auditory brainstem response in moments of the precollection and collections A and B, according to the evaluated groups

Collection	Control (n=12)	Sepsis (n=12)	Ischemia (n=12)	Sham (n=12)	Ischemia/sepsis (n=12)	Sepsis/sham (n=12)
Precollection	3.03±0.18	3.03±0.20	2.99±0.12	2.99±0.23	3.03±0.18	3.01±0.19
A	3.00±0.13	3.08±0.19	3.04±0.16	2.97±0.15	3.15±0.33	3.07±0.14
B	2.98±0.23	3.99±1.03	3.02±0.21	3.10±0.30	4.42±0.93	4.52±0.61



**Figure 1.** Layout of the average of the potential of the auditory brainstem response in each study group in all the moments (precollection, A, and B)

the precollection and collection B. In this study, the animals underwent carotid occlusion on the first day of the experiment, and the ABR was collected 24 hours after the procedure. The analysis was also performed just for wave IV.

In a study made with cats subjected to ischemia by occlusion of the common carotid, the authors<sup>(17)</sup> found the presence of wave I and absence of the other waves in the ABR. In this study, the ABR was evaluated immediately after the surgical procedure.

In a later study, the authors found the latencies of ABR waves in Wistar rats subjected to occlusion of the anterior cerebellar artery. They divided the study into two groups according to collection: ipsilateral to the occlusion (group A) and contralateral to the occlusion (collection B). The authors observed different types of waves in group A, from total absence of ABR waves to waves I, II, III and IV, with an increase in interpeak latencies I-IV and II-IV. In relation to group B (whose collection was contralateral to the occlusion), the authors observed the presence of waves I, II, III and IV, with an increase in interpeak latencies I-IV and II-IV. All samples were taken within 60 minutes after occlusion. The authors justified the response variability in group A because of possible variations in vessels in the anterior cerebellar artery. According to them, when there is a significant contralateral movement, there may be blood supply to the cochlea, which maintains the presence of ABR waves.

In our study, we believe that the ABR values held before and after ischemia can be justified, because the potential assesses the brainstem region. It is known that ischemic damage are related mainly to the cortical regions (CA1 area of the cortex)<sup>(19)</sup>. In addition, the ABR in this study was collected 24 hours after the ischemia procedure; thus, there may have been injuries during the lack of blood supply, as cited by Inui et al.<sup>(18)</sup>, but the body itself may have recovered from event. In this study, the ABR was not sensitive to ischemia.

## CONCLUSION

This study found a strong correlation between the increases in latency of wave IV of ABR in animals induced to sepsis. Thus, the evaluation of this potential can serve as a prognostic factor for identifying the damage to the brainstem in gerbils with sepsis. However, the same effect was not observed in animals subjected to ischemia, whose wave IV value did not change.

*\*JPL conducted the tests in gerbils and drafted the manuscript; ITV and ES corrected the manuscript; DFB was responsible for the process of inducing ischemia and sepsis in the animal studied.*

## REFERENCES

1. Casali RL, Santos MFC. Potencial evocado auditivo de tronco encefálico: padrão de respostas de lactentes termos e prematuros. *Braz J Otorrinolaryngol.* 2010;76(6):729-38.
2. Pedriali IVG, Kozłowski L. Potenciais evocados auditivos de tronco encefálico na detecção do neurinoma do acústico. *Arquivos Int Otorrinolaryngol.* 2005;9(1):303-6.

3. Seelig VC. Questões atuais relacionadas ao uso de modelos animais em pesquisa científica [monografia]. Porto Alegre: Universidade Federal do Rio Grande do Sul; 2007.
4. Lenzi GL, Frackowiak RS, Jones T. Cerebral oxygen metabolism and blood flow in human cerebral ischemic infarction. *J Cereb Blood Flow Metab.* 1982;2(3):321-35.
5. Joaquim MAS, Patriota GC, Bianco AM. Isquemia encefálica, cascatas vasodilatadoras e alterações bioquímicas. *Arq Bras Neurocir.* 2010;29(2):58-63.
6. Zipfel GJ, Lee JM, Choi DW. Reducing calcium overload in the ischemic brain. *N Engl J Med.* 1999;341(20):1543-4.
7. Stehno-Bittel L. Neuroplasticidade. In: Lundy-Ekman L. *Neurociência: fundamentos para reabilitação.* Rio de Janeiro: Elsevier; 2008. p. 61-70.
8. Basile-Filho A, Suen VMM, Martins MA, Coletto FA, Marson F. Monitorização da resposta orgânica ao trauma e à sepse. *Medicina (Ribeirão Preto).* 2001;34:5-17.
9. Juncal VR, Britto Neto LA, Camelier AA, Messeder OHC, Farias AMC. Impacto clínico do diagnóstico de sepse à admissão em UTI de um hospital privado em Salvador, Bahia. *J Bras Pneumol.* 2011;37(1):85-92.
10. Jardim M, Person OC, Rapoport PB. Potencial evocado auditivo de tronco encefálico como auxílio diagnóstico de morte encefálica. *Pró-Fono R Atual Cient.* 2008;20(2):123-8.
11. Jasper HH. The ten-twenty electrode system of the International Federation. *Electroenceph Clin Neurophysiol.* 1958;10:371-5.
12. Smith DI, Kraus N. Postnatal development of the auditory brainstem response (ABR) in the unanesthetized gerbil. *Hear Res.* 1987;27(2):157-64.
13. Field A. *Descobrimos a estatística usando o SPSS. 2ª edição.* São Paulo: Artmed; 2009.
14. Hotchkiss RS, Tinsley KW, Swanson PE, Karl IE. Endothelial cell apoptosis in sepsis. *Crit Care Med.* 2002;30(5 Suppl):S225-8.
15. Silva FP. Nomenclatura e Epidemiologia. In: Silva FP, Velasco IT. *Sepse.* Barueri: Manole; 2007. p. 12-8.
16. Zauner C, Gendo A, Kramer L, Funk GC, Bauer E, Schenk P, et al. Impaired subcortical and cortical sensory evoked potential pathways in septic patients. *Crit Care Med.* 2002;30(5):1136-9.
17. Sohmer H, Gafni M, Havatselet G. Persistence of auditory nerve response and absence of brain-stem response in severe cerebral ischaemia. *Electroencephalogr Clin Neurophysiol.* 1984;58(1):65-72.
18. Inui H, Murai T, Matsunaga T. Auditory brainstem response findings in brainstem ischemia following selective occlusion of the anterior inferior cerebellar artery in the rat. *Eur Arch Otorhinolaryngol.* 1995;252(3):181-5.
19. O'Sullivan SB, Schmitz TJ. *Fisioterapia, avaliação e tratamento. 2ª ed.* São Paulo: Manole; 1993.