



Erratum

In the article "Fungi on irrigated rice seeds produced in the pre-germinated system in the Alto Vale do Itajaí region, Santa Catarina state, Brazil" published in *Ciência Rural*, volume 50, number 8, DOI <http://doi.org/10.1590/0103-8478cr20190903>.

In the table 1, where we read:

Table 1 - Average incidence of fungi associated with seeds of irrigated rice cultivars produced in the Alto Vale do Itajaí Region, Santa Catarina State. Lages, 2019.

Cultivar	Nº Lotes	Incidence of fungi (%)										
		<i>Mic</i>	<i>Alt</i>	<i>Cur</i>	<i>Bip</i>	<i>Sar</i>	<i>Nig</i>	<i>Pen</i>	<i>Asp</i>	<i>Altsp</i>	<i>Cer</i>	<i>Fus</i>
-----Safra 2015 / 2016-----												
SCS121 CL	57	61.9	6.0	6.6	7.3	2.2	0.8	1.1	0.7	0.1	0.2	0.0
SCS116 Satoru	33	72.3	2.9	1.7	1.6	2.2	1.6	1.3	1.2	0.4	0.3	0.0
SCSBRS Tio Taka	17	55.6	7.7	5.7	7.3	2.7	3.0	0.5	0.7	0.9	0.4	0.4
Epagri 109	11	68.1	2.0	2.0	2.5	3.9	1.2	2.1	2.3	1.0	0.0	0.0
SCS117 CL	11	60.5	9.0	3.9	5.1	2.5	1.4	0.4	0.3	0.5	0.4	0.0
SCS118 Marquês	10	57.2	7.6	9.6	3.3	3.7	0.7	1.8	1.8	0.0	0.2	0.0
Average (%)		62.6	5.8	4.9	4.5	2.8	1.4	1.2	1.2	0.5	0.3	0.1
-----Safra 2016 / 2017-----												
SCS121 CL	55	530	7.4	3.0	2.7	4.2	0.9	0.6	1.0	0.7	1.2	0.1
SCS116 Satoru	33	506	9.1	2.4	2.8	2.7	0.7	1.3	1.9	0.9	0.8	0.2
Epagri 109	31	455	11.3	3.0	3.2	3.9	0.6	0.8	1.3	1.5	0.6	0.3
SCSBRS Tio Taka	26	627	6.5	2.5	3.4	3.4	0.7	0.5	0.7	1.0	0.5	0.3
SCS122 Miura	12	431	11.7	5.6	5.1	7.1	0.0	1.2	2.1	2.3	1.1	0.0
SCS118 Marquês	12	453	9.3	3.0	3.3	4.8	1.6	0.4	0.8	1.6	1.1	0.0
Average (%)		500	9.2	3.2	3.4	4.3	0.7	0.8	1.3	1.3	0.9	0.2
-----Safra 2017 / 2018-----												
SCS122 Miura	73	535	7.2	1.7	1.6	1.1	0.6	0.1	0.0	0.3	1.2	0.0
SCS121 CL	53	516	5.9	0.8	3.3	0.5	0.8	0.1	0.2	0.4	1.0	0.1
Epagri 109	16	446	14.3	1.7	1.9	0.3	0.7	0.1	0.0	0.8	1.2	0.0
SCS116 Satoru	13	489	8.9	1.2	2.6	0.3	0.0	0.2	0.4	0.7	2.5	0.1
Primoriso CL	10	577	7.2	1.0	0.9	0.9	0.5	0.2	0.0	0.4	2.8	0.0
SCSBRS Tio Taka	6	562	4.5	0.7	2.5	2.2	0.3	0.3	0.0	0.3	1.2	0.0
Average (%)		521	8.0	1.2	2.1	0.9	0.5	0.2	0.1	0.5	1.6	0.0
Overall Average (%)		550	7.7	3.1	3.3	2.7	0.9	0.7	0.9	0.8	0.9	0.1

Mic - *Microdochium albescens*; *Alt* - *Alternaria padwickii*; *Cur* - *Curvularia sp.*; *Bip* - *Bipolaris oryzae*; *Sar* - *Sarocladium oryzae*; *Nig* - *Nigrospora oryza*; *Pen* - *Penicillium spp.*; *Asp* - *Aspergillus flavus*; *Alt_{sp}* - *Alternaria sp.*; *Cer* - *Cercospora janseana* e *Fus* - *Fusarium sp.*

Read:

Table 1 - Average incidence of fungi associated with seeds of irrigated rice cultivars produced in the Alto Vale do Itajaí Region, Santa Catarina State. Lages, 2019.

Cultivar	Nº Lotes	Incidence of fungi (%)										
		<i>Mic</i>	<i>Alt</i>	<i>Cur</i>	<i>Bip</i>	<i>Sar</i>	<i>Nig</i>	<i>Pen</i>	<i>Asp</i>	<i>Altsp</i>	<i>Cer</i>	<i>Fus</i>
-----Safra 2015 / 2016-----												
SCS121 CL	57	61.9	6.0	6.6	7.3	2.2	0.8	1.1	0.7	0.1	0.2	0.0
SCS116 Satoru	33	72.3	2.9	1.7	1.6	2.2	1.6	1.3	1.2	0.4	0.3	0.0
SCSBRS Tio Taka	17	55.6	7.7	5.7	7.3	2.7	3.0	0.5	0.7	0.9	0.4	0.4
Epagri 109	11	68.1	2.0	2.0	2.5	3.9	1.2	2.1	2.3	1.0	0.0	0.0
SCS117 CL	11	60.5	9.0	3.9	5.1	2.5	1.4	0.4	0.3	0.5	0.4	0.0
SCS118 Marquês	10	57.2	7.6	9.6	3.3	3.7	0.7	1.8	1.8	0.0	0.2	0.0
Average (%)		62.6	5.8	4.9	4.5	2.8	1.4	1.2	1.2	0.5	0.3	0.1
-----Safra 2016 / 2017-----												
SCS121 CL	55	53.0	7.4	3.0	2.7	4.2	0.9	0.6	1.0	0.7	1.2	0.1
SCS116 Satoru	33	50.6	9.1	2.4	2.8	2.7	0.7	1.3	1.9	0.9	0.8	0.2
Epagri 109	31	45.5	11.3	3.0	3.2	3.9	0.6	0.8	1.3	1.5	0.6	0.3
SCSBRS Tio Taka	26	62.7	6.5	2.5	3.4	3.4	0.7	0.5	0.7	1.0	0.5	0.3
SCS122 Miura	12	43.1	11.7	5.6	5.1	7.1	0.0	1.2	2.1	2.3	1.1	0.0
SCS118 Marquês	12	45.3	9.3	3.0	3.3	4.8	1.6	0.4	0.8	1.6	1.1	0.0
Average (%)		50.0	9.2	3.2	3.4	4.3	0.7	0.8	1.3	1.3	0.9	0.2
-----Safra 2017 / 2018-----												
SCS122 Miura	73	53.5	7.2	1.7	1.6	1.1	0.6	0.1	0.0	0.3	1.2	0.0
SCS121 CL	53	51.6	5.9	0.8	3.3	0.5	0.8	0.1	0.2	0.4	1.0	0.1
Epagri 109	16	44.6	14.3	1.7	1.9	0.3	0.7	0.1	0.0	0.8	1.2	0.0
SCS116 Satoru	13	48.9	8.9	1.2	2.6	0.3	0.0	0.2	0.4	0.7	2.5	0.1
Primoriso CL	10	57.7	7.2	1.0	0.9	0.9	0.5	0.2	0.0	0.4	2.8	0.0
SCSBRS Tio Taka	6	56.2	4.5	0.7	2.5	2.2	0.3	0.3	0.0	0.3	1.2	0.0
Average (%)		52.1	8.0	1.2	2.1	0.9	0.5	0.2	0.1	0.5	1.6	0.0
Overall Average (%)		55.0	7.7	3.1	3.3	2.7	0.9	0.7	0.9	0.8	0.9	0.1

Mic - *Microdochium albescens*; *Alt* - *Alternaria padwickii*; *Cur* - *Curvularia sp.*; *Bip* - *Bipolaris oryzae*; *Sar* - *Sarocladium oryzae*; *Nig* - *Nigrospora oryza*; *Pen* - *Penicillium spp.*; *Asp* - *Aspergillus flavus*; *Alt_{sp}* - *Alternaria sp.*; *Cer* - *Cercospora janseana* e *Fus* - *Fusarium sp.*

In the table 2, where we read:

Table 2 - Average prevalence of fungi associated with seeds of irrigated rice cultivars produced in the Alto Vale do Itajaí Region, Santa Catarina State. Lages, 2019.

Cultivar	Nº Lotes	Prevalence of fungi (%)										
		<i>Mic</i>	<i>Alt</i>	<i>Cur</i>	<i>Bip</i>	<i>Sar</i>	<i>Nig</i>	<i>Pen</i>	<i>Asp</i>	<i>Altsp</i>	<i>Cer</i>	<i>Fus</i>
-----Safra 2015 / 2016-----												
SCS121 CL	57	100	84.7	66.7	39.4	63.2	36.4	33.3	48.5	21.2	9.1	0.0
SCS116 Satoru	33	100	100	70.6	76.5	57.6	64.7	29.4	23.5	17.7	17.7	0.0
SCSBRS Tio Taka	17	100	100	90.9	45.5	64.7	54.6	36.4	9.1	18.2	18.2	0.4
Epagri 109	11	100	36.4	45.5	54.6	81.8	27.3	54.6	63.6	27.3	9.1	0.0
SCS117 CL	11	100	89.5	89.5	77.2	72.7	26.3	45.6	22.8	8.8	14.0	0.0
SCS118 Marquês	10	100	90.0	100	50.0	70.0	40.0	50.0	40.0	0.0	100	0.0
Average (%)		100	83.4	77.2	57.2	68.3	41.6	41.6	34.6	15.5	13.0	0.1
-----Safra 2016 / 2017-----												
SCS121 CL	55	100	89.1	80.0	63.6	63.6	23.6	27.3	36.4	32.7	40.0	3.6
SCS116 Satoru	33	100	90.9	69.7	60.6	45.5	21.2	48.5	42.4	36.4	21.2	6.1
Epagri 109	31	100	92.3	69.2	53.8	38.5	19.2	21.9	19.2	46.2	34.6	7.7
SCSBRS Tio Taka	26	100	96.8	77.4	67.7	58.1	16.1	32.3	45.2	54.8	32.3	9.7
SCS122 Miura	12	100	100	91.7	100	91.7	0.0	58.3	50.0	75.0	33.3	16.7

SCS118 Marquês	12	100	100	91.7	75.0	50.0	50.0	16.2	25.0	50.0	25.0	16.7
Average (%)		100	94.8	79.9	70.1	57.9	21.7	34.1	36.4	49.2	31.1	10.1
-----Safra 2017 / 2018-----												
SCS122 Miura	73	100	868	26.4	66.0	18.9	22.6	5.7	7.6	18.9	32.1	5.7
SCS121 CL	53	100	100	53.1	69.2	15.4	23.1	15.4	15.4	30.8	53.8	0.0
Epagri 109	16	100	100	16.6	50.0	0.0	16.7	16.7	0.0	16.7	50.0	0.0
SCS116 Satoru	13	100	937	50.0	56.3	12.5	18.8	6.3	0.0	31.3	50.0	0.0
Primoriso CL	10	100	918	57.5	45.2	34.3	12.3	6.9	1.4	10.9	49.3	2.7
SCSBRS Tio Taka	6	100	90.0	50.0	30.0	50.0	0.0	10.0	0.0	20.0	90.0	0.0
Average (%)		100	93.7	42.3	52.8	21.9	15.6	10.2	4.1	21.4	54.2	1.4
Overall Average (%)		100	90.6	66.5	59.9	49.4	26.3	28.6	25.0	28.7	32.8	11.6

Mic - *Microdochium albescens*; Alt - *Alternaria padwiicki*; Cur - *Curvularia sp.*; Bip - *Bipolaris oryzae*; Sar - *Sarocladium oryzae*; Nig - *Nigrospora oryza*; Pen - *Penicillium spp.*; Asp - *Aspergillus flavus*; Alt_{sp} - *Alternaria sp.*; Cer - *Cercospora janseana* e Fus - *Fusarium sp.*

Read:

Table 2 - Average prevalence of fungi associated with seeds of irrigated rice cultivars produced in the Alto Vale do Itajaí Region, Santa Catarina State. Lages, 2019.

Cultivar	N° Lotes	-----Prevalence of fungi (%)-----										
		Mic	Alt	Cur	Bip	Sar	Nig	Pen	Asp	Altsp	Cer	Fus
-----Safra 2015 / 2016-----												
SCS121 CL	57	100	84.7	66.7	39.4	63.2	36.4	33.3	48.5	21.2	9.1	0.0
SCS116 Satoru	33	100	100	70.6	76.5	57.6	64.7	29.4	23.5	17.7	17.7	0.0
SCSBRS Tio Taka	17	100	100	90.9	45.5	64.7	54.6	36.4	9.1	18.2	18.2	0.4
Epagri 109	11	100	36.4	45.5	54.6	81.8	27.3	54.6	63.6	27.3	9.1	0.0
SCS117 CL	11	100	89.5	89.5	77.2	72.7	26.3	45.6	22.8	8.8	14.0	0.0
SCS118 Marquês	10	100	90.0	100	50.0	70.0	40.0	50.0	40.0	0.0	100	0.0
Average (%)		100	83.4	77.2	57.2	68.3	41.6	41.6	34.6	15.5	13.0	0.1
-----Safra 2016 / 2017-----												
SCS121 CL	55	100	89.1	80.0	63.6	63.6	23.6	27.3	36.4	32.7	40.0	3.6
SCS116 Satoru	33	100	90.9	69.7	60.6	45.5	21.2	48.5	42.4	36.4	21.2	6.1
Epagri 109	31	100	92.3	69.2	53.8	38.5	19.2	21.9	19.2	46.2	34.6	7.7
SCSBRS Tio Taka	26	100	96.8	77.4	67.7	58.1	16.1	32.3	45.2	54.8	32.3	9.7
SCS122 Miura	12	100	100	91.7	100	91.7	0.0	58.3	50.0	75.0	33.3	16.7
SCS118 Marquês	12	100	100	91.7	75.0	50.0	50.0	16.2	25.0	50.0	25.0	16.7
Average (%)		100	94.8	79.9	70.1	57.9	21.7	34.1	36.4	49.2	31.1	10.1
-----Safra 2017 / 2018-----												
SCS122 Miura	73	100	86.8	26.4	66.0	18.9	22.6	5.7	7.6	18.9	32.1	5.7
SCS121 CL	53	100	100	53.1	69.2	15.4	23.1	15.4	15.4	30.8	53.8	0.0
Epagri 109	16	100	100	16.6	50.0	0.0	16.7	16.7	0.0	16.7	50.0	0.0
SCS116 Satoru	13	100	93.7	50.0	56.3	12.5	18.8	6.3	0.0	31.3	50.0	0.0
Primoriso CL	10	100	91.8	57.5	45.2	34.3	12.3	6.9	1.4	10.9	49.3	2.7
SCSBRS Tio Taka	6	100	90.0	50.0	30.0	50.0	0.0	10.0	0.0	20.0	90.0	0.0
Average (%)		100	93.7	42.3	52.8	21.9	15.6	10.2	4.1	21.4	54.2	1.4
Overall Average (%)		100	90.6	66.5	59.9	49.4	26.3	28.6	25.0	28.7	32.8	11.6

Mic - *Microdochium albescens*; Alt - *Alternaria padwiicki*; Cur - *Curvularia sp.*; Bip - *Bipolaris oryzae*; Sar - *Sarocladium oryzae*; Nig - *Nigrospora oryza*; Pen - *Penicillium spp.*; Asp - *Aspergillus flavus*; Alt_{sp} - *Alternaria sp.*; Cer - *Cercospora janseana* e Fus - *Fusarium sp.*