


# Revealing of free radical scavenging and angiotensin I-converting enzyme inhibitor potency of pigmented rice seed protein

Erlin SUSILOWATI<sup>1</sup>, Bella Rhea Lavifa SANJAYA<sup>2</sup>, Ari Satia NUGRAHA<sup>1,2,3</sup>,  
Mohammad UBAIDILLAH<sup>1,2,4</sup>, Tri Agus SISWOYO<sup>1,2,4\*</sup> 

## Abstract

The potential source of bioactive protein from pigmented rice seed was investigated by buffered extracting from nine varieties (Merah-SP, Gogo Niti-2, Merah Wangi, Super Manggis, Lamongan-1, Mota, Ketan Hitam-2, Beureum Taleus, and Aek Sibundong) and non-pigmented rice (IR-64) as a control. The potent contributor to free radical scavenging of extracted proteins was evaluated by analyzing their free amino acid composition. The free radical scavenging and angiotensin-I converting enzyme (ACE-I) inhibitory activity of these proteins were analyzed *in-vitro*. The free radical scavenging activity was analyzed using various standard methods, including radical cation 2,2-azino-bis(3-ethylbenzothiazoline-6-sulphonate acid (ABTS<sup>+</sup>), hydroxyl radical (OH<sup>•</sup>) and defense against radical-mediated DNA damages by hydroxyl. Total free amino acid content ranged between 77.7-181.2 g/100 g protein, and among them, Ketan Hitam-2 displayed the highest abundant antioxidant amino acid grouping (23.23%) and shown significantly higher ABTS<sup>+</sup> activity (IC<sub>50</sub>=8.64 µg/mL) and OH<sup>•</sup> activity (IC<sub>50</sub>=20.33 µg/mL). Moreover, Ketan Hitam-2 also exhibited notable ACE-I inhibitory activity (IC<sub>50</sub>=6.20 µg/mL) and protected hydroxyl-induced oxidative damage to DNA. The *in-vitro* systems for free radical scavenging and ACE-I inhibitory were used to acquire the data. The potency of Ketan Hitam-2 seed protein could be utilized as a natural nutraceuticals compound.

**Keywords:** ACE-I inhibitory; free radical scavenging; pigmented rice; protein.

**Practical Application:** The pigmented rice seed proteins have a high potential for a human nutraceutical health supplement.

## 1 Introduction

Rice (*Oryza sativa* L., family: *Poaceae*) is an important cereal crop in the global area. Rice is mainly used as a major staple globally, including in China (143.790 metric tons), India (100.000 metric tons), and Indonesia (38.100 metric tons), according to Global Rice Consumption 2018/2019 (Shahbandeh, 2019). In the Indonesian Center for Rice Research, the germplasm of rice consists of 2095 accession of local rice, 804 accessions of introduction rice, and 270 varieties of superior rice. However, information regarding pigmented rice varieties are inadequate. Farmers identified the name briefly based on the colors; black rice, red rice, and black glutinous rice.

Pigmented rice naturally contains higher anthocyanin, protein, and polyphenols than non-pigmented rice (Sati & Singh, 2019). Previous studies have evaluated physicochemical properties (Murdifin et al., 2015), Nuclear Magnetic Resonance (NMR) based on metabolomes information (Wijaya et al., 2018), and morphological variation (Shinta et al., 2014) of pigmented rice in Indonesia. However, further explorations of pigmented rice in Indonesia are necessary to reveal its health benefits for reducing the risk of human disease. The presence of bioactive

compounds in pigmented rice can act be as a scavenger to free radicals caused by oxidative stress leading to cell damage.

The usage of naturally sourced bioactive compounds as an alternative drug is in demand to reduce the dependency on synthetic chemical compounds (Jamshidi-Kia et al., 2018). A plant can produce bioactive peptides as nutrient supplements which have health benefits as an antioxidant, anti-inflammatory, and antihypertensive agents (Zou et al., 2020). Pigmented rice is well known in possessing high antioxidant capacity in which it can effectively prevent the unnecessary inflammatory response. The previous report on Lingkang taker and Umling ame of Indian pigmented rice varieties showed antioxidant constituents such as gallic acid, salicylic acid, quercetin, ferulic acid, caffeic acid, apigenin, and anthocyanin, which was proposed to possess a potency in preventing radical-induced related diseases, including hypertension (Samyot et al., 2016).

Bioactive peptides as antihypertensive agents generally composed of short amino acids structure, which is described by the presence of hydrophobic, aromatic, positive charged, and aliphatic amino acids (Rai et al., 2017). Also, the presence of aromatic amino acids in the C-terminus, positively charged amino acids in the middle,

Received 16 Dec., 2020

Accepted 14 June, 2021

<sup>1</sup>Graduate School of Biotechnology, University of Jember, Jember, Indonesia

<sup>2</sup>The Center of Excellence on Crop Industrial Biotechnology – PUI-PT BioTIn, University of Jember, Jember, Indonesia

<sup>3</sup>Drug Utilisation and Discovery Research Group, Faculty of Pharmacy, University of Jember, Jember, Indonesia

<sup>4</sup>Faculty of Agriculture, University of Jember, Jember, Indonesia

\*Corresponding author: [triagus.faperta@unej.ac.id](mailto:triagus.faperta@unej.ac.id), [siswoyo.triagus@gmail.com](mailto:siswoyo.triagus@gmail.com)

and hydrophobic amino acids in the N-terminus of the peptides demonstrated high potential as an ACE inhibitor (Wijesekara et al., 2011). According to (Ahn et al., 2014), a bioactive peptide composed of more than 60% of hydrophobic amino acid could increase the antioxidant activity by transferring electrons from the benzene ring to the radical electrons. However, the anti-oxidative effects of bioactive compounds in local Indonesian pigmented rice remain unclear. Therefore, this study aimed to investigate the antioxidant activity of bioactive peptide of pigmented rice and its role as ACE-inhibitors, which attributed to preventing or delaying hypertension progression.

## 2 Materials and methods

### 2.1 Plants collection and reagents

The nine varieties of pigmented rice seeds (Merah-SP, Gogo Niti-2, Merah Wangi, Super Manggis, Lamongan-1, Mota, Ketan Hitam-2, Beureum Taleus, Aek Sibundong) and non-pigmented rice seeds (IR-64) as control were cultivated with conventional plantation measure and their seeds were harvested on August 2019 at the Center of Excellence on Crop Industrial Biotechnology (PUI-PT BioTIn) in Agrotechno Park Research Area, University of Jember, Jember, East Java, Indonesia. The reagents used in this study are: bovine serum albumin (BSA), hydrogen peroxide ( $H_2O_2$ ), 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulphonic acid ( $ABTS^{++}$ ), pyrogallol, thiobarbituric acid (TBA), trichloroacetic acid (TCA), 2-deoxy-D-ribose, ACE Kit (Dojindo Molecular Technology), glutathione (G-SH), and ethylenediaminetetraacetic acid (EDTA) were purchased from Sigma-Aldrich, Singapore and other supporting chemicals were analytical grade product from Merck Co., USA.

### 2.2 Amino acid compositions analysis

Amino acid hydrolysis was analyzed by the method of (Szkudzińska et al., 2017). The rice samples were weighing (0.1 g), and HCl (6N, 5 mL) was added. The hydrolysis process is conducted at 110 °C for 22 h. The mixture was cooled at room temperature after the hydrolysis, then transferred to a measuring flask (50 mL) with aquabidest added up to boundary marker and then filtered using a filter of 0.45 micron. Into the supernatant, alpha aminobutyric acid (50 mM, 0.4 mL) was added as an internal testing standard. A portion of hydrolysate (20 µL) was injected into a thermostat, autosampler, high-pressure binary pump and photodiode array detector (PDA) in the UPLC system (Waters 2475, US). The chromatographic separation was achieved using the AccQ-Tag Ultra C-18 column (2.1 x 100 mm; 1.7 µm). The column heater was set at 55 °C, and the mobile phase flow rate was maintained at 0.7 mL/min. Eluent A was 10% AccQ-Tag Ultra concentrate solvent A, and eluent B was 100% AccQ-Tag Ultra solvent B. The non-linear separation gradient was 0-0.54 min (99.9% A), 5.74 min (90.0% A), 7.74 min (78.8% A), 8.04-8.64 min (40.4% A), 8.73-10 min (99.9% A). A VanGuard™ Waters column (2.1 mm i.d. x 5 mm, 1.7 µm particles) was used as the guard column. One microliter of the sample was injected for analysis. The PDA detector was set at 260 nm to classify the composition of the amino acids, with a sampling rate of 20 points/sec.

### 2.3 Protein extraction

Rice protein extraction was performed by grounding and homogenizing fresh seed (1 g) with a phosphate buffer (3 mL, 50 mM, pH 6.8). The mixture then centrifugated at 10.000 rpm for 15 min, in which its supernatant was then preserved under liquid nitrogen. The total protein content of the supernatant was measured using the Bradford method (Bonjoch & Tamayo, 2001). Into a portion of the sample (5 µL), aquadest (45 µL) was added prior to Bradford solution addition (950 µL). In order to determine the dissolved protein content, absorbance was recorded at a wavelength of 595 nm, and the result was compared with the BSA standard.

### 2.4 $ABTS^{++}$ radical scavenging activity assay

The free radical scavenging activity was performed based on the  $ABTS^{++}$  activity, using the method Karami et al., (2019). The radical cations were prepared by mixing  $ABTS$  solution (7 mM, 1 mL) with potassium persulfate (2.45 mM, 1 mL) followed by incubation for 12-16 h in the dark place. The  $ABTS$  working solution was diluted with phosphate saline buffer (0.2 M, pH 7) to produce an absorbance of 0.700-0.750 at the wavelength of 734 nm. The photometric assay was conducted on  $ABTS$  solution (950 µL) and protein extracts (20 µg), in which the absorbance was recorded at a wavelength of 734 nm. The antioxidant activity of the tested samples was calculated by using the following Formula 1:

$$(\%) ABTS^{++} : \left[ \frac{(Ac - As)}{Ac} \right] \times 100\% \quad (1)$$

where is Ac = absorbance control, and As = absorbance sample.

### 2.5 Hydroxyl radical scavenging activity assay

Hydroxyl radical scavenging activity was evaluated through the analytical protocol described by (Hazra et al., 2008). Hydroxyl radical was generated by mixing 2-deoxyribose (28 mM, 50 µL),  $FeCl_3$  (10 mM, 10 µL), EDTA (1 mM, 100 µL),  $H_2O_2$  (1 mM, 10 µL), ascorbic acid (1 mM, 100 µL), and protein sample (20 µg). The mixture was then diluted until the final volume of 1 mL with a phosphate buffer (pH 7.4) and was incubated at 37 °C for 1 h. Into the mixture, TBA (1%, 500 µL) and TCA (2.8%, 500 µL) were added. The mixture was incubated for 30 min at 80 °C, and its absorbance at a wavelength of 532 nm was recorded. Percentage inhibition was assessed by comparing the sample absorbance with blank solution sample by using the following Formula 2:

$$(\%) Hydroxyl : \left[ \frac{(Ab - As)}{Ab} \right] \times 100\% \quad (2)$$

where is Ab = absorbance blank and As = absorbance sample.

### 2.6 Angiotensin I-Converting Enzyme (ACE-I) inhibitor activity assay

The activity of ACE-I inhibitor was performed using the ACE Kit (Dojindo Molecular Technology). Crude protein sample (20 µg) was loaded into a sample well followed by the

addition of substrat buffer and enzyme working solution. The mixture was than incubated for 1 h at 37 °C, which was then an indicator working solution (200 µL) applied. The mixture was then re-incubated for 10 min at room temperature, followed by recording the absorbance at wavelength of 450 nm under a microplate reader system (ZN-320, Zenix). Blank 1 (positive control) was made without a sample solution, and blank 2 was made without a sample solution and enzyme working solution. ACE inhibition was evaluated by the following Formula 3:

$$\left[ \frac{(Ab_1 - As)}{(Ab_1 - Ab_2)} \right] \times 100\% \quad (3)$$

where is  $Ab_1$  = absorbance blank 1,  $Ab_2$  = Absorbance blank 2, and  $As$  = absorbance sample and the result was compared with the captopril as reference drug.

### 2.7 Protective DNA assay

Protective DNA was analyzed using the method described by (Siswoyo et al., 2011). The pBT7 plasmid from the collection of Nutraceutical and Pharmaceutical Laboratory, Center for Development of Advanced Science and Technology, University of Jember, Jember, East Java, Indonesia was used. The DNA plasmid (2.5 µg) was treated by a Fenton's reagent (80 mM  $FeCl_3$ , 30 mM  $H_2O_2$ , and 50 mM ascorbic acid) until the final volume of 1 mL and was incubated at 37 °C for 15 min. The protein samples were then added and made the final volume up to 20 µL with ddH<sub>2</sub>O. The mixture was then incubated at 37 °C for 15 min. The mixture then running to 1.5% gel electrophoresis and was visualized through the gel documentation system (Major Science, USA).

### 2.8 Statistical analysis

All values are expressed as the mean of three replicates ± standard deviations (SD). Each data was analyzed statistically

using analysis of variance (ANOVA). Duncan's multiple range tests with a significance level of  $p < 0.05$  were followed. The SPSS® 20.0 Package (IBM Cooperation, Chicago, USA) was used throughout.

## 3 Results


### 3.1 Variation shapes and color morphology of rice seed

The rice seeds of each variety were hulled to determine the variation of shape and color morphologies. The results showed that rice seed of each variety has variation in the shape and color. Visually, the shape could be determined as slender, medium, and round. Whereas, the color of the seeds varies from brown to black (Figure 1).

### 3.2 Variation and ratio of antioxidant amino acid to total amino acid

The seventeen amino acid compositions of each rice were analyzed by UPLC. The data were analyzed by the coefficient variance (CV) value. The CV value is used to determine the percentage of data variability, and the high CV values indicate a wide variety of data. The results showed that the CV value of each amino acid compositions of nine rice is between 19.91-59.52% (Table 1). It was indicated that there is a high variation of amino acid concentrations among rice varieties.

It was known that arginine, lysine, tyrosine, histidine, cysteine, and methionine are the amino acids that correlate with antioxidant activity. The ratio of antioxidant amino acid group to a total amino acid of each variety was analyzed. The results showed the highest ratio of antioxidant amino acid to total amino acids found in Ketan Hitam-2, followed by Beureum Taleus, 23.23%, and 23.17% (Table 1). It was indicated that Ketan Hitam-2 and Beureum Taleus rice seed has a high potency of antioxidant.



**Figure 1.** Variation shape and color morphology of pigmented (A-I) and non-pigmented (J) rice seeds. (A) Merah-SP; (B) Gogo Niti-2; (C) Merah Wangi; (D) Super Manggis; (E) Lamongan; (F) Mota; (G) Ketan Hitam-2; (H) Beureum Taleus; (I) Aek Sibundong; (J) IR-64.



**Table 1.** The amino acid composition of rice seed protein.

Amino acid*	Variety										Mean	SE	CV (%)
	A	B	C	D	E	F	G	H	I	J			
L-Arginine**	9.34	14.22	9.72	12.61	11.76	16.60	10.36	9.96	7.48	10.56	11.26	0.83	23.44
L-Aspartic acid	7.56	10.96	8.61	10.24	9.35	15.72	7.73	7.89	5.90	8.56	9.25	0.85	29.02
L-Cysteine**	0.40	0.67	0.49	0.63	0.53	0.80	0.65	0.55	0.40	0.57	0.57	0.04	21.85
L-Glutamic acid	16.18	24.43	19.02	23.02	20.71	33.32	16.38	16.89	12.40	18.58	20.09	1.84	28.96
L-Histidine**	2.89	4.57	3.38	4.10	3.52	5.20	3.23	3.47	2.58	3.60	3.65	0.25	21.34
L-Lysine**	3.75	4.55	5.30	5.32	4.33	6.71	3.77	3.95	2.86	4.64	4.52	0.34	23.71
L-Serine	6.37	9.61	7.55	8.82	7.58	10.45	6.45	7.26	5.41	7.72	7.72	0.49	19.91
L-Threonine	4.19	6.73	5.07	5.96	5.81	7.40	4.64	4.71	3.64	5.30	5.35	0.36	21.56
L-Alanine	5.53	8.23	6.61	7.70	6.72	10.33	5.76	5.82	4.29	6.42	6.74	0.53	24.93
L-Phenylalanine	5.76	9.64	6.13	8.90	7.90	12.40	6.62	6.84	5.13	7.10	7.64	0.69	28.40
L-Tyrosine**	4.23	6.73	4.06	6.62	5.77	8.42	4.98	5.70	3.51	4.75	5.48	0.47	27.14
Glycine	5.45	8.22	6.29	7.34	6.47	8.90	5.68	5.79	4.42	6.42	6.50	0.42	20.57
L-Isoleucine	4.59	6.89	5.16	6.46	5.73	8.07	4.67	4.64	3.52	5.19	5.49	0.42	24.20
L-Leucine	9.19	14.19	10.36	13.20	11.91	16.55	9.44	9.19	7.08	10.47	11.16	0.89	25.25
L-Methionine**	0.01	0.86	0.78	0.51	0.68	0.70	0.96	0.62	0.37	0.02	0.55	0.10	59.52
L-Valine	6.68	10.07	7.63	9.22	8.41	11.52	6.97	6.60	5.09	7.57	7.98	0.60	23.60
L-Proline	4.69	6.96	5.42	6.44	5.72	8.13	4.80	4.80	3.62	5.39	5.60	0.41	23.12
TAA (g/100 g Protein)	96.81	147.53	111.58	137.09	122.9	181.22	103.09	104.68	77.7	112.86	119.55	9.29	24.58
TAAAnt (g/100 g Protein)	20.62	31.60	23.73	29.79	26.59	38.43	23.95	24.25	17.20	24.14	26.03	1.89	23.01
TAAAnt /TAA (%)	21.30	21.42	21.27	21.73	21.64	21.21	23.23	23.17	22.14	21.39	21.85	0.24	3.49

A: Merah-SP; B: Gogo Niti-2; C: Merah Wangi; D: Super Manggis; E: Lamongan-1; F: Mota; G: Ketan Hitam-2; H: Beureum Taleus; I: Aek Sibundong; J: IR-64; SE: Standard Error; CV: Coefficient of Variance; TAA: Total Amino Acid, TAAAnt: Total Amino Acid Antioxidant.

\*g/100 g protein; \*\*Antioxidant amino acid grouping.

### 3.3 Free radical scavenging activity of pigmented rice seed protein

Free radical scavenging activity was analyzed by ABTS<sup>•+</sup> and OH<sup>•</sup> assay. The results showed that the IC<sub>50</sub> value of all pigmented rice seed protein was significantly lower than non-pigmented rice (control). The lowest IC<sub>50</sub> value of ABTS<sup>•+</sup> scavenging activity was shown in the seed protein of Ketan Hitam-2 (8.64 µg/mL), while the IC<sub>50</sub> value of G-SH as positive control is 0.22 µg/mL (Table 2). It was indicated that the seed protein of Ketan Hitam-2 has the capability to scavenge radical.

The IC<sub>50</sub> value of ABTS<sup>•+</sup> and OH<sup>•</sup> scavenging of seed protein of Ketan Hitam-2 variant is significantly less than other pigmented rices. The IC<sub>50</sub> rate of OH<sup>•</sup> scavenging of the seed protein of Aek Sibundong variant has the same significance as the seed protein of Ketan Hitam-2 variant. The low IC<sub>50</sub> value of ABTS<sup>•+</sup> and OH<sup>•</sup> scavenging in seed protein of Ketan Hitam-2 and Aek Sibundong indicates a potential antioxidant agent as cellular protection against oxidative stress (Table 2). Also, the oxidative DNA damage protection against hydroxyl radicals was evaluated by a fenton's reagent. Results suggested that the pBT7 plasmid incubation of in fenton's reagent for 15 min resulted in supercoiled (SC) cleavage to create the open circular (OC) form (Figure 2), indicating that single-strand DNA breaks were created by the hydroxyl radical formed by the fenton reaction. As shown in Figure 2, extra protein does not always respond to DNA damage. However, the addition of Ketan Hitam-2 seed protein may decrease SC to OC conversion. Its ability is almost the same as shown by G-SH as the positive control.


**Table 2.** The free radical scavenging activities of pigmented rice seed protein.

Variety	Free Radical (*IC <sub>50</sub> , µg/mL)	
	ABTS <sup>•+</sup> (µg/mL)	Hydroxyl (µg/mL)
Merah-SP	45.66 ± 0.93 <sup>c</sup>	25.30 ± 0.53 <sup>d</sup>
Gogo Niti-2	47.34 ± 1.15 <sup>c</sup>	33.08 ± 0.26 <sup>a</sup>
Merah Wangi	44.63 ± 1.09 <sup>c</sup>	23.43 ± 0.15 <sup>f</sup>
Super Manggis	43.99 ± 0.24 <sup>c</sup>	22.11 ± 0.17 <sup>g</sup>
Lamongan-1	61.62 ± 1.02 <sup>b</sup>	24.44 ± 0.15 <sup>e</sup>
Mota	43.61 ± 0.34 <sup>c</sup>	30.30 ± 0.19 <sup>b</sup>
Ketan Hitam-2	8.64 ± 0.68 <sup>e</sup>	20.33 ± 0.23 <sup>i</sup>
Beureum Taleus	36.87 ± 0.35 <sup>d</sup>	20.90 ± 0.18 <sup>h</sup>
Aek Sibundong	63.66 ± 1.94 <sup>b</sup>	20.35 ± 0.17 <sup>i</sup>
IR-64	72.53 ± 1.67 <sup>a</sup>	29.47 ± 0.23 <sup>c</sup>
G-SH	0.22 ± 0.08 <sup>f</sup>	8.24 ± 0.12 <sup>j</sup>

\*IC<sub>50</sub> value (µg/mL) is the effective concentration of antioxidants to inhibit 50% radical ABTS and hydroxyl. Glutathione (G-SH) was used as a positive control. All values are expressed as the mean of three replicates ± standard deviations (SD), the Duncan's multiple range tests with a significance level of *p*<0.05 were followed. The different letters in the column are represented significantly different.

### 3.4 Angiotensin I-Converting Enzyme (ACE-I) inhibitory activity

This study also investigated each pigmented rice seed protein's potency to inhibited the activity of ACE-I. The crude protein of each variety has been analyzed *in-vitro* using the ACE kit. The IC<sub>50</sub> value of the Ketan Hitam-2 seed protein is significantly less than other seed proteins of pigmented rice



**Figure 2.** Inhibitory effect of pigmented rice seed protein on DNA damage treat by hydroxyl radicals. DNA damage was introduced by mixing the fenton reagent into the plasmid DNA for 0 and 15 minutes (lanes 1 and 2, respectively). Seed protein of Merah SP, Gogo Niti 2, Merah Wangi, Super Manggis, Lamongan, Mota, Ketan Hitam-2, Beureum Taleus, Aek Sibundong, and IR-64 (lanes A-J respectively) and Gluthatione as a positive control (lane +). OC: Open Circular; SC: Supercoiled.

**Table 3.** The ACE-I inhibitor activities of pigmented rice seed protein.

Variety	ACE-I inhibitor (*IC <sub>50</sub> , µg/mL)
Merah-SP	15.75 ± 0.04 <sup>f</sup>
Gogo Niti-2	7.66 ± 0.63 <sup>e</sup>
Merah Wangi	6.39 ± 0.06 <sup>c</sup>
Super Manggis	6.76 ± 0.05 <sup>d</sup>
Lamongan-1	6.31 ± 0.02 <sup>c</sup>
Mota	7.05 ± 0.02 <sup>c</sup>
Ketan Hitam-2	6.20 ± 0.05 <sup>b</sup>
Beureum Taleus	17.30 ± 0.11 <sup>g</sup>
Aek Sibundong	26.60 ± 0.14 <sup>i</sup>
IR-64	21.63 ± 0.02 <sup>h</sup>
Captopril	2.08 ± 0.20 <sup>a</sup>

\*IC<sub>50</sub> value (µg/mL) is the effective concentration of antioxidants to inhibit 50% ACE-I activity. Captopril was used as a positive control of ACE-I. All values are expressed as the mean of three replicates ± standard deviations (SD), the Duncan's multiple range tests with a significance level of  $p < 0.05$  were followed. The different letters in the column are represented significantly different.

(Table 3). This result demonstrated that the seed protein of Ketan Hitam-2 has the highest capability to inhibit the ACE-I enzyme than other pigmented rice, therefore this variant was selected as an antihypertensive compound. Nevertheless, the activity still three folds higher than the control (IC<sub>50</sub> value of captopril is 2.08 µg/mL).

#### 4 Discussion

Previous reports showed rice to contain relatively low amounts of proteins within the range of 4-10% (Balindong et al., 2018). This protein content considerably varies among different varieties and cultivar depending on climatic, adaptive, and growth conditions as well as a level of maturity at harvest (Dabi & Khanna, 2018). Compared to non-pigmented rice, pigmented rice varieties tendentially have higher protein content with a well-balanced composition of amino acids (Samyori et al., 2017). The amount of amino acid in each variety of pigmented rice is very different in which CV value are ranging from 19.91% to 59.52% (Table 1) and could represent different potencies. The amino acid profiling in this current study (Table 1) showed higher abundant of L-tyrosine and L-methionine amino acids in pigmented rice. Previous research reported tyrosine and methionine have positive effects on antioxidant capacity due to its structures. Tyrosine have capability to act as hydrogen transferor, while methionine tends to oxidize methionine sulphoxide (Karami et al., 2019). In addition, the amino acid component was previously reported to have high correlation with radical scavenging capacity (Pérez et al., 2007). In the current study, high of L-methionine reported in

pigmented rice protein could play a role as an antioxidant. According to Xu et al. (2017), there are several amino acid which were classified as antioxidant group, these includes L-arginin, L-lysine, L-tyrosine, L-histidine, L-cysteine, and L-methione. As a result, the abundances of TAAAnt/TAA in pigmented rice could contribute to the antioxidant peptides activity.

The scavenging of ABTS<sup>•+</sup> and OH<sup>•</sup> radicals were carried out to confirm antioxidant potential in each rice seed protein. The results showed that the IC<sub>50</sub> value in ABTS<sup>•+</sup> radical and OH<sup>•</sup> radical of Ketan Hitam-2 is the lowest; it is mean that Ketan Hitam-2 protein has the best antioxidant capabilities (Table 2). Several previous studies documented a good association between the percentage of hydrophobic amino acid and the OH<sup>•</sup> radical activity (Ajibola et al., 2011; Zou et al., 2016). Based on data shown in Table 1, the variety with high percentage of hydrophobic amino acid are Gogo Niti-2, Super Manggis and Ketan Hitam-2 with percentage value of 81.37%, 80.47% and 80.35%, respectively. This suggested the high hydroxyl scavenger activity in Ketan Hitam-2 seed protein was related to its high content of hydrophobic amino acids. The mechanism of action may be that antioxidant peptides can enter smoothly through hydrophobic interactions with membrane lipid bilayers by their hydrophobicity, where they can exercise significant scavenging capacity (Pouzo et al., 2016). Bioactive peptides with antioxidant IC<sub>50</sub> values under 500 µg/mL suggested to possess a significant potency as nutraceutical resources (Supriyadi et al., 2019; Xu et al., 2017).

Studies have shown the presence of natural antioxidants as nutraceutical ingredients was able to reduce chronic diseases such as mutagenesis, cancer, and DNA damage (Gupta et al., 2013; Tan et al., 2018). Evaluation of the protective role of rice protein against hydroxyl radicals caused by oxidative DNA damage was performed which in the case of plasmid, the OC DNA conformation resulted in a single-strand break in hydroxyl radical-exposed SC plasmid DNA due to the Fenton reaction. The study indicated incubation of pBT7 in 15 min Fenton's reagent resulted in SC cleavage and created OC form (Figure 2, lane 2). This suggested that the Fenton reaction-formed hydroxyl radical which induced DNA damage in single-strand. Nevertheless, adding of Ketan Hitam-2 seed protein to the Fenton reagent mixture clearly reduced the SC converting to OC DNA significantly (Figure 2). This can be caused by the high antioxidant amino acid group contents, L-tyrosine and L-methionine amino acids. The activities of antioxidants contribute to fenton reaction inhibition, protecting SC DNA from radical-induced hydroxyl breaks in the strands. Hydroxyl radicals are highly reactive and toward biological molecules, such as DNA, proteins, and lipids, which can cause chemical cation changes (Dizdaroglu

& Jaruga, 2012). Hydroxyl radicals may react with sugar units through base groups or hydrogen abstraction. DNA damage occurred as an unpleasant oxygen-metabolism by product and other environmental insults. Most of the DNA damage that is regularly repaired and removed may not be adequately repaired, accumulating inside cells, causing genetic damage (Chatgililoglu & O'Neill, 2001). Previous study has reported that increased antioxidant capacity could be responsible for reducing cell damage (Samsampour et al., 2018).

Apart from free radical scavenging study, the activity of ACE-I inhibitor of each rice protein was also analysed. ACE-I inhibitors related to decreasing the blood pressure (Arora et al., 2014) in which ACE-I enzyme is responsible for the development of angiotensin II from angiotensin I. Angiotensin II is a vasoconstrictor causing blood pressure to increase. ACE-I inhibitors prevent renal disease development by decreasing intraglomerular pressure induced by angiotensin II (Tutor & Chichioco-Hernandez, 2017). Captopril was selected as the standard ACE inhibitor drug and was proved to be effective in inhibiting ACE activity at relatively lower concentrations than sample extracts. This was confirmed from the IC<sub>50</sub> values (Table 3), where captopril showed a significant low inhibitory concentration compared to the other samples. Several studies stated that some ACE inhibitory peptides were directly isolated from food materials without *in vitro* proteolysis (Maruyama & Suzuki, 1982; Shin et al., 2001). Li et al. (2004) reviewed that the ACE-inhibitory peptide activities were ranged from 0.20 to 246.7 mg/mL on with the IC<sub>50</sub> values. Most food-protein-derived antihypertensive peptides with high ACE-inhibitory activity were purified to have a relatively low molecular weight (Pihlanto-Leppälä et al., 2000). The current study showed inhibitor activity Ketan Hitam-2 was the highest indicated by its lowest IC<sub>50</sub> value and this indicated Ketan Hitam-2 seed protein has the best potency to inhibit the activity of ACE-I. As well as antioxidant properties, it has been reported that ACE-I inhibitor peptides are also known to have high hydrophobic amino acids (López-Barrios et al., 2014). Another study has published that ACE-I inhibitor peptides have high hydrophobicity, aromatics, and branched side chains amino acids (Rai et al., 2017).

In this study, the analysed were carried out on crude protein of pigmented rice seed. The high activity reported were the activity of crude protein which was isolated from pigmented rice seeds. Thus, it is a premature conclusion if consuming pigmented rice directly could reduce the blood pressure.

## 5 Conclusion

In conclusion, the present study successfully explored and demonstrated bioactive protein in pigmented rice seed related to their potential protective effect against free radicals and ACE activity-induced by oxidative stress. Seed protein of Ketan Hitam-2 showed a higher potency as a protection agent with high content of the antioxidant amino acid group, scavenge of ABTS •+ and OH• radicals, protect DNA damage, and increase the activity of ACE-I. These results provide information related to nutraceutical sources in the prevention of cellular disease including hypertension.

## Conflict of interest

The authors reported no potential conflict of interest.

## Acknowledgements

This research was supported by the University of Jember and the Ministry of Culture and Education (DRPM), The Republic of Indonesia.

## References

- Ahn, C. B., Kim, J. G., & Je, J. Y. (2014). Purification and antioxidant properties of octapeptide from salmon byproduct protein hydrolysate by gastrointestinal digestion. *Food Chemistry*, 147, 78-83. <http://dx.doi.org/10.1016/j.foodchem.2013.09.136>. PMID:24206688.
- Ajibola, C. F., Fashakin, J. B., Fagbemi, T. N., & Aluko, R. E. (2011). Effect of peptide size on antioxidant properties of African yam bean seed (*Sphenostylis stenocarpa*) protein hydrolysate fractions. *International Journal of Molecular Sciences*, 12(10), 6685-6702. <http://dx.doi.org/10.3390/ijms12106685>. PMID:22072912.
- Arora, P. K., Lifesciences, N., & Chauhan, A. (2014). ACE inhibitors: a comprehensive review. *International Journal of Pharmaceutical Sciences and Research*, 4(2), 532-548.
- Balindong, J. L., Ward, R. M., Liu, L., Rose, T. J., Pallas, L. A., Ovenden, B. W., Snell, P. J., & Waters, D. L. E. (2018). Rice grain protein composition influences instrumental measures of rice cooking and eating quality. *Journal of Cereal Science*, 79, 35-42. <http://dx.doi.org/10.1016/j.jcs.2017.09.008>.
- Bonjoch, N. P., & Tamayo, P. R. (2001). Protein content quantification by bradford method. In M. J. R. Roger (Ed.), *Handbook of plant ecophysiology techniques* (pp. 283-295). Dordrecht: Springer. [https://doi.org/10.1007/0-306-48057-3\\_19](https://doi.org/10.1007/0-306-48057-3_19).
- Chatgililoglu, C., & O'Neill, P. (2001). Free radicals associated with DNA damage. *Experimental Gerontology*, 36(9), 1459-1471. [http://dx.doi.org/10.1016/S0531-5565\(01\)00132-2](http://dx.doi.org/10.1016/S0531-5565(01)00132-2). PMID:11525869.
- Dabi, T., & Khanna, V. (2018). Effect of climate change on rice. *Agrotechnology*, 7(2), 1-7. <http://dx.doi.org/10.4172/2168-9881.1000181>.
- Dizdaroglu, M., & Jaruga, P. (2012). Mechanisms of free radical-induced damage to DNA. *Free Radical Research*, 46(4), 382-419. <http://dx.doi.org/10.3109/10715762.2011.653969>. PMID:22276778.
- Gupta, C., Prakash, D., & Gupta, S. (2013). Relationships between bioactive food components and their health benefits. In D. M. Martirosyan (Ed.), *Introduction to functional food science textbook*. Richardson: Food Science Publisher, pp. 66-85.
- Hazra, B., Biswas, S., & Mandal, N. (2008). Antioxidant and free radical scavenging activity of *Spondias pinnata*. *BMC Complementary and Alternative Medicine*, 8, 63. <http://dx.doi.org/10.1186/1472-6882-8-63>. PMID:19068130.
- Jamshidi-Kia, F., Lorigooini, Z., & Amini-Khoei, H. (2018). Medicinal plants: past history and future perspective. *Journal of HerbMed Pharmacology*, 7(1), 1-7. <http://dx.doi.org/10.15171/jhp.2018.01>.
- Karami, Z., Peighambari, S. H., Hesari, J., & Akbari-Adergani, B. (2019). Response surface methodology to optimize hydrolysis parameters in production of antioxidant peptides from wheat germ protein by alcalase digestion and identification of antioxidant peptides by LC-MS/MS. *Journal of Agricultural Science and Technology*, 21(4), 829-844.
- Li, G. H., Le, G. W., Shi, Y. H., & Shrestha, S. (2004). Angiotensin I-converting enzyme inhibitory peptides derived from food proteins



- and their physiological and pharmacological effects. *Nutrition Research*, 24(7), 469-486. [http://dx.doi.org/10.1016/S0271-5317\(04\)00058-2](http://dx.doi.org/10.1016/S0271-5317(04)00058-2).
- López-Barrios, L., Gutiérrez-Urbe, J. A., & Serna-Saldívar, S. O. (2014). Bioactive peptides and hydrolysates from pulses and their potential use as functional ingredients. *Shipin Kexue*, 79(3), R273-R283. <http://dx.doi.org/10.1111/1750-3841.12365>.
- Maruyama, S., & Suzuki, H. (1982). A peptide inhibitor of angiotensin I converting enzyme in the tryptic hydrolysate of casein. *Agricultural and Biological Chemistry*, 46(5), 1393-1394. <http://dx.doi.org/10.1080/00021369.1982.10865255>.
- Murdifin, M., Pakki, E., Rahim, A., Syaiful, S. A., Ismail, E., Evary, Y. M., & Bahar, M. A. (2015). Physicochemical properties of Indonesian pigmented rice (*Oryza sativa* Linn.) varieties from south Sulawesi. *Asian Journal of Plant Sciences*, 14(2), 59-65. <http://dx.doi.org/10.3923/ajps.2015.59.65>.
- Pérez, R. A., Iglesias, M. T., Pueyo, E., González, M., & De Lorenzo, C. (2007). Amino acid composition and antioxidant capacity of Spanish honeys. *Journal of Agricultural and Food Chemistry*, 55(2), 360-365. <http://dx.doi.org/10.1021/jf062055b>. PMID:17227066.
- Pihlanto-Leppälä, A., Koskinen, P., Piilola, K., Tupasela, T., & Korhonen, H. (2000). Angiotensin I-converting enzyme inhibitory properties of whey protein digests: Concentration and characterization of active peptides. *The Journal of Dairy Research*, 67(1), 53-64. <http://dx.doi.org/10.1017/S0022029999003982>. PMID:10717843.
- Pouzo, L. B., Descalzo, A. M., Zaritzky, N. E., Rossetti, L., & Pavan, E. (2016). Antioxidant status, lipid and color stability of aged beef from grazing steers supplemented with corn grain and increasing levels of flaxseed. *Meat Science*, 111, 1-8. <http://dx.doi.org/10.1016/j.meatsci.2015.07.026>. PMID:26318758.
- Rai, A. K., Sanjukta, S., & Jeyaram, K. (2017). Production of angiotensin I converting enzyme inhibitory (ACE-I) peptides during milk fermentation and their role in reducing hypertension. *Critical Reviews in Food Science and Nutrition*, 57(13), 2789-2800. <http://dx.doi.org/10.1080/10408398.2015.1068736>. PMID:26463100.
- Samsampour, D., Sadeghi, F., Asadi, M., & Ebrahimzadeh, A. (2018). Effect of nitric oxide (NO) on the induction of callus and antioxidant capacity of *Hyoscyamus Niger* under in vitro salt stress. *Journal of Applied Botany and Food Quality*, 91, 24-32. <http://dx.doi.org/10.5073/JABFQ.2018.091.004>.
- Samyori, D., Das, A. B., & Deka, S. C. (2017). Pigmented rice a potential source of bioactive compounds: a review. *International Journal of Food Science & Technology*, 52(5), 1073-1081. <http://dx.doi.org/10.1111/ijfs.13378>.
- Samyori, D., Deka, S. C., & Das, A. B. (2016). Phytochemical and antioxidant profile of pigmented and non-pigmented rice cultivars of Arunachal Pradesh, India. *International Journal of Food Properties*, 19(5), 1104-1114. <http://dx.doi.org/10.1080/10942912.2015.1055761>.
- Sati, R., & Singh, S. (2019). Pigmented rice: a potential ingredient for extruded products review paper. *Journal of Pharmacognosy and Phytochemistry*, 8(3), 700-702.
- Shahbandeh, M. (2019). *Statista: global rice consumption 2018/2019, by country*. Retrieved from <https://www.statista.com/statistics/255971/top-countries-based-on-rice-consumption-2012-2013/>
- Shin, Z. I., Yu, R., Park, S. A., Chung, D. K., Ahn, C. W., Nam, H. S., Kim, K. S., & Lee, H. J. (2001). His-His-Leu, an angiotensin I converting enzyme inhibitory peptide derived from Korean soybean paste, exerts antihypertensive activity in vivo. *Journal of Agricultural and Food Chemistry*, 49(6), 3004-3009. <http://dx.doi.org/10.1021/jf001135r>. PMID:11410001.
- Shinta, S., Indriyani, S., & Arisoelaningsih, E. (2014). Morphological variation of six pigmented rice local varieties grown in organic rice field at Sengguruh Village, Kepanjen Subdistrict, Malang District. *Journal of Tropical Life Science*, 4(2), 149-160. <http://dx.doi.org/10.11594/jtls.04.02.12>.
- Siswoyo, T. A., Mardiana, E., Lee, K. O., & Hoshokawa, K. (2011). Isolation and characterization of antioxidant protein fractions from melinjo (*Gnetum gnemon*) seeds. *Journal of Agricultural and Food Chemistry*, 59(10), 5648-5656. <http://dx.doi.org/10.1021/jf2000647>. PMID:21486028.
- Supriyadi, A., Arum, L. S., Nugraha, A. R. I. S., Agung, A., Ratnadewi, I., & Siswoyo, T. R. I. A. (2019). Current research in nutrition and food science revealing antioxidant and antidiabetic potency of Melinjo (*Gnetum gnemon*) seed protein hydrolysate at different stages of seed maturation. *Current Research in Nutrition and Food Science Journal*, 7(2), 479. <http://dx.doi.org/10.12944/CRNFSJ.7.2.17>.
- Szkudzińska, K., Smutniak, I., Rubaj, J., Korol, W., & Bielecka, G. (2017). Method validation for determination of amino acids in feed by UPLC. *Accreditation and Quality Assurance*, 22(5), 247-252. <http://dx.doi.org/10.1007/s00769-017-1281-9>.
- Tan, B. L., Norhaizan, M. E., Liew, W. P. P., & Sulaiman Rahman, H. (2018). Antioxidant and oxidative stress: a mutual interplay in age-related diseases. *Frontiers in Pharmacology*, 9, 1162. <http://dx.doi.org/10.3389/fphar.2018.01162>. PMID:30405405.
- Tutor, J. T., & Chichioco-Hernandez, C. L. (2017). Angiotensin-converting enzyme inhibition of fractions from *Eleusine indica* leaf extracts. *Pharmacognosy Journal*, 10(1), 25-28. <http://dx.doi.org/10.5530/pj.2018.1.5>.
- Wijaya, D. N., Susanto, F. A., Purwestri, Y. A., Ismoyowati, D., & Nuringtyas, T. R. (2018). NMR metabolite comparison of local pigmented rice in Yogyakarta. *Indonesian Journal of Biotechnology*, 22(2), 68. <http://dx.doi.org/10.22146/ijbiotech.27308>.
- Wijesekara, I., Qian, Z. J., Ryu, B. M., Ngo, D. H., & Kim, S. K. (2011). Purification and identification of antihypertensive peptides from seaweed pipefish (*Syngnathus schlegelii*) muscle protein hydrolysate. *Food Research International*, 44(3), 703-707. <http://dx.doi.org/10.1016/j.foodres.2010.12.022>.
- Xu, N., Chen, G., & Liu, H. (2017). Antioxidative categorization of twenty amino acids based on experimental evaluation. *Molecules*, 22(12), 2066. <http://dx.doi.org/10.3390/molecules22122066>.
- Zou, T.-B., He, T.-P., Li, H.-B., Tang, H.-W., & Xia, E.-Q. (2016). The structure-activity relationship of the antioxidant peptides from natural proteins. *Molecules*, 21(1), 72. <http://dx.doi.org/10.3390/molecules21010072>. PMID:26771594.
- Zou, Z., Wang, M., Wang, Z., Aluko, R. E., & He, R. (2020). Antihypertensive and antioxidant activities of enzymatic wheat bran protein hydrolysates. *Journal of Food Biochemistry*, 44(1), e13090. <http://dx.doi.org/10.1111/jfbc.13090>. PMID:31663146.