

Inactivation of *E. coli* and *B. subtilis* spores in ozonized cassava starch

Emanuele Oliveira Cerqueira AMORIM¹, Alline Artigiani Lima TRIBST¹,
Pedro Esteves Duarte AUGUSTO^{1,2}, Marcelo CRISTIANINI^{1*}

Abstract

In the present study, the efficacy of ozone inactivation of *B. subtilis* spores and *E. coli* in cassava starch was evaluated. Cassava starch with 18 and 30% moisture content was processed with ozone at concentrations of 40-118 ppm and exposure times of 15-120 minutes. The processing at 113 ppm/120 minutes (maximum exposure level to ozone evaluated) at 18% of moisture content did not cause significant reduction of *B. subtilis* spores and caused the reduction of only 2 decimal of *E. coli*. On the other hand, when the ozonation process was carried out for 120 minutes at 30% of moisture content, 3.6 decimal reduction of *B. subtilis* was achieved at 40 ppm of ozone and total *B. subtilis* load reduction (>5 log cycles) was observed at 118 ppm of ozone. Similarly, total *E. coli* load reduction (>7 log cycles) was achieved at 40 ppm of ozone exposure for 60 minutes. Therefore, the results indicate that the ozone efficacy against microorganisms in cassava starch was mainly dependent on the sample moisture content and to ozone concentration and exposure time. Moreover, it was observed that ozone is a promising technology to reduce microbial counts in dried food.

Keywords: ozonation; food processing; non-thermal technologies.

1 Introduction

Today's consumers have demanded processed products very similar to fresh food (TRIBST; SANT'ANA; DE MASSAGUER, 2009). For such reason, the development of non-thermal methods for food processing has been extensively studied – e.g. high-hydrostatic pressure, irradiation, UV radiation, pulsed electric field, pulsed light, ultrasound, and high-pressure homogenization, and ozone (DIELS; WUYTACK; MICHIELS, 2003; TRIBST; SANT'ANA; DE MASSAGUER, 2009; KLOCKOW; KEENER, 2009).

It is expected that products processed by such technologies maintain the nutrients and sensory characteristics better than thermal process. Among these technologies, ozone stands out for its antimicrobial properties and for being totally degradable in oxygen with no waste/toxic products (TIWARI et al., 2008, 2010); thus it is generally recognized as safe (GRAS) for food applications (GUZEL-SEYDİM; GREENE; SEYDİM, 2004; DHILLON et al., 2009; TIWARI et al., 2010). An additional advantage is the lower cost of ozone equipment (GUZEL-SEYDİM; GREENE; SEYDİM, 2004).

The ozone is a high oxidative compound with a broad antimicrobial spectrum; it is able to inactivate bacterial vegetative cells and spores, yeast, molds, and viruses and to kill insects in stored grains and degrade mycotoxins (TIWARI et al., 2010). Ozone inactivates microorganisms by the progressive oxidation of vital cellular components. The microbial surface is the primary target of ozonation with the oxidation of the polyunsaturated fatty acids and consequent loss of selective permeability and cell disruption. Additionally, ozone causes

the oxidation of sulfhydryl groups and amino acids of enzymes, peptides, and proteins including nucleic acids and vital enzymes (KHADRE; YOUSEF; KIM, 2001; GUZEL-SEYDİM; GREENE; SEYDİM, 2004). In general, Gram negative bacteria are more sensitive to ozone treatment than Gram positive ones (KHADRE; YOUSEF; KIM, 2001). However, several other factors affect the ozone efficacy including the strain of the microorganism, age of the culture, density of the treated population, and presence of ozone-demanding medium components and method of applying ozone (KHADRE; YOUSEF; KIM, 2001)

In spores, ozone degrades the coat layer components thus exposing the cortex and the core to oxidation by ozone (KHADRE; YOUSEF, 2001; TIWARI et al., 2010). The molecular ozone and the reactive by-products of ozone decomposition (free radicals: OH[·], O₂ and HO₃[·]) have antimicrobial activity. Due to its high efficacy, low concentration and short contact time are normally enough to achieve the desired microbial inactivation.

Ozone has been previously used in food industry to inactivate microorganisms in packages and stainless steel surfaces (KHADRE; YOUSEF, 2001), for microbial decontamination of poultry carcasses, on chilled water treatment and recycling (GUZEL-SEYDİM; GREENE; SEYDİM, 2004; KHADRE; YOUSEF, 2001), for fish shelf-life extension (PASTORIZA et al., 2008), for fruit juice processing (TIWARI et al., 2009a, b), for sanitation of fruit and vegetables (ACHEN; YOUSEF, 2001; BIALKA; DEMIRCI, 2007; NAJAFI; KHODAPARAST, 2009; KIM; YOUSEF; CHISM, 1999; KOSEKI; ISOBE, 2006), and

Received 10/7/2012

Accepted 14/2/2013 (005780)

¹ Department of Food Technology – DTA, School of Food Engineering – FEA, University of Campinas – UNICAMP, Rua Monteiro Lobato, 80, CP 6121, Cidade Universitária Zeferino Vaz, CEP 13083-862, Campinas, SP, Brazil, e-mail: olecram@fea.unicamp.br

² Technical School of Campinas – COTUCA, University of Campinas – UNICAMP, Campinas, SP, Brazil

*Corresponding author

DOI: <http://dx.doi.org/10.1590/S0101-20612013005000043>

for decontamination of dry products as seeds, grains, and pepper (TIWARI et al., 2010, AKBAS; OZDEMIR, 2006; EMER; AKBAS; OZDEMIR, 2008; IBANOGLU, 2001, 2002; KETTERINGHAM et al., 2006; ZHAO; CRANSTON, 1995).

In addition to ozone antimicrobial potential, this gas can also be used to change physicochemical characteristics of corn, sago, and tapioca starch (CHAN; BHAT; KARIM, 2009) and wheat flour (LI et al., 2012) by oxidation improving their functional performance and presenting better results than those obtained by the traditional oxidation process (using chlorine).

Cassava starch is an important raw material in the food industry due its absence of "cereal flavor", typically observed in corn starch, its viscoamylographic standard during cooking (DEMIATE et al., 2005), its *pasta* low temperature, and its syneresis low susceptibility (SILVA et al., 2003).

Considering that cassava starch is extracted from plant root, it is potentially contaminated with many microorganisms, mainly bacterial spores. Therefore, some processes are required to guarantee the microbial stability of cassava starch before its use as raw material in the food industry. Due to low temperature gelation, the thermal process application for microbial inactivation of cassava starch is not feasible (CEREDA; VILPOUX, 2005); therefore, the development of a non-thermal process is necessary.

The *Bacillus subtilis* is considered an ideal target for the chemical process of microbial inactivation due to its safety and high resistance to chemical compounds (MOREAU; ORANGE; FEUILLOLEY, 2008). In addition to the analysis of the spore forming microorganisms, which are potentially starch pathogens or spoilers, the evaluation of *Escherichia coli* was carried out considering the the Brazilian food legislation (BRASIL, 2001), which establishes a maximum limit of coliforms able to growth at 45 °C of 10² CFU/g of starch.

Therefore, based on the ozone properties as a sanitizer and the requirements of a non-thermal process to stabilize cassava starch, this study aimed to evaluate the efficacy of ozone inactivation of *B. subtilis* spores and *E. coli* in cassava starch.

2 Materials and methods

2.1 Starch and microorganisms

Cassava starch was supplied by the company Corn Products Brasil (Conchal, Brazil). It was previously sterilized using γ -irradiation (10 kGy) performed by the Brazilian Radiation company (EMBRARAD- Cotia, Brazil) to inactivate its native microorganisms.

The lyophilized strains of *Bacillus subtilis* ATCC 6633 and *Escherichia coli* ATCC 8739 were donated by André Tosello Research and Technology Foundation – FAT (Campinas, Brazil – www.fat.org.br).

2.2 Cell culture suspensions

The *E. coli* and *B. subtilis* cultures were activated using nutrient broth and kept under refrigeration on nutrient agar slants.

Petri dishes containing nutrient agar were inoculated with *E. coli* in and incubated at 37 °C for 24 hours. To prepare *E. coli* suspension, the cell mass was harvested in sterile peptone water and used on the same day. The *B. subtilis* spores suspension was prepared as described by Wuytach, Boven and Michiels (1998), with few modifications: *B. subtilis* sporulation was carried out on nutrient agar supplement with 20 mg.L⁻¹ of manganese sulphate, incubation was carried out at 30 °C for 14 days and, after harvesting, spores were three times centrifuged at 10,000 rpm for 10 minutes at 10 °C. A heat shock at 70 °C for 20 min was carried out before *B. subtilis* count. Both microorganisms were plate counted on nutrient agar and incubated at 30 °C for 48 hours (*B. subtilis*) or 37 °C for 24 hours (*E. coli*).

2.3 Ozonation

Ozone was produced from concentrated O₂ air (Model NewLife Elite, AirSep Corporation, New York, United States) using an Interzone generator (Line DEVOC®, Model SGOZ-123, Interzone Group of Brazil, Jundiaí, Brazil) with 5 g·h⁻¹ of production capacity. The ozonized air flow was 14 m³·h⁻¹, and the ozone concentration in the gas was continuously monitored by an ozone analyzer (Model UV-100, Eco Sensors, Inc., Santa Fe, USA). Ozonized air was injected in a stainless steel powder mixer of 25 L through two air inlet devices located in the mixer lid. The spinning blades promoted an efficient contact between starch and ozonized air.

2.4 Ozone treatment of cassava starch inoculated with *E. coli*

1 kg of cassava starch (initial moisture of 11%) was inoculated with 100 mL of *E. coli* suspension reaching a final count of 10⁶-10⁸ CFU·g⁻¹ and moisture of 18% (maximum limit of starch moisture allowed by the Brazilian law). For assays using cassava starch moisture at 30% (moisture of starch immediately after cassava centrifugation carried out in the starch industry just before the drying processing), the samples were previously humidified with sterile water, followed by inoculation with 100 mL of *E. coli* suspension. After inoculation, cassava starch rested for one hour before ozonation to enable the adherence of microorganism to starch.

The starch of 18% moisture content was processed at ozone concentration of 40 and 113 mg.L⁻¹ for 30, 60, 90, and 120 minutes. The starch of 30% moisture content was processed at ozone concentration of 40 mg.L⁻¹ for 30, 60, 90, and 120 minutes or at ozone concentration of 118 mg.L⁻¹ for 15, 30, 45, and 60 minutes. The 113 and 118 mg.L⁻¹ were the higher ozone levels reached in the equipment for the samples at 18 and 30% of moisture. This small variation in the maximum ozone concentration can be attributed to few oscillations in the oxygen concentration and ozone generated in the equipment.

Survivor counts of *E. coli* were performed on nutrient agar just after the ozonation process. The control samples were obtained by exposing previously inoculated cassava starch to air (without ozone) under the same conditions used for the ozonized samples. Initial count was determined by starch plating immediately after inoculation. The samples were incubated at

37 °C for 24 h. The Number of Decimal Reductions (NDR) of each process was determined according to Equation 1.

$$\text{NDR} = \log(\text{initial count}) - \log(\text{survivor's count}) \quad (1)$$

2.5 Ozone treatment of cassava starch inoculated with *B. subtilis*

1 kg of cassava starch was inoculated with *B. subtilis* spores suspension at the same conditions described for *E. coli*. *B. subtilis* final count was 10^7 CFU·g⁻¹, and cassava starch moisture was 18 and 30%. The inoculated sample was allowed to rest for 1 hour for the adherence of microorganism to starch, and the starch of 18% moisture content was then ozonized at ozone concentration of 118 mg.L⁻¹ for 30, 60, 90, and 120 minutes, and starch of 30% moisture content was processed at 40 and 118 mg.L⁻¹ for 30, 60, 90, and 120 minutes.

The *B. subtilis* enumeration was carried out after a heat shock (70 °C/20 minutes) for spore activation. The samples were plated on nutrient agar and incubated at 30 °C for 48 hours. The control samples were obtained by exposing previously inoculated cassava starch to air (without ozone) under the same conditions used for the ozonized samples. Initial count was determined by starch plating immediately after inoculation. The Number of Decimal Reductions (NDR) of each process was determined by Equation 1.

2.6 Statistical analysis

The analysis of variance (ANOVA) was performed to compare the effects of the different treatments, and the Tukey test was used to determine their differences at a 5% confidence level. Statistical analyses were carried out using STATISTICA 5.0 software (StatSoft, Inc., Tulsa, Okla., U.S.A). All of the tested conditions were evaluated in triplicate. The results were presented as mean ± standard deviation.

3 Results and discussion

3.1 *E. coli* inactivation

The *E. coli* inactivation on ozonized 18% moisture cassava starch is shown in Figure 1. Statistical analysis indicated that 30, 60, and 90 minutes of contact time with 40 mg.L⁻¹ ozone caused about one decimal reduction, with no differences between




Figure 1. Inactivation of *E. coli* in ozonized 18% moisture cassava starch. Vertical bars represent the standard deviation of each value.

them ($p > 0.05$). On the other hand, 120 minutes of contact at this same concentration caused about 2 decimal reduction, and it was different from other conditions evaluated ($p < 0.05$). The results of cassava starch ozonized at the concentration of 113 mg.L⁻¹ showed 1-2 decimal reduction, with no significant differences between different ozone contact times. Considering the control sample results (<0.5 NDR), it was observed that ozone was able to reduce *E. coli* counts in 18% moisture cassava starch, but these reductions (<2 decimal reductions) were not enough to guarantee the cassava starch safety when used as food raw material. Moreover, no significant differences were observed between *E. coli* inactivation in the samples ozonized at 40 and 113 mg.L⁻¹, which probably indicates ozone saturation at concentration of 40 mg.L⁻¹.

The effect of starch humidification on the ozone efficacy was evaluated through ozonation process carried out on previously humidified starch at 30% moisture (moisture of the starch after centrifugation). The *E. coli* inactivation at this condition is shown in Table 1.

The control treatment (carried out with no ozone injection) showed 1.3-1.6 decimal reductions, indicating that at 30% moisture, the microorganism was more sensitive in non-ideal growth media or its recovery after the process was reduced. Comparing the results of control and ozonized samples, on the other hand, it was observed that ozonation was able to inactivate the microorganism.

The results showed that increases in ozone concentration and exposure time improve *E. coli* inactivation during the process. Processes applying ozone concentration at 40 mg.L⁻¹ for 60 minutes or 118 mg.L⁻¹ for 30 minutes are enough to guarantee an adequate level of *E. coli* inactivation assuring the safety of foods with this starch as raw material.

E. coli inactivation by ozone has been previously studied by several authors in different products including water, fruits, vegetables, meat, and dry products (MUTHUKUMARAPPAN; HALAWEISH; NAIDU, 2000; ZHAO; CRANSTON, 1995; KHADRE; YOUSEF; KIM, 2001; AKBAS; OZDEMIR, 2006, 2008b; EMER; AKBAS; OZDEMIR, 2008). These studies were carried out using different equipment and strains of *E. coli*, but, in general, it was observed that higher inactivation occurred in water or products with high moisture (MUTHUKUMARAPPAN; HALAWEISH; NAIDU, 2000).

The use of 0.35 ppm of ozone resulted in 5 decimal reductions of *E. coli* in water, and the use of 18 ppm for 50 minutes resulted in total load inactivation of *E. coli* O157:H7

Table 1. Inactivation of *E. coli* in ozonized 30% moisture cassava starch.

Time (minutes)	NRD at different ozone concentration		
	*0 mg.L ⁻¹	40 mg.L ⁻¹	118 mg.L ⁻¹
15	-	-	3.57
30	1.26	3.72	>6
45	-	-	>7
60	1.45	>7	>7
90	1.51	>7	-
120	1.64	>7	-

*Results of control sample. These values are the NDR average of two processes.

in several foods (MUTHUKUMARAPPAN; HALAWEISH; NAIDU, 2000). The concentration of 1.0 ppm of ozone applied for 360 minutes resulted in 3.5 decimal reduction in dried figs (AKBAS; OZDEMIR, 2008b). For pistachio (AKBAS; OZDEMIR, 2006), red pepper (AKBAS; OZDEMIR, 2008b), and black pepper (EMER; AKBAS; OZDEMIR, 2008), similar inactivation level (about 2 decimal reductions) was obtained after ozonation for 120-360 minutes.

Therefore, the presence of water in food is very important to reach high microbial inactivation level by ozone, as observed in the present study. Similarly, previous results showed that microbial inactivation by ozone on black pepper was improved at high moisture content (ZHAO; CRANSTON, 1995). It is known that ozone is a powerful oxidative compound that reacts with microorganism and organic materials (KHADRE; YOUSEF; KIM, 2001; MUTHUKUMARAPPAN; HALAWEISH; NAIDU, 2000). Thus, the increase in the organic material content may have affected the ozone efficacy.

Additionally, the ozone gas is rapidly decomposed in oxygen (MUTHUKUMARAPPAN; HALAWEISH; NAIDU, 2000). It is possible that ozone partial dissolution in the food free water results in the increase in the antimicrobial half-life, which can become more effective. Another possible effect is that moisture can improve the gas-product contact area or promote water free-radical formation that was effective for microbial inactivation in addition to the ozone effects.

Comparing the results obtained for *E. coli* inactivation in the cassava starch and other dry products (AKBAS; OZDEMIR, 2006, 2008b; EMER; AKBAS; OZDEMIR, 2008), it was observed that, for the other products, lower ozone concentration of 0.5-1 mg.L⁻¹ was required to reach the same level of inactivation (i.e., 40 times smaller than that required for cassava starch). Additionally to the differences in the moisture content and water activity of these foods and cassava starch, the differences between the results obtained can be explained by different ozonized air flow and the amount of product ozonized, which interfere in the ozone available amount and also on ozone mass/product mass ratio. Moreover, the *E. coli* adherence in cassava starch for one hour before ozonation and the powder small dimension with rough surface can also affect ozone activity.

3.2 *B. subtilis* inactivation

Considering the results obtained for *E. coli* inactivation in ozonized 18% moisture cassava starch and the expected higher ozone resistance of *B. subtilis*, its inactivation in 18% moisture cassava starch was evaluated only at the ozone concentration of 118 mg.L⁻¹ (which was the maximum ozone concentration reached in the ozonation equipment). No *B. subtilis* spore reduction was observed in cassava starch of 18% moisture content, even after ozonation at 118 mg.L⁻¹ concentration for 120 minutes (data not shown). This indicates that, at this cassava starch moisture, ozone was not effective for *B. subtilis* inactivation.

Comparing the results obtained for *B. subtilis* and *E. coli* inactivation (113 mg.L⁻¹ ozone concentration), it was confirmed that *Bacillus* spores are more resistant to ozone than *E. coli*, as previously described by Akbas and Ozdemir (2008a, b) and Muthukumarappan, Halaweish and Naidu (2000). This can be




Figure 2. Inactivation of *B. subtilis* in ozonized 30% moisture cassava starch. Vertical bars represent the standard deviation of each value.

explained by the high resistance of the spore coat and outer membrane to the oxidative mechanisms of the ozone (AKBAS; OZDEMIR, 2008a, b; MUTHUKUMARAPPAN; HALAWEISH; NAIDU, 2000), compared with oxidation of vegetative cell envelope and membrane.

The effect of 30% moisture starch on the ozone efficacy against *B. subtilis* was also evaluated, and the results are shown in Figure 2. The results showed that the control processes caused no significant reduction on *B. subtilis* spores. On the other hand ozone promoted *B. subtilis* spores inactivation at the concentration of 40 and 118 mg.L⁻¹, and it was higher at higher ozone concentration ($p < 0.05$). In addition, increased contact time up to 60 and 120 minutes caused a significant increase in spore inactivation at ozone concentration of 40 and 118 mg.L⁻¹, respectively. Again, with the increase in the starch moisture, a significant increase in the microbial inactivation was observed. Data on *B. subtilis* spores in water also demonstrated higher level of spore inactivation, with 3 log reductions after processing with 0.7 ppm of ozone, indicating that the water content is important for spore inactivation.

It is interesting to highlight that by increasing ozone concentration by three times (40 mg.L⁻¹ to 118 mg.L⁻¹) it was able to reduce 50% of process time aiming to achieve the same level of *B. subtilis* spores inactivation, which can be a possible solution when low time processing is required.

Previous data on ozone treatment of wheat and wheat flour showed low level of inactivation of total aerobic microorganism and yeast and moulds (<1 decimal reduction) using ozonated water at concentrations up to 16.5 ppm (IBANOGLU, 2001, 2002; LI et al., 2012). These results corroborate the findings that food with low water content possible slows ozone reaction speed and therefore reduces the disinfection ability of the ozone gas.

4 Conclusion

The ozonation process can be effective for *E. coli* and *B. subtilis* inactivation in cassava starch, and it is highly dependent on starch moisture. At low starch moisture (18%), the process was not enough to guarantee microbial inactivation; however, at 30% moisture starch, it was possible to reach >5 decimal reductions, ensuring safe products. *E. coli* were less resistant to ozone treatment, and it was inactivated after 30 minutes treatment with 118 ppm of ozone. On the other hand, 90 minutes are required to reach an appropriated inactivation of *B. subtilis*. Therefore, the application of ozone at these conditions makes cassava starch a safe product to be used as a raw material in the food industry.

Acknowledgments

The authors would like to thank the São Paulo Research Foundation (FAPESP) for financial support to the project # 2008/10942-5, the Brazilian National Research Council (CNPq) for the E.O.C Amorim fellowship, and the Corn Products Brasil for the cassava starch donation.

References

- ACHEN, M.; YOUSEF, A. E. Efficacy of ozone against *Escherichia coli* O157:H7 on apples. **Journal of Food Science**, v. 66, n. 9, p. 1380-1384, 2001. <http://dx.doi.org/10.1111/j.1365-2621.2001.tb15218.x>
- AKBAS, M. Y.; OZDEMIR, M. Effectiveness of ozone for inactivation of *Escherichia coli* and *Bacillus cereus* in pistachios. **International Journal of Food Science and Technology**, v. 41, n. 5, p. 513-519, 2006. <http://dx.doi.org/10.1111/j.1365-2621.2005.01099.x>
- AKBAS, M. Y.; OZDEMIR, M. Application of gaseous ozone to control populations of *Escherichia coli*, *Bacillus cereus* and *Bacillus cereus* spores in dried figs. **Food Microbiology**, v. 25, n. 2, p. 386-391, 2008a. <http://dx.doi.org/10.1016/j.fm.2007.09.007>
- AKBAS, M. Y.; OZDEMIR, M. Effect of gaseous ozone on microbial inactivation and sensory of flaked red peppers. **International Journal of Food Science and Technology**, v. 43, n. 9, p. 1657-1662, 2008b. <http://dx.doi.org/10.1111/j.1365-2621.2008.01722.x>
- BIALKA, K. L.; DEMIRCI, A. Utilization of gaseous ozone for the decontamination of *Escherichia coli* O157:H7 and *Salmonella* on raspberries and strawberries. **Journal of Food Protection**, v. 70, n. 5, p. 1093-1098, 2007.
- BRASIL. Ministério da Saúde. Resolução RDC nº 12, de 02 de janeiro de 2001. Aprova o regulamento técnico sobre padrões microbiológicos para alimentos. **Diário Oficial da República Federativa do Brasil, Brasília**, DF, 10 jan. 2001.
- CEREDA, M. P.; VILPOUX, O. **Uso de lâmpada ultravioleta Germetec para redução de conteúdo microbiano de amido comercial de mandioca**. Centro de Tecnologias para o Agro-negócio, Universidade Católica Dom Bosco, 2005. Nota prévia.
- CHAN, H. T.; BHAT, R.; KARIM, A. A. Physicochemical and functional properties of ozone-oxidized starch. **Journal of Agricultural and Food Chemistry**, v. 57, p. 5965-5970, 2009. <http://dx.doi.org/10.1021/jf9008789>
- DEMIATE, I. M. et al. Viscographic characteristics of chemically modified cassava starches assessed by RVA. **Publicio UEPG - Ciências Exatas e da Terra**, v. 11, n. 1, p. 7-17, 2005.
- DHILLON, B. et al. Development and evaluation of an ozonated water system for antimicrobial treatment of durum wheat. **Journal of Food Science**, v. 74, n. 7, p. E396-403, 2009. <http://dx.doi.org/10.1111/j.1750-3841.2009.01275.x>
- DIELS, A. M. J.; WUYTACK, E. Y.; MICHIELS, C. W. Modeling inactivation of *Staphylococcus aureus* and *Yersinia enterocolitica* by high-pressure homogenization at different temperatures. **International Journal of Food Microbiology**, v. 87, n. 1-2, p. 55-62, 2003. [http://dx.doi.org/10.1016/S0168-1605\(03\)00050-3](http://dx.doi.org/10.1016/S0168-1605(03)00050-3)
- EMER, Z.; AKBAS, M. Y.; OZDEMIR, M. Bactericidal activity of ozone against *Escherichia coli* in whole and ground black peppers. **Journal of Food Protection**, v. 71, n. 5, p. 914-917, 2008.
- GUZEL-SEYDIM, Z. B.; GREENE, A. K.; SEYDIM, A. C. Use of ozone in the food industry. **LWT - Food Science and Technology**, v. 37, n. 4, p. 453-460, 2004.
- IBANOGLU, S. Influence of tempering with ozonated water on the selected properties of wheat flour. **Journal of Food Engineering**, v. 48, n. 4, p. 345-350, 2001. [http://dx.doi.org/10.1016/S0260-8774\(00\)00177-1](http://dx.doi.org/10.1016/S0260-8774(00)00177-1)
- IBANOGLU, S. Wheat washing with ozonated water: effects on selected flour properties. **International Journal of Food Science and Technology**, v. 37, n. 5, p. 579-584, 2002. <http://dx.doi.org/10.1046/j.1365-2621.2002.00593.x>
- KHADRE, M. A.; YOUSEF, A. E. Decontamination of a multilaminated aseptic food packaging material and stainless steel by ozone. **Journal of Food Safety**, v. 21, p. 1-13, 2001. <http://dx.doi.org/10.1111/j.1745-4565.2001.tb00304.x>
- KHADRE, M. A.; YOUSEF, A. E.; KIM, J-G. Microbiological aspects of ozone applications in food: a review. **Journal of Food Science**, v. 66, n. 9, p. 1242-1252, 2001. <http://dx.doi.org/10.1111/j.1365-2621.2001.tb15196.x>
- KETTERINGHAM, L. et al. Application of aqueous ozone for treating pre-cut green peppers (*Capsicum annum* L.). **Journal of Food Engineering**, v. 76, p. 104-111, 2006. <http://dx.doi.org/10.1016/j.jfoodeng.2005.05.019>
- KIM, J.-G.; YOUSEF, A. E.; CHISM, G. W. Use of ozone to inactivate microorganisms on lettuce. **Journal of Food Safety**, v. 19, p. 17-34, 1999. <http://dx.doi.org/10.1111/j.1745-4565.1999.tb00231.x>
- KLOCKOW, P. A.; KEENER, K. M. Safety and quality assessment of packaged spinach treated with a novel ozone-generation system. **LWT - Food Science and Technology**, v. 42, n. 6, p. 1047-1053, 2009.
- KOSEKI, S.; ISOBE, S. Effect of ozonated water treatment on microbial control and on browning of Iceberg Lettuce (*Lactuca sativa* L.). **Journal of Food Protection**, v. 69, n. 1, p. 154-160, 2006.
- LI, M. et al. Evaluation the quality characteristics of wheat flour and shelf-life of fresh noodles as affected by ozone treatment. **Food Chemistry**, v. 135, p. 2163-2169, 2012. <http://dx.doi.org/10.1016/j.foodchem.2012.06.103>
- MOREAU, M.; ORANGE, N.; FEUILLOLEY, M. G. J. Non-thermal plasma technologies: New tools for bio-decontamination. **Biotechnology Advances**, v. 26, n. 6, p. 610-617, 2008. <http://dx.doi.org/10.1016/j.biotechadv.2008.08.001>
- MUTHUKUMARAPPAN, K.; HALAWEISH, F.; NAIDU, A. S. Ozone. In: NAIDU, A. S. **Natural Food Antimicrobial Systems**. CRC Press, 2000. <http://dx.doi.org/10.1201/9781420039368.ch29>
- NAJAFI, M. B. H.; KHODAPARAST, M. H. H. Efficacy of ozone to reduce microbial populations in date fruits. **Food Control**, v. 20, p. 27-30, 2009. <http://dx.doi.org/10.1016/j.foodcont.2008.01.010>
- PASTORIZA, L. et al. Use of sterile and ozonized water as a strategy to stabilize the quality of stored refrigerated fresh fish. **Food Control**, v. 19, p. 772-780, 2008. <http://dx.doi.org/10.1016/j.foodcont.2007.08.003>
- SILVA, G. O. et al. Amidos nativos e modificados: propriedades e aplicações em alimentos. **Boletim da Sociedade Brasileira de Ciência e Tecnologia de Alimentos**, v. 37, p. 101-106, 2003. Suplemento.
- TIWARI, B. K. et al. Effect of ozonation on the rheological and colour characteristics of hydrocolloid dispersions. **Food Research International**, v. 41, n. 10, p. 1035-1043, 2008. <http://dx.doi.org/10.1016/j.foodres.2008.07.011>
- TIWARI, B. K. et al. Anthocyanin and colour degradation in ozone treated blackberry juice. **Innovative Food Science and Emerging Technologies**, v. 10, n. 1, p. 70-75, 2009a. <http://dx.doi.org/10.1016/j.ifset.2008.08.002>

- TIWARI, B. K. et al. Effect of ozone processing on anthocyanins and ascorbic acid degradation of strawberry juice. **Food Chemistry**, v. 113, n. 4, p. 1119-1126, 2009b. <http://dx.doi.org/10.1016/j.foodchem.2008.08.085>
- TIWARI, B. K. et al. Review: Application of ozone in grain processing. **Journal of Cereal Science**, v. 51, p. 248-255, 2010. <http://dx.doi.org/10.1016/j.jcs.2010.01.007>
- TRIBST, A. A. L.; SANT'ANA, A. S.; DE MASSAGUER, P. R. Review: Microbiological quality and safety of fruit juices – past, present and future perspectives. **Critical Reviews in Microbiology**, v. 35, n. 4, p. 310-339, 2009. PMID:19863382. <http://dx.doi.org/10.3109/10408410903241428>
- WUYTACH, E. Y.; BOVEN, S.; MICHIELS, C. W. Comparative Study of Pressure-Induced Germination of *Bacillus subtilis* Spores at Low and High Pressures. **Applied and Environmental Microbiology**, v. 64, n. 9, p. 3220-3224, 1998.
- ZHAO, J.; CRANSTON, P. M. Microbial decontamination of black pepper by ozone and the effect of the treatment on volatile oil constituents of the spice. **Journal of the Science of Food and Agriculture**, v. 68, n. 1, p. 11-18, 1995. <http://dx.doi.org/10.1002/jsfa.2740680103>