



Cytogenetic data of *Partamona peckolti* (Hymenoptera, Apidae, Meliponini) by C banding and fluorochrome staining with DA/CMA₃ and DA/DAPI

Rute Magalhães Brito¹, Ana Paula de Almeida Caixeiro², Silvia das Graças Pompolo² and Giselle Garcia Azevedo³

¹Departamento de Biologia/Genética, USP, Cidade Universitária, São Paulo, SP, Brazil.

²Departamento de Biologia Geral, UFV, Viçosa, MG, Brazil.

³Setor de Entomologia, Faculdade de Filosofia Ciências e Letras de Ribeirão Preto, USP, Ribeirão Preto, SP, Brazil.

Abstract

The stingless bees of the *Partamona* genus have been studied taxonomically, ecologically and behaviourally, but cytogenetic studies are still rare. The objective of this study was to obtain cytogenetic data to contribute to *Partamona peckolti* species characterization. Heterochromatin was localized in all chromosome pericentromeric regions but some blocks could be visualized on some large chromosome arms. A large heterozygous DA-CMA₃-positive band was observed on one large chromosome arm, but was completely absent when C banding was applied before fluorochrome staining, with only one small positive band being visualized. Sequential DA-CMA₃-NOR staining of interphase nuclei provided coincident positive responses. This suggests that DA-CMA₃-positive bands of *P. peckolti* correspond to nucleolar organizer regions, as previously confirmed for another *Partamona* species by FISH.

Key words: Meliponini, *Partamona*, karyotype, C-banding, fluorochromes.

Received: May 31, 1999; accepted: December 4, 2002.

Introduction

The first karyotype evolution hypothesis elaborated for Meliponini bees postulated that the different chromosome numbers observed today are the product of polyploidy events from a basic number, $n = 8$ of *Melipona* species (Kerr *et al.*, 1952). Several Meliponini genus have been studied in detail at the cytogenetic level aiming at the comprehension of the karyotype evolution of stingless bees such as: *Leurotrigona* (Pompolo and Campos, 1995); *Frieseomelitta* (Moreira, 1997), *Plebeia* (Caixeiro and Pompolo, 1998) and *Melipona* (Rocha and Pompolo, 1998; Rocha *et al.*, 2002). These modern cytogenetic data have shown interespecific differences that cannot sustain the Polyploidy Hypothesis mentioned. Alternatively another karyotype evolution theory, the Minimum Interaction Theory (Imai *et al.* 1986, 1988, 1994), postulated for Australian ants, has also been accepted as a possible model for the karyotype evolution of Meliponini bees (Pompolo, 1992). This theory states that primitive karyotypes had small numbers of large chromosomes and, as time went by, these chromosomes got smaller and increased in number by

centric fission that would prevent deleterious interactions within the interphase nucleus, such as translocations.

Detailed study of each Meliponini genus at the cytogenetic level will provide the necessary data to verify if this Theory could also be applied to the stingless bees or whether it would be necessary to postulate another hypothesis to understand the karyotype evolution of this important group which is responsible for the pollination of 40 to 90% of wild plant species, depending on the ecosystem (Kerr, *et al.* 1996).

The objective of the present scientific note was to contribute to the knowledge of the *Partamona* genus at the cytogenetic level. According to Pedro (1998), 32 species of *Partamona* have been described so far, but few have been characterized cytogenetically yet: *P. personi* (Tarelho, 1973); *P. mulata*, *P. vicina*, *P. ayilae*, *Partamona* sp.n. (Brito-Ribon, *et al.*, 1999); and *Partamona* sp. aff. *nigrrior* (Brito, 1998), all presenting $2n = 34$ e $n = 17$ chromosomes. There is also *Partamona helleri* (Costa *et al.*, 1992, Brito *et al.* 1997, Tosta *et al.*, 1998) which presents $2n = 34$ chromosomes and a B chromosome system varying from 0 to 4.

In this short communication, we describe the karyotype of *Partamona peckolti*, a species that occurs in the tropical rainforests of central America, the Caribbean

and Pacific coasts, Colombia and Ecuador (Camargo, 1980), presenting the chromosome number, C banding pattern and fluorochrome AT and GC specific staining.

Material and Methods

The biological material was collected from a nest associated with a Bromeliacea plant, in San Francisco de Las Pampas village (00°25'182" S and 79°00'183" W), Cotopaxi province, Ecuador. Slides with metaphases were prepared from 30 post-defecant larvae cerebral ganglion according to the technique described by Imai *et al.* (1988). Chromosome banding techniques were applied as C banding according to Sumner (1972) and fluorochrome sequential staining DA - DAPI - Chromomycin A₃ (Schweizer, 1980). Sequential treatment, C banding followed by DA - DAPI - CMA₃ staining was done using the techniques in the same way as done separately. Another sequential staining with DA - CMA₃ followed by AgNOR treatment was also applied using the Schweizer (1980) protocol and silver impregnation as described by Howell and Black (1980). The metaphases were photographed in photomicroscope Olympus BX60, with FUJI HR II ISO 40 and KODAK GOLD ULTRA ISO 100 films. The karyotypes were mounted in increasing order of the chromosome length.

Results and Discussion

Partamona peckolti presented $2n = 34$ chromosomes as observed in other species of the genus (Tarelho, 1973; Brito 1998; Brito-Ribon *et al.*, 1999) and conventional

staining revealed a secondary constriction in the first pair of the karyotype not observed in any other *Partamona* species studied cytogenetically (Brito 1998, Brito-Ribon *et al.*, 1999) (Figure 1A).

The C-banding pattern showed heterochromatin in the pericentromeric region of all chromosomes, as already observed in *P. helleri* (Brito, 1998) and *Partamona* sp.n. (Brito-Ribon *et al.*, 1999) as well as a large heterochromatic block on the first karyotype pair, similarly to *P. aylaie*, *P. mulata* and *P. vicina* (Brito-Ribon *et al.*, 1999) (Table 1 and Figure 1B).

Sequential staining with DA-DAPI-CMA₃ revealed a wide positive band in one large chromosome with both fluorochromes (Figure 2A and B, arrows). A similar wide CMA₃-positive band has already been observed in other *Partamona* species (Table 1). However, this large band disappeared when C banding was applied before sequential staining with DA-DAPI-CMA₃ (Figure 3A, B and C). Another small DA/CMA₃⁺ band seen with or without the C banding treatment which presented the shape and richness in GC base pairs led us to believe that we were evidencing a nucleolar organizer region (NOR) (Figure 3 arrowhead). This correlation between CMA₃⁺ bands and NORs is quite common among animals and in insects has been observed in: the grasshoppers *Eyprepocnemis plorans* and *Locusta migratoria* (Camacho, *et al.*, 1991); the wasp *Trypoxylon albitarse* (Araújo *et al.*, 2000; 2002); the lady beetle *Olla v-nigrum* (Maffei *et al.*, 2001a); and in other Meliponini bees such as *Melipona* (Rocha and Pompolo, 1998; Rocha *et al.*, 2002); *Plebeia* (Caixeiro, 1999) and *Friesella*




Figure 1 - *Partamona peckolti* karyotypes mounted from metaphases submitted to: A) Conventional staining, and B) C banding. Bar = 5 μ m.




Figure 2 - *Partamona peckolti* metaphase submitted to sequential staining with: a) DA-DAPI, and B) CMA₃. The arrow points at a large DA-DAPI-CMA₃-positive band. Bar = 5 μm.

schrottkyi (Mampumbu, 2002). The location of these regions by the classical technique of Howell and Black (1980) is difficult on Meliponinae chromosomes that was effectively achieved only in *Tetragonisca angustula* (Menezes, 1997), *Melipona* (Maffei *et al.*, 2001b; Rocha *et al.*, 2002); *Plebeia* (Maffei *et al.*, 2001b) and *Friesella schrottkyi* (Mampumbu, 2002). Conversely, correspondence of DA-CMA₃ and AgNOR staining could be observed at interphase nuclei (Figure 4A and B). As the correlation among positive staining with DA-CMA₃; AgNOR and fluorescent in situ hybridization with 18S rDNA probe has been observed for two another *Partamona* species, *P. helleri* and *P. aff. nigrior* (Brito, 1988), we indeed accepted that structure (Figure 3C) as a NOR.






Figure 3 - *Partamona peckolti* metaphase submitted to C banding followed by sequential staining. A) DA-DAPI; B) CMA₃; C) GIEMSA. The arrow points at a NOR-like CMA₃-positive band. Bar = 5 μm.

Table 1 - Comparative data obtained for the *Partamona* genus with C-banding; DA/CMA₃⁺ staining and AgNO₃ impregnation. *: Not assayed.

<i>Partamona</i> species	Pericentromeric C ⁺ bands	1 st pair with broadly heterochromatic long arm	Majority of pairs with heterochromatic short arms	Large heteromorphic DA/CMA ₃ ⁺ band	Positive band correlation C ⁺ -DA/CMA ₃ ⁺	Positive nucleolus staining correlation DA/CMA ₃ ⁺ /AgNO ₃	References
<i>P. aff. nigrrior</i>	yes	no	no	no	no	yes	Brito (1998)
<i>P. ayilae</i>	no	yes	yes	no	*	*	Brito-Ribon (1999)
<i>P. helleri</i>	yes	no	no	yes	yes	yes	Brito (1998)
<i>P. mulata</i>	no	yes	yes	yes	*	*	Brito-Ribon (1999)
<i>P. peckolti</i>	yes	yes	no	yes	no	yes	Present work
<i>P. sp.n.</i>	yes	no	no	no	*	*	Brito-Ribon (1999)
<i>P. vicina</i>	no	yes	yes	yes	*	*	Brito-Ribon (1999)

**A****B****Figure 4** - Interphase nucleus submitted to: A) DA-CMA₃ staining, and B) AgNOR staining. The arrows point at coincident positive staining points. Bar = 5 μm.

Many studies have been carried out for a better understanding of the evolution of the *Partamona* genus: taxonomy revision (Pedro, 1998), provisioning and oviposition process behavior of nine species (Azevedo, 2001); molecular study of B chromosomes of *P. helleri* (Tosta, 2001); mi-

tochondrial genome characterization of *P. mulata* and *P. helleri* (Brito and Arias, 2001). In this context, Cytogenetics can still contribute greatly to the knowledge of the relations among *Partamona* species to further cytotaxonomy treatment of the data. Nevertheless, to achieve this purpose, more karyotype characterizations are needed as many *Partamona* species persist even without their chromosome number known. Furthermore, the better comprehension of the relations among *Partamona* species, especially at the cytogenetic level, will contribute to the comprehension of the karyotype evolution of the Meliponinae bees as a whole.

Acknowledgments

Research supported by Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), Fundação Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Conselho Nacional de Pesquisa (CNPq) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). We are grateful to Dr. Giovanni Onore (Universidad Católica Del Ecuador) for receiving Giselle A. Azevedo in Quito, Dr. Ronaldo Zucchi (Departamento de Biologia - FFCLRP) for encouragement, to MS Sidnei Mateus (Departamento de Biologia - FFCLRP), and to Mr. Cesar Tapi for help with nest collection.

References

- Araújo SMR, Pompolo SG, Dergam JAS and Campos, LAO (2000) The B chromosome system of *Trypoxylon (Trypargilum) albitarse* (Hymenoptera, Sphecidae). 1. Banding analysis. *Cytobios*, 101(1):7-13.
- Araújo SMR, Pompolo SG, Perfectti F and Camacho JPM (2002) Genetic load caused by variation in the amount of rDNA in a wasp. *Chromosome Res* 10:607-613.
- Azevedo GG (2001) Estudo etológico-comparado de alguns táxons de *Partamona* Schwarz e *Parapartamona* Schwarz (Hymenoptera, Apidae, Meliponini). PhD Thesis, Universidade de São Paulo, Ribeirão Preto.

- Brito RM, Costa MA and Pompolo SG (1997) Characterization and distribution of supernumerary chromosomes in 23 colonies of *Partamona helleri* (Hymenoptera, Apidae, Meliponinae). *Brazil J Genet* 20(2):185-188.
- Brito, RM (1998) Caracterização citogenética de duas espécies do gênero *Partamona*, Schwarz 1939 (Hymenoptera, Apidae, Meliponinae). MS Dissertation, Universidade Federal de Viçosa, Viçosa.
- Brito-Ribon RM, Miyazawa CS and Pompolo SG (1999) First Karyotype Characterization of four Species of *Partamona* (Friese, 1980) (Hymenoptera, Apidae, Meliponini) in Mato Grosso State - Brazil. *Cytobios* 100:19-26.
- Brito RM and Arias MC (2001) Caracterização parcial do genoma mitocondrial de duas espécies do gênero *Partamona* (Hymenoptera, Apidae, Meliponinae) por meio de RFLP. CD-ROM.47° Congresso Nacional de Genética, Águas de Lindóia, Brazil.
- Camacho JPM, Cabrero J and Viseras E (1991) G banding in two species of grasshoppers and its relationships to C, N, and fluorescence banding techniques. *Genome* 34:638-643.
- Caixeiro APA and Pompolo SG (1998) Análise citogenética da heterocromatina constitutiva de duas espécies do gênero *Plebeia* (Hymenoptera, Apidae, Meliponinae) por meio de bandamento C e fluorocromos. *Genet Mol Biol* 21(3):49.
- Caixeiro APA (1999) Caracterização citogenética da heterocromatina constitutiva e sua implicação na evolução do cariótipo de espécies do gênero *Plebeia* (Hymenoptera: Apidae, Meliponini). MS Dissertation, Universidade Federal de Viçosa, Viçosa.
- Costa MA, Pompolo SG and Campos LAO (1992) Supernumerary chromosomes in *Partamona cupira* (Hymenoptera, Apidae, Meliponinae). *Rev Brasil Genet* 15:801-806.
- Howell W and Black DA (1980) Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 36:1014-1015.
- Imai HT, Maruyama T, Gojobori T, Inoue Y and Crozier RH (1986) Theoretical bases for karyotype evolution. The minimum-interaction hypothesis. *Am Nat* 128:900-920.
- Imai HT, Taylor RW and Crozier RH (1988) Modes of spontaneous chromosomal mutation and karyotype evolution in ants with reference to the Minimum Interaction Hypothesis. *Jpn J Genet* 63:159-185.
- Imai HT, Taylor RW and Crozier RH (1994) Experimental bases for the minimum interaction theory. I. Chromosome evolution in ants of the *Myrmecia pilosula* species complex (Hymenoptera: Formicidae: Myrmecinae). *Jpn J Genet* 69:137-182.
- Kerr WE (1952) A variação do número de cromosomas na evolução dos Hymenoptera. *Estrato Scientia Genética* IV(3):182-190.
- Kerr WE, Carvalho GA and Nascimento VA (1996) Abelha urucu. *Biologia, manejo e conservação*. Acangaú, Belo Horizonte, 144 pp.
- Maffei EMD, Pompolo SG, Campos LAO and Petitpierre E (2001a) Sequential FISH analysis with rDNA genes and Ag-NOR banding in the lady beetle *Olla v-nigrum* (Coleoptera: Coccinellidae). *Hereditas* 135:13-18.
- Maffei EMD, Pompolo SG, Silva-Jr JC, Caixeiro APA, Rocha MP and Dergam JA (2001b) Silver staining of nucleolar organizer regions (NORs) in some species of Hymenoptera (bees and parasitic wasps) and Coleoptera (lady beetle). *Cytobios* 104:119-125.
- Mampumbu AR (2002) Análise citogenética da heterocromatina e da NOR em populações de abelhas sem ferrão *Friesella schrottkyi* (FRIESE, 1900) (Hymenoptera: Apidae: Meliponini). MS Dissertation, Unicamp, Campinas.
- Menezes MBF (1997) Caracterização citogenética e análise da heterocromatina constitutiva em *Tetragonisca angustula* (Hymenoptera, Meliponinae). MS Dissertation. Universidade Federal de Viçosa, Viçosa.
- Moreira CMLC (1997) Caracterização cariotípica de três espécies do gênero *Friesomelitta* (Hymenoptera, Apidae, Meliponinae). MS Dissertation. Universidade Federal de Viçosa, Viçosa.
- Pedro SRM (1998) Meliponini neotropicais: o gênero *Partamona* Schwarz, 1939 (Hymenoptera, Apidae): taxonomia e biogeografia. PhD Thesis. Universidade de São Paulo, Ribeirão Preto.
- Pompolo SG (1992) Estudos citogenéticos em Meliponinae. *Naturalia* (special ed.): 62-66.
- Pompolo SG and Campos LAO (1995) Karyotypes of two species of stingless bees, *Leurotrigona muelleri* and *Leurotrigona pusilla* (Hymenoptera, Meliponinae). *Rev Brasil Genet* 18(2):181-184.
- Rocha MP and Pompolo SG (1998) Karyotypes and heterochromatin variation (C-bands) in *Melipona* species (Hymenoptera, Apidae, Meliponinae). *Genet Mol Biol* 21(1):41-45.
- Rocha MP, Pompolo SG, Dergam JA, Fernandes A and Campos LAO (2002) DNA characterization and karyotypic evolution in the bee genus *Melipona* (Hymenoptera, Meliponini). *Hereditas* 136:19-27.
- Schweizer D (1980) Simultaneous fluorescent staining of R bands and specific heterochromatic regions (DA DAPI-bands) in human chromosomes. *Cytogenet Cell Genet* 27:190-193.
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Exp Cell Res* 75:304-306.
- Tarelho, ZVS (1973) Contribuição ao estudo citogenético dos Apoidea. MS Dissertation, Universidade de São Paulo, Ribeirão Preto.
- Tosta VC, Waldschmidt AM Brito, RM Pompolo SG, Salomão TMF, Barros EG and Campos LAO (1998) Marcadores moleculares RAPD ligados à presença de cromossomos supranumerários em *Partamona helleri* (Hymenoptera, Apidae). *Genet Mol Biol* 21, suppl.: 83. 44° Congresso Nacional de Genética, Águas de Lindóia, Brazil.
- Tosta VC, Salomão TMF, Tavares MG, Pompolo SG, Barros EG and Campos LAO (2001) Identificação, isolamento e clonagem de marcadores moleculares RAPD associados a cromossomos B em *Partamona helleri* (Hymenoptera, apidae, Meliponinae). CD-ROM. 47° Congresso Nacional de Genética, Águas de Lindóia, Brazil.