



## Evaluation of the mutagenicity and antimutagenicity of *Ziziphus joazeiro* Mart. bark in the micronucleus assay

Marcelo Fabiano Gomes Boriollo<sup>1,2</sup>, Marielly Reis Resende<sup>4</sup>, Thaísia Andrielle da Silva<sup>1,2</sup>, Juliana Yoshida Públio<sup>2</sup>, Luiz Silva Souza<sup>1,2</sup>, Carlos Tadeu dos Santos Dias<sup>4</sup>, Nelma de Mello Silva Oliveira<sup>1,3</sup> and João Evangelista Fiorini<sup>1,2</sup>

<sup>1</sup>Laboratório de Farmacogenômica e Biologia Molecular, Faculdade de Ciências Médicas, Universidade José do Rosário Vellano, Alfenas, MG, Brazil.

<sup>2</sup>Centro de Pesquisa e Pós-Graduação em Ciência Animal, Área de Patologia e Farmacologia Animal, Universidade José do Rosário Vellano, Alfenas, MG, Brazil.

<sup>3</sup>Laboratório de Ecotoxicologia e Microbiologia Ambiental, Faculdade de Tecnologia, Universidade Estadual de Campinas, Limeira, SP, Brazil.

<sup>4</sup>Departamento de Ciências Exatas, Escola de Agricultura “Luiz de Queiroz”, Universidade de São Paulo, Piracicaba, SP, Brazil.

### Abstract

The aim of this study was to evaluate the mutagenicity (clastogenicity/aneugenicity) of a glycolic extract of *Ziziphus joazeiro* bark (GEZJ) by the micronucleus assay in mice bone marrow. Antimutagenic activity was also assessed using treatments associated with GEZJ and doxorubicin (DXR). Mice were evaluated 24-48 h after exposure to positive (N-nitroso-N-ethylurea, NEU - 50 mg.kg<sup>-1</sup> and DXR - 5 mg.kg<sup>-1</sup>) and negative (150 mM NaCl) controls, as well as treatment with GEZJ (0.5-2 g.kg<sup>-1</sup>), GEZJ (2 g.kg<sup>-1</sup>) + NEU and GEZJ (2 g.kg<sup>-1</sup>) + DXR. There were no significant differences in the frequencies of micronucleated polychromatic erythrocytes in mice treated with GEZJ and GEZJ + DXR compared to the negative controls, indicating that GEZJ was not mutagenic. Analysis of the polychromatic:normochromatic erythrocyte ratio revealed significant differences in the responses to doses of 0.5 g.kg<sup>-1</sup> and 1-2 g.kg<sup>-1</sup> and the positive control (NEU). These results indicated no systemic toxicity and moderate toxicity at lower and higher doses of GEZJ. The lack of mutagenicity and systemic toxicity in the antimutagenic assays, especially for treatment with GEZJ + DXR, suggested that phytochemical compounds in *Z. joazeiro* bark attenuated DXR-induced mutagenicity and the moderate systemic toxicity of a high dose of *Z. joazeiro* bark (2 g.kg<sup>-1</sup>). Further studies on the genotoxicity of *Z. joazeiro* extracts are necessary to establish the possible health risk in humans and to determine the potential as a chemopreventive agent for therapeutic use.

**Keywords:** antimutagenicity, bone marrow, doxorubicin, micronucleus assay, mutagenicity, *Ziziphus joazeiro* Mart. (raspa-de-Juá).

Received: October 3, 2013; Accepted: March 17, 2014.

### Introduction

Many species of medicinal plants, such as *Amburana cearensis*, *Anadenanthera colubrina*, *Mentha x villosa*, *Myracrodruon urundeuva*, *Plectranthus amboinicus*, *Ruta graveolens*, *Ximения americana* and *Ziziphus joazeiro*, are widely used by communities in the Brazilian Caatinga to treat a large spectrum of clinical conditions ranging from diseases requiring palliative care to general aches, e.g., bronchitis, sinusitis, rhinitis, nasal congestion, headaches,

flu, fever, expectorant, colic, hypertension, thrombosis, indigestion, intestinal dysfunction, liver and kidney problems, infectious and inflammatory processes and pain in general (Cartaxo *et al.*, 2010). *Ziziphus joazeiro* Mart. (Rhamnaceae) is a native Brazilian tree resistant to dry environments (Cartaxo *et al.*, 2010). This species is an important source of water and food for animals in arid habitats (Braga, 1960; Cruz, 1985; Nunes *et al.*, 1987).

A phytochemical analysis of *Z. joazeiro* Mart. has shown that the leaf epicuticular wax is rich in *n*-alkanes (78.6%), very efficient compounds for impermeabilizing the leaf surface, and triterpenoids (Oliveira *et al.*, 2003). A similar analysis of a dichloromethane extract of *Z. joazeiro* Mart. bark identified triterpenoids with weak antibacterial activity (e.g., betulinic, aliphatic and ursolic acids) and re-

Send correspondence to Marcelo Fabiano Gomes Boriollo. Laboratório de Farmacogenômica e Biologia Molecular, Faculdade de Ciências Médicas & Centro de Pesquisa e Pós-Graduação, Universidade José do Rosário Vellano, Rodovia MG 179, km 0, Campus Universitário, 37130-000 Alfenas, MG, Brazil. E-mail: marcelo.boriollo@unifenas.br.

markable activity against *Staphylococcus epidermidis* [e.g., betulinic acid ester derivatives such as 7 $\beta$ -(4-hydroxy-benzoyloxy), 7 $\beta$ -(4-hydroxy-3-methoxy-benzoyloxy) and 27-(4-hydroxy-3-methoxy-benzoyloxy)] (Schuhly *et al.*, 1999). *Ziziphus joazeiro* Mart. bark also contains an abundance of saponins that have been used as toothpastes, with aqueous extracts showing antimicrobial action against bacteria (planktonic cells and artificial biofilms) related to dental caries and periodontal diseases (Alviano *et al.*, 2008). Other popular therapeutic applications of *Z. joazeiro* Mart. include the treatment of dandruff, rheumatism, flu, fever, chronic bronchitis, gastric ulcers, indigestion, heartburn and headaches (Schuhly *et al.*, 1999; Cartaxo *et al.*, 2010). In addition, experimental studies have identified potential antifungal (Cruz *et al.*, 2007), antibacterial (Schuhly *et al.*, 1999; Alviano *et al.*, 2008; Leal *et al.*, 2010), antioxidant (Alviano *et al.*, 2008) and antipyretic (Nunes *et al.*, 1987) activities, as well as low toxicity (Alviano *et al.*, 2008).

Biologically active compounds have been recognized for their pharmacological properties, but many of them are of limited therapeutic use because of their toxicological, carcinogenic and mutagenic properties (Ames, 1983; Konstantopoulou *et al.*, 1992; Tavares, 1996). The analysis of genotoxicity is a major aspect of drug development since most pharmaceutical companies evaluate the potential of a new therapeutic agent based on its genotoxicity *in vitro* and *in vivo* (Purves *et al.*, 1995). In this context, the screening of popularly used plants and their isolated components for mutagenic activity is necessary and important for establishing adequate control measures. This screening can also provide insights into the mechanisms involved in the biological effects of plants used as therapeutic agents (Varanda, 2006).

As far as genotoxicity studies are concerned, the *in vivo* micronucleus (MN) assay in rodent bone marrow is a crucial part of the battery of tests used to identify hazardous mutagens (Mateuca *et al.*, 2006). This assay is especially suited for assessing mutagenic hazards because it contemplates various factors, such as *in vivo* metabolism, pharmacokinetics and DNA repair mechanisms, even though these processes vary among species and tissues and have different genetic endpoints (OECD, 1997a,b; Ribeiro *et al.*, 2003). Since bone marrow erythroblasts develop into polychromatic erythrocytes (PCEs), *i.e.*, cells generated by extrusion of the main nucleus, micronuclei may remain in an otherwise anucleated cytoplasm. Consequently, the frequency of micronucleated polychromatic erythrocytes (MNPCEs) has been the principal endpoint for MN assays. The measurement of MNPCEs in peripheral blood is possible in any species in which the spleen does not remove micronucleated erythrocytes, or that is sufficiently sensitive to agents that cause structural or numerical chromosomal aberrations. An increase in the frequency of MNPCEs in treated animals, *i.e.*, a positive result, indicates

that a substance can cause the formation of micronuclei through chromosomal damage or damage to the mitotic apparatus of erythroblasts. On the other hand, a negative result implies that the test substance does not cause micronucleus formation in immature erythrocytes. The number of normochromatic erythrocytes (NCEs) in peripheral blood that contain micronuclei for a given number of mature erythrocytes can also be used as the endpoint of this assay (OECD, 1997c; Ribeiro *et al.*, 2003). Several studies have used the mammalian *in vivo* MN assay to understand the mutagenic effects induced by phytotherapeutics and foods (Indart *et al.*, 2007; Venkatesh *et al.*, 2007; Chandrasekaran *et al.*, 2011; Silva *et al.*, 2011; Alves *et al.*, 2012).

Although several studies have examined the potential therapeutic effectiveness of *Z. joazeiro* Mart., there has been no systematic investigation of the genotoxic and mutagenic effects of this plant. In this work, we examined the mutagenic effects of a glycolic extract of *Z. joazeiro* Mart. bark as part of a wider study on the genotoxic potential of herbal extracts. The effect of the maximum permissible concentration of *Z. joazeiro* Mart. on the mutagenicity of doxorubicin (DXR) in mouse bone marrow, *i.e.*, its antimutagenic activity, was also examined.

## Material and Methods

### Raw material and sample preparation

A glycolic extract of *Z. joazeiro* bark (GEZJ) was purchased commercially and stored according to the manufacturer's recommendations (AKSY Comercial Ltda., São Bernardo do Campo, SP, Brazil). Aliquots (1.5 L) of this extract were submitted to solvent removal proceedings by rotary evaporation (40 rpm) (Rotavapor model R-215) coupled to a bath heating system maintained at 50-60 °C (Bath Heating model B-491), a vacuum pump (vacuum of 500 mm Hg; Vacuum Pump V-700 with Automatic Vacuum Controller V-855), a water recirculator (Recirculator Chiller F-100) and an evaporation bottle (Büchi Labor-technik AG, Switzerland). The final product was transferred to a 1 L reaction bottle (SCHOTT® DURAN®) and kept at -20 °C for 24 h in order to evaluate the freezing of the final product and the efficacy of solvent evaporation (Agência Nacional de Vigilância Sanitária (ANVISA) (2010)). Aliquots (40 mL) of this final product were transferred to penicillin-type glass vials (50 mL) and lyophilized (Lyophilizer model Alpha 1-2 LDPlus, Martin Christ Gefriertrocknungsanlagen GmbH, Germany) and the dry mass were measured (Electronic Analytical Balance AUW-220D, Shimadzu Corp., Kyoto, Japan). Aqueous solutions of the lyophilized product were prepared in type 1 water at twice the final concentration, sterilized by filtration (Millipore Corporation, hydrophilic Durapore® PVDF, 0.22  $\mu$ m,  $\pm$  47 mm, cat. no. GVWP 047 00) and stored in sterile polypropylene tubes (50 mL) at -70 °C until used.

### In vivo assays

Healthy, heterogeneous, young adult male and female Swiss mice (Unib:SW) 7-12 weeks old (pubescent period) weighing 30-40 g (weight variation among mice of each sex was < 20% of the mean weight) were provided by CEMIB (Centro Multidisciplinar para Investigação Biológica - UNICAMP) and erythrocytes from the bone marrow of these mice were used in the MN assay (Collaborative Study Group for the Micronucleus Test (CSGMT), 1986; Chorilli *et al.*, 2007).

Animals of the same sex were housed in polypropylene boxes in an air-conditioned environment to  $22 \pm 3$  °C, with a relative air humidity of  $50\% \pm 20\%$  and a 12 h light/dark cycle. The mice were fed commercial rodent chow (Purina® Labina, Nestlé Purina Pet Care Company) and water *ad libitum*, and were acclimated to laboratory conditions for seven days prior to use in the experiments. At the end of this period, each mouse was weighed and then received 2 mL of liquid (containing the desired test agent) per 100 g body weight.

All animals were properly identified by numerical markings on their tails to ensure continuity of the records and reliable interpretation of the results throughout the study (OECD, 1997c). After the period of treatment, the mice were euthanized by inhalation of carbon dioxide in adapted acrylic chambers as described in the Report of the American Veterinary Medical Association panel on euthanasia (Beaver *et al.*, 2000). This study was done in accordance with the Universal Declaration of Animal Rights (UNESCO, 1978), the ethical principles for animal experimentation established by the Brazilian Society of Labora-


tory Animal Science (SBCAL - *Sociedade Brasileira de Ciência em Animais de Laboratório*), the Brazilian Environmental Crimes Law (Law no. 9.605, February 12, 1998), the Brazilian standards for Didactic-Scientific Practice of Vivisection of Animals (Law no. 6.638, May 8, 1979), and was approved by the Committee for Ethics in Research Involving Animals at UNIFENAS (CEPEAU Protocol no. 04A/2008).

### Experimental groups

The experimental groups of mice (3 males and 3 females each) were assessed 24 h and 48 h after a single treatment administered by gavage (Figure 1). The mutagenic activity of GEZJ was assessed in mice that received doses of 0.5-2 g.kg<sup>-1</sup> (groups 7-14) and the antimutagenic activity was assessed in mice that received NEU (50 mg.kg<sup>-1</sup>) + GEZJ (2 g.kg<sup>-1</sup>) (groups 15 and 16) and DXR (5 mg.kg<sup>-1</sup>) + GEZJ (2 g.kg<sup>-1</sup>) (groups 17 and 18). The doses of GEZJ were chosen based on previous acute toxicity experiments in mice that yielded LD<sub>50</sub> values of 2.0-3.5 g/kg for several plant extracts, including *Z. joazeiro* (Alviano *et al.*, 2008). Negative controls (groups 1 and 2: 150 mM NaCl in type 1 water) and positive controls (groups 3 and 4: 50 mg.kg<sup>-1</sup> of NEU; groups 5 and 6: 5 mg.kg<sup>-1</sup> of DXR) were also included as single treatments administered by gavage (NaCl) and intraperitoneally (NEU and DXR) (OECD, 1997c).

### Processing of bone marrow

MN assays using bone marrow erythrocytes were done 24 h and 48 h after treatment, using previously described methodology (Schmid, 1976; Zambrano *et al.*,



**Figure 1** - Experimental protocol for assessing the mutagenic and antimutagenic activity of a glycolic extract of *Z. joazeiro* bark. T - treatment, E - euthanasia and *n* - number of mice.



1982). Shortly after euthanasia, the femora were surgically and aseptically removed and the mice were appropriately discarded. Each femur was sectioned at the proximal end and the contents of the spinal canal were washed with 1.5 mL of 150 mM NaCl and transferred to a 15 mL centrifuge tube. This material was resuspended with a Pasteur pipette to ensure a homogenous distribution of bone marrow cells. The suspension was then centrifuged at 1,000 rpm (Bench centrifuge, model NT 810, Nova Técnica Ind. e Com. de Equip. para Laboratório Ltda., Piracicaba, SP, Brazil) for 5 min. The supernatant was discarded and the resulting pellet was resuspended in 500  $\mu$ L of 150 mM NaCl solution added 4% formaldehyde. The slides (two per animal) were prepared by smearing, dried at room temperature for 24 h and stained with Leishman's eosin methylene blue dye [pure dye for 3 min followed by diluted dye in distilled water (1:6) for 15 min] to differentiate polychromatic erythrocytes (PCEs) from monochromatic erythrocytes (NCEs).

PCEs were observed by light microscopy (Nikon Eclipse E-200 microscope) at a magnification of 1000x, counted (at least 2000 anucleated polychromatic erythrocytes per animal) with the aid of a digital cell counter (Contador Diferencial CCS02, Kacil Indústria e Comércio Ltda., PE, Brazil) and photographed using an 8.1 Megapixel Digital Camera (DC FWL 150). The number of PCEs, the number and frequency of MNPCEs and the ratio of polychromatic to monochromatic erythrocytes (PCE/NCE) were determined.

### Statistical analysis

The data from the MN assay were analyzed by one-way analysis of variance (ANOVA) using a 9 x 2 x 2 (treatment x gender x time) factorial scheme followed by multiple comparisons with the Tukey test ( $\alpha = 0.05$ ). All analyses were done using SAS<sup>®</sup> version 9.2 computer software.

### Results and Discussion

*Ziziphus joazeiro* Mart. has been popularly used to treat dandruff, rheumatism, flu, fever, chronic bronchitis, gastric ulcers, indigestion, heartburn and headaches and to clean teeth (Schuhly *et al.*, 1999; Cartaxo *et al.*, 2010). In addition, *Z. joazeiro* has potential antifungal (Cruz *et al.*, 2007), antibacterial (Schuhly *et al.*, 1999; Alviano *et al.*, 2008; Leal *et al.*, 2010), antioxidant (Alviano *et al.*, 2008) and antipyretic (Nunes *et al.*, 1987) activities, as well as low toxicity (Alviano *et al.*, 2008). This information partly supports the popular use of *Z. joazeiro* for certain treatments and agrees with ethnopharmacological studies designed to select plants for bioactivity screening (Cruz *et al.*, 2007). In contrast, few studies have examined the mutagenic and antimutagenic effects of *Z. joazeiro* Mart.

In the present study, the number and frequency of MNPCEs and the PCE/NCE ratios in mouse bone marrow were analyzed in mutagenic and antimutagenic assays of a glycolic extract of *Z. joazeiro* bark (Table 1 and Figure 2). Analysis of the MNPCEs revealed no significant differences between the 24 h and 48 h results for the negative (NaCl) and positive (DXR and NEU) controls. However, there were significant differences ( $p < 0.05$ ) between the negative and positive controls at the two time intervals. There were no differences between the negative controls and the treatments with GEJZ (0.5-2 g.kg<sup>-1</sup>) or with GEJZ (2 g.kg<sup>-1</sup>) + DXR (5 mg.kg<sup>-1</sup>): these responses showed no dose or time dependence, but varied between male and female mice. Mice treated with GEJZ (2 g.kg<sup>-1</sup>) + NEU (50 mg.kg<sup>-1</sup>) had intermediate values ( $n$  and %) that differed significantly from the negative and positive controls. These results suggest absence of mutagenicity (clastogenicity and/or aneugenicity) for GEJZ, regardless of the extract dose and time interval, although the responses varied between sexes. In contrast, GEJZ (2 g.kg<sup>-1</sup>) showed antimutagenic activity (anticlastogeny and/or antianeugeny) towards the chemotherapeutic agent DXR (5 mg.kg<sup>-1</sup>) or NEU (50 mg.kg<sup>-1</sup>), regardless of the time interval, although once again intersex variation was observed. These findings indicate that compounds in GEJZ can act against DXR-induced mutagenic effects in mouse bone marrow. Such compounds could include  $n$ -alkanes, triterpenoids [*i.e.*, betulinic acid, aliphatic acid, ursolic acid, ester derivatives of betulinic acid such as 7 $\beta$ -(4-hydroxy-benzoyloxy)-betulinic acid, 7 $\beta$ -(4-hydroxy-3-methoxy-benzoyloxy)-betulinic acid and 27-(4-hydroxy-3-methoxy-benzoyloxy)-betulinic acid] (Oliveira *et al.*, 2003; Schuhly *et al.*, 1999) and saponins (Alviano *et al.*, 2008). DXR has been reported to induce micronuclei, chromatid and chromosomal aberrations, and DNA single- and double-strand breaks *in vitro* and *in vivo* (Bean *et al.*, 1992; Al-Harbi, 1993; Al-Shabanah, 1993; Delvaeye *et al.*, 1993; Jagetia and Nayak, 1996, 2000; Shan *et al.*, 1996; Dhawan *et al.*, 2003; Jagetia and Aruna, 2000). In addition, the major acute toxicity induced by DXR is bone marrow suppression, while the long-term clinical usefulness is limited by a cumulative, dose-dependent, irreversible, chronic cardiotoxicity that manifests itself as congestive heart failure or cardiomyopathy (Van Acker *et al.*, 1995, 2000).

For the PCE/NCE ratio, there were no significant differences between the negative controls (NaCl), the positive control DXR (5 mg.kg<sup>-1</sup>), the GEJZ (0.5 mg.kg<sup>-1</sup>) group, and mice treated with GEJZ (2 g.kg<sup>-1</sup>) + NEU (50 mg.kg<sup>-1</sup>) or with GEJZ (2 g.kg<sup>-1</sup>) + DXR (5 mg.kg<sup>-1</sup>) (Table 1 and Figure 1). For the treatments with GEJZ, there was a significant difference between the dose of 500 mg.kg<sup>-1</sup> and the doses of 1.5 g.kg<sup>-1</sup> and 2 g.kg<sup>-1</sup>. Although there were no significant intersex differences, the responses did vary with time (24 h vs. 48 h). Lower doses of GEJZ (0.5-1 g.kg<sup>-1</sup>) were not toxic to bone marrow compared to higher doses

**Table 1** - MNPCE frequencies and PCE/NCE ratios in mouse bone marrow in mutagenic and antimutagenic assays of *Z. joazeiro* bark.

Treatment	Number of PCEs analyzed		MNPCEs*				PCE / (PCE + NCE)**		NCE (n)	
	24 h	48 h	24 h (n) <sup>A</sup>	48 h (n) <sup>A</sup>	24 h (%) <sup>A</sup>	48 h (%) <sup>A</sup>	24 h <sup>A</sup>	48 h <sup>B</sup>	24 h	48 h
150 mM NaCl										
F <sub>1</sub>	2095	2097	7	10	0.33	0.48	1.00	1.00	5	3
F <sub>2</sub>	2094	2095	9	10	0.43	0.48	1.00	1.00	6	5
F <sub>3</sub>	2087	2089	11	8	0.53	0.38	0.99	0.99	13	11
Σ F	Σ 6276	Σ 6281	Σ 27	Σ 28	0.43 ± 0.10	0.45 ± 0.05	1.00 ± 0.00	1.00 ± 0.00	Σ 24	Σ 19
M <sub>1</sub>	2095	2088	9	13	0.43	0.62	1.00	0.99	5	12
M <sub>2</sub>	2055	2088	12	11	0.58	0.53	0.98	0.99	45	12
M <sub>3</sub>	2058	2084	7	11	0.34	0.53	0.98	0.99	42	16
Σ M	Σ 6208	Σ 6260	Σ 28	Σ 35	0.45 ± 0.12	0.56 ± 0.06	0.99 ± 0.01	0.99 ± 0.00	Σ 92	Σ 40
Σ	Σ 12484	Σ 12541	Σ 55 <sup>A</sup>	Σ 63 <sup>A</sup>	0.44 ± 0.08 <sup>A</sup>	0.50 ± 0.06 <sup>A</sup>	0.99 ± 0.01 <sup>A</sup>	1.00 ± 0.00 <sup>A</sup>	Σ 116	Σ 59
N-Nitroso-N-ethylurea (NEU, 50 mg.kg <sup>-1</sup> )										
F <sub>1</sub>	2148	2075	38	36	1.77	1.73	0.49	0.65	2252	1125
F <sub>2</sub>	1884	2032	32	34	1.70	1.67	0.54	0.81	1616	468
F <sub>3</sub>	2002	1948	15	31	0.75	1.59	0.61	0.93	1298	152
Σ F	Σ 6034	Σ 6055	Σ 85	Σ 101	1.41 ± 0.57	1.67 ± 0.07	0.54 ± 0.06	0.80 ± 0.14	Σ 5166	Σ 1745
M <sub>1</sub>	2025	1999	64	31	3.16	1.55	0.41	0.36	2875	3501
M <sub>2</sub>	2028	1916	105	40	5.18	2.09	0.51	0.55	1972	1584
M <sub>3</sub>	2004	2069	25	38	1.25	1.84	0.67	0.65	996	1131
Σ M	Σ 6057	Σ 5984	Σ 194	Σ 109	3.20 ± 1.97	1.83 ± 0.27	0.53 ± 0.13	0.52 ± 0.14	Σ 5843	Σ 6216
Σ M and F	Σ 12091	Σ 12039	Σ 279 <sup>C</sup>	Σ 210 <sup>C</sup>	2.30 ± 1.66 <sup>C</sup>	1.75 ± 0.18 <sup>C</sup>	0.54 ± 0.06 <sup>D</sup>	0.66 ± 0.16 <sup>D</sup>	Σ 11009	Σ 7961
Doxorubicin hydrochloride (DXR, 5 mg.kg <sup>-1</sup> )										
F <sub>1</sub>	2091	2017	49	36	2.34	1.78	0.72	0.96	809	83
F <sub>2</sub>	2106	2077	73	63	3.47	3.03	0.98	0.99	44	23
F <sub>3</sub>	2056	2092	57	50	2.77	2.39	0.84	0.95	394	108
Σ F	Σ 6253	Σ 6186	Σ 179	Σ 149	2.86 ± 0.57	2.40 ± 0.62	0.85 ± 0.13	0.97 ± 0.02	1247	214
M <sub>1</sub>	2067	2086	53	61	2.56	2.92	0.98	0.95	33	114
M <sub>2</sub>	2063	2042	56	70	2.71	3.43	0.98	0.97	37	58
M <sub>3</sub>	2082	2075	46	50	2.21	2.41	0.99	0.99	18	25
Σ M	Σ 6212	Σ 6203	Σ 155	Σ 181	2.50 ± 0.26	2.92 ± 0.51	0.99 ± 0.00	0.97 ± 0.02	88	197
Σ M and F	Σ 12465	Σ 12389	Σ 334 <sup>D</sup>	Σ 330 <sup>D</sup>	2.68 ± 0.42 <sup>D</sup>	2.66 ± 0.43 <sup>D</sup>	0.92 ± 0.07 <sup>AB</sup>	0.97 ± 0.01 <sup>AB</sup>	1335	411
Glycolic extract of <i>Z. joazeiro</i> Mart. bark (0.5 mg.kg <sup>-1</sup> )										
F <sub>1</sub>	2051	2085	8	8	0.39	0.38	0.77	0.95	620	115

**Table 1 (cont.)**

Treatment	Number of PCEs analyzed			MNPCEs*			PCE / (PCE + NCE)**			NCE (n)		
	24 h	48 h	Σ	24 h (n) <sup>A</sup>	48 h (n) <sup>A</sup>	Σ	24 h (%) <sup>A</sup>	48 h (%) <sup>A</sup>	24 h <sup>A</sup>	48 h <sup>B</sup>	24 h	48 h
F <sub>2</sub>	2082	2035		4	8		0.19	0.39	0.94	0.97	129	65
F <sub>3</sub>	2083	2005		11	2		0.53	0.10	0.94	0.95	134	95
Σ F <sup>A</sup> , A**	Σ 6216	Σ 6125		Σ 23	Σ 18		0.37 ± 0.17	0.29 ± 0.17	0.88 ± 0.10	0.96 ± 0.01	Σ 883	Σ 275
M <sub>1</sub>	2076	2060		9	7		0.43	0.34	0.99	0.98	24	40
M <sub>2</sub>	2002	2046		7	7		0.35	0.34	0.95	0.97	98	54
M <sub>3</sub>	2038	2071		7	17		0.34	0.82	0.97	0.99	62	29
Σ M <sup>B</sup> , A**	Σ 6116	Σ 6177		Σ 23	Σ 31		0.38 ± 0.05	0.50 ± 0.28	0.97 ± 0.02	0.98 ± 0.01	Σ 184	Σ 123
Σ M and F	Σ 12332	Σ 12302		Σ 46 <sup>A</sup>	Σ 49 <sup>A</sup>		0.37 ± 0.00 <sup>A</sup>	0.40 ± 0.15 <sup>A</sup>	0.92 ± 0.07 <sup>AB</sup>	0.97 ± 0.02 <sup>AB</sup>	Σ 1067	Σ 398
Glycolic extract of <i>Z. joazeiro</i> Mart. bark (1 g.kg <sup>-1</sup> )												
F <sub>1</sub>	2062	2047		9	10		0.44	0.49	0.80	0.89	531	253
F <sub>2</sub>	2044	2053		10	9		0.49	0.44	0.91	0.93	190	147
F <sub>3</sub>	2023	2067		11	8		0.54	0.39	0.86	0.94	337	138
Σ F <sup>A</sup> , A**	Σ 6129	Σ 6167		Σ 30	Σ 27		0.49 ± 0.05	0.44 ± 0.05	0.86 ± 0.06	0.92 ± 0.03	Σ 1058	Σ 538
M <sub>1</sub>	2073	2014		15	12		0.72	0.60	0.81	0.92	489	175
M <sub>2</sub>	2056	2032		3	13		0.15	0.64	0.98	0.95	37	110
M <sub>3</sub>	2101	2075		20	16		0.95	0.77	0.84	0.97	389	68
Σ M <sup>B</sup> , A**	Σ 6230	Σ 6121		Σ 38	Σ 41		0.61 ± 0.42	0.67 ± 0.09	0.88 ± 0.09	0.95 ± 0.02	Σ 915	Σ 353
Σ M and F	Σ 12359	Σ 12288		Σ 68 <sup>A</sup>	Σ 68 <sup>A</sup>		0.55 ± 0.09 <sup>A</sup>	0.55 ± 0.16 <sup>A</sup>	0.86 ± 0.01 <sup>BC</sup>	0.93 ± 0.02 <sup>BC</sup>	Σ 1973	Σ 891
Glycolic extract of <i>Z. joazeiro</i> Mart. bark (1.5 mg.kg <sup>-1</sup> )												
F <sub>1</sub>	2082	2041		21	16		1.01	0.78	0.80	0.84	518	401
F <sub>2</sub>	2134	2107		18	15		0.84	0.71	0.89	0.80	266	528
F <sub>3</sub>	2095	2075		16	14		0.76	0.67	0.81	0.91	505	195
Σ F <sup>A</sup> , A**	Σ 6311	Σ 6223		Σ 55	Σ 45		0.87 ± 0.12	0.72 ± 0.06	0.83 ± 0.05	0.85 ± 0.06	Σ 1289	Σ 1124
M <sub>1</sub>	2045	2075		12	27		0.59	1.30	0.82	0.96	455	84
M <sub>2</sub>	2125	2048		14	16		0.66	0.78	0.86	0.92	352	173
M <sub>3</sub>	2108	2022		16	8		0.76	0.40	0.85	0.69	374	928
Σ M <sup>B</sup> , A**	Σ 6278	Σ 6145		Σ 42	Σ 51		0.67 ± 0.09	0.83 ± 0.45	0.84 ± 0.02	0.84 ± 0.15	Σ 1181	Σ 1185
Σ M and F	Σ 12589	Σ 12368		Σ 97 <sup>AB</sup>	Σ 96 <sup>AB</sup>		0.77 ± 0.14 <sup>AB</sup>	0.78 ± 0.08 <sup>AB</sup>	0.84 ± 0.01 <sup>C</sup>	0.84 ± 0.01 <sup>C</sup>	Σ 2470	Σ 2309
Glycolic extract of <i>Z. joazeiro</i> Mart. bark (2 g.kg <sup>-1</sup> )												
F <sub>1</sub>	2032	2162		21	14		1.03	0.65	0.69	0.85	922	388
F <sub>2</sub>	2173	2037		17	23		0.78	1.13	0.93	0.79	176	532
F <sub>3</sub>	2020	2070		18	18		0.89	0.87	0.84	0.85	387	378


Table 1 (cont.)

Treatment	Number of PCEs analyzed			MNPCEs*			PCE / (PCE + NCE)**			NCE (n)	
	24 h	48 h	24 h (n) <sup>A</sup>	48 h (n) <sup>A</sup>	24 h (%) <sup>A'</sup>	48 h (%) <sup>A'</sup>	24 h <sup>A''</sup>	48 h <sup>B''</sup>	24 h	48 h	
Σ F <sup>A, A**</sup>	Σ 6225	Σ 6269	Σ 56	Σ 55	0.90 ± 0.13	0.88 ± 0.24	0.82 ± 0.12	0.83 ± 0.03	Σ 1485	Σ 1298	
M <sub>1</sub>	2033	2010	17	14	0.84	0.70	0.84	0.77	383	590	
M <sub>2</sub>	2058	2056	18	17	0.87	0.83	0.79	0.86	542	344	
M <sub>3</sub>	2020	2037	10	26	0.50	1.28	0.70	0.88	880	277	
Σ M <sup>B*, A**</sup>	Σ 6111	Σ 6103	Σ 45	Σ 57	0.74 ± 0.21	0.93 ± 0.30	0.78 ± 0.07	0.83 ± 0.06	Σ 1805	Σ 1211	
Σ M and F	Σ 12336	Σ 12372	Σ 101 <sup>AB</sup>	Σ 112 <sup>AB</sup>	0.82 ± 0.12 <sup>AB'</sup>	0.91 ± 0.04 <sup>AB'</sup>	0.79 ± 0.03 <sup>C''</sup>	0.83 ± 0.00 <sup>C''</sup>	Σ 3290	Σ 2509	
Glycolic extract of <i>Z. joazeiro</i> Mart. bark (2 g.kg <sup>-1</sup> ) + NEU (50 mg.kg <sup>-1</sup> )											
F <sub>1</sub>	2052	2079	31	24	1.51	1.15	0.93	0.99	148	21	
F <sub>2</sub>	2072	2055	26	16	1.25	0.78	0.99	0.98	28	45	
F <sub>3</sub>	2071	2167	23	16	1.11	0.74	0.99	0.99	29	33	
Σ F <sup>A, A**</sup>	Σ 6195	Σ 6301	Σ 80	Σ 56	1.29 ± 0.20	0.89 ± 0.23	0.97 ± 0.03	0.98 ± 0.01	Σ 205	Σ 99	
M <sub>1</sub>	2138	2241	32	43	1.50	1.92	0.97	0.90	62	259	
M <sub>2</sub>	2144	2103	29	28	1.35	1.33	0.97	0.73	56	797	
M <sub>3</sub>	2072	2076	32	27	1.54	1.30	0.94	0.99	128	24	
Σ M <sup>B*, A**</sup>	Σ 6354	Σ 6420	Σ 93	Σ 98	1.46 ± 0.10	1.53 ± 0.35	0.96 ± 0.02	0.86 ± 0.13	Σ 246	Σ 1080	
Σ M and F	Σ 12549	Σ 12721	Σ 173 <sup>B</sup>	Σ 154 <sup>B</sup>	1.38 <sup>B'</sup>	1.21 <sup>B'</sup>	0.97 ± 0.00 <sup>AB''</sup>	0.92 ± 0.09 <sup>AB''</sup>	Σ 451	Σ 1179	
Glycolic extract of <i>Z. joazeiro</i> Mart. bark (2 g.kg <sup>-1</sup> ) + DXR (5 mg.kg <sup>-1</sup> )											
F <sub>1</sub>	2086	2090	23	18	1.10	0.86	0.99	0.99	14	10	
F <sub>2</sub>	2080	2100	23	20	1.11	0.95	0.99	1.00	20	1	
F <sub>3</sub>	2080	2075	21	17	1.01	0.82	0.99	0.99	20	24	
Σ F <sup>A, A**</sup>	Σ 6246	Σ 6265	Σ 67	Σ 55	1.07 ± 0.05	0.88 ± 0.07	0.99 ± 0.00	0.99 ± 0.01	Σ 54	Σ 35	
M <sub>1</sub>	2086	2088	24	12	1.15	0.57	0.99	0.99	14	12	
M <sub>2</sub>	2083	2076	15	18	0.72	0.87	0.99	0.99	17	24	
M <sub>3</sub>	2065	2096	17	14	0.82	0.67	0.98	1.00	35	4	
Σ M <sup>B*, A**</sup>	Σ 6234	Σ 6260	Σ 56	Σ 44	0.90 ± 0.22	0.70 ± 0.15	0.99 ± 0.01	0.99 ± 0.00	Σ 66	Σ 40	
Σ M and F	Σ 12480	Σ 12525	Σ 123 <sup>AB</sup>	Σ 99 <sup>AB</sup>	0.99 ± 0.12 <sup>AB'</sup>	0.79 ± 0.12 <sup>AB'</sup>	0.99 ± 0.00 <sup>A''</sup>	0.99 ± 0.00 <sup>A''</sup>	Σ 120	Σ 75	

Means with different letters are significantly different (p &lt; 0.05). M: Male. F: Female.

(1.5-2 g.kg<sup>-1</sup>), regardless of sex, but varied between time intervals. Thus, the PCE/NCE ratio at higher doses was significantly lower than observed in positive the controls treated with NEU. These results suggest the absence of systemic toxicity at GEZJ doses of 0.5-1 g.kg<sup>-1</sup> and moderate toxicity at doses of 1.5-2 g.kg<sup>-1</sup>, regardless of mouse gen-

der, with variable responses over time (24-48 h). Whereas treatment with GEZJ (2 g.kg<sup>-1</sup>) + DXR (5 mg.kg<sup>-1</sup>) significantly reduced the MNPCEs (*n* and %), there was a significant increase in the PCE/NCE ratio with this same treatment, indicating that this combination was not toxic to mouse bone marrow. These results also suggest that the



**Figure 2** - Box-plots showing the MNPCE frequencies and PCE/NCE ratios in mouse bone marrow in mutagenic and antimutagenic assays of *Z. joazeiro* bark. Means with different letters are significantly different ( $p < 0.05$ ). NaCl - control group treated with 150 mM NaCl, NEU - N-nitroso-N-ethylurea (50 mg.kg<sup>-1</sup>), DXR - doxorubicin hydrochloride (5 mg.kg<sup>-1</sup>), GEZJ - Glycolic extract of *Z. joazeiro* Mart. bark (0.5-2 g.kg<sup>-1</sup>), GEZJ (2 g.kg<sup>-1</sup>) + NEU (50 mg.kg<sup>-1</sup>) and GEZJ (2 g.kg<sup>-1</sup>) + DXR (5 mg.kg<sup>-1</sup>).



phytochemical compounds responsible for the moderate toxicity (altered PCE/NCE ratio) of GEZJ (2 g.kg<sup>-1</sup>) in bone marrow may also have an important role in attenuating the mutagenicity (*n* and % of MNPCE) of DRX (5 mg.kg<sup>-1</sup>).

The acute toxicity of different plant extracts, including *Z. joazeiro*, has previously been based on doses (1 to 4 or 5 g/kg) administered orally to different groups of mice (one dose per mouse, with each group containing eight animals: four males and four females) (Alviano *et al.*, 2008). Behavioral parameters, including convulsion, hyperactivity, sedation, grooming, loss of righting reflex, increased or decreased respiration, and changes in food and water intake were also noted. These animals were observed and weighed over a period of 14 days; no weight loss was detected. Treated mice showed no behavioral alterations and the extract LD<sub>50</sub> values ranged from 2.0-3.5 g/kg. None of the extracts was lethal to mice at the doses tested and the data from the *in vivo* assays indicated that the extracts had low toxicity (Alviano *et al.*, 2008). The data from the MN assays presented here provides additional information on the systemic toxicity of *Z. joazeiro* in mouse bone marrow based on the PCE/NCE ratio that suggested moderate toxicity of GEZJ at doses of 1.5-2 g.kg<sup>-1</sup> that was independent of mouse gender but varied with time (24-48 h).

The PCE/NCE ratio is an indicator of the acceleration or inhibition of erythropoiesis and varies with the scoring interval. A continuous decline in the PCE/NCE ratio may reflect the inhibition of cell division, the killing of erythroblasts, the removal of damaged cells, or dilution of the existing cell pool with newly formed cells (Venkatesh *et al.*, 2007). Several mechanisms may contribute to the cytotoxicity of DXR and MN induction (Gewirtz, 1999), including the intercalation of DXR in cellular DNA (Painter, 1978; Kiyomiya *et al.*, 2001), stabilization of the topoisomerase II-DNA complex (Pommier *et al.*, 1985; Guano *et al.*, 1999), free radical-mediated toxicity caused by redox cycling of the semiquinone radical (Bachur *et al.*, 1979), or the formation of reactive oxygen species by the DXR-iron complex (Eliot *et al.*, 1984; Myers, 1998; Konorev *et al.*, 1999). On the other hand, chemicals such as captopril and desferrioxamine (Al-Harbi, 1993; Al-Shabanah, 1993), β-carotene and vitamins A, C and E (Lu *et al.*, 1996; Gulkac *et al.*, 2004; Costa and Nepomuceno, 2006), thiol N-acetylcysteine, probucol, lovastatin and hydrophilic flavonoids such as rutin and luteolin (Al-Gharably, 1996; Sadzuka *et al.*, 1997; D'Agostini *et al.*, 1998; Bardeleben *et al.*, 2002) can also reduce DXR-induced MN formation, genotoxicity and cytotoxicity. However, proponents of herbal medicine always claim that mixtures are better than pure chemicals because the dozens of biologically active compounds in plants work together to produce a greater effect than any one chemical on its own (Mackenzie, 2001).

Screening for newer pharmacological agents that can protect normal cells against DXR-induced cumulative toxicity is essential. Many plants widely used in traditional

medicine are less toxic than pharmaceutical agents and have recently attracted the attention of researchers around the world. Plants contain many compounds and it is likely that these can provide better protection than a single molecule (Vidhya and Devraj, 1999). The presence of many molecules in plants may be advantageous, as some of them may counteract the toxicity of others so that the net effect may be therapeutically beneficial. For example, the effect of various concentrations (200, 250, 300, 350 and 400 mg/kg body weight) of *Aegle marmelos* on DXR-induced mutagenicity in mouse bone marrow was studied (Venkatesh *et al.*, 2007). Mice treated with different concentrations of DXR (5, 10 or 15 mg.kg<sup>-1</sup> body weight) showed a dose-dependent elevation in the frequency of PCE and NCE in their bone marrow, and this was accompanied by a DXR-mediated dose-dependent decline in the PCE/NCE ratio. In contrast, the treatment of mice with *A. marmelos* orally once a day for five consecutive days before treatment with DXR significantly reduced the frequency of DXR-induced micronuclei and significantly increased the PCE/NCE ratio at all time intervals. This chemoprotective effect may reflect the sum of interactions between different components of this complex mixture. The degree of protection may depend on the individual or collective interaction of components with the genotoxic agent. The plausible mechanisms of action of *A. marmelos* in protecting against DXR-induced damage included the scavenging of O<sub>2</sub><sup>•-</sup>, •OH and other free radicals, an increase in antioxidant status, restoration of topoisomerase II activity and inhibition of the formation of the DXR-iron complex (Venkatesh *et al.*, 2007). More recently, Alves *et al.* (2012) evaluated the genotoxic potential of a hydroalcoholic extract of *Copaifera lansdorffii* Desf. leaves and its influence on the genotoxicity of DXR (MN test) in peripheral blood from Swiss mice. Their findings demonstrated that *C. lansdorffii* Desf. was not genotoxic but that the extract significantly reduced the number of micronuclei in DXR-treated mice. The putative antioxidant activity of one or more of the active compounds of *C. lansdorffii* Desf., including two major flavonoid heterosides (quercitrin and afzelin), may explain the effect of this plant on DXR genotoxicity.

## Conclusions

This study used the MN assay to evaluate the mutagenic (clastogeny and/or aneugeny) and antimutagenic activity of an extract of *Z. joazeiro* bark in mouse bone marrow. The *Z. joazeiro* bark extract was not mutagenic at the doses and time intervals tested, although sex-related variation was observed. The antimutagenic effect (anti-clastogeny and/or anti-aneugeny) of *Z. joazeiro* bark extract against DXR-induced genotoxicity was observed at a high dose of extract (2 g.kg<sup>-1</sup>), but was independent of the duration of treatment and animal sex. Low concentrations of GEZJ (0.5-1 g.kg<sup>-1</sup>) were not toxic, regardless of mouse gender and duration of treatment, whereas moderate toxic-

ity was observed at doses of 1.5-2 g.kg<sup>-1</sup>. Together, these findings indicate that phytochemical compounds in *Z. joazeiro* bark can attenuate DRX-induced mutagenicity and that a high dose of extract (2 g.kg<sup>-1</sup>) showed no toxicity in the conditions tested here.

Other studies on the genotoxicity and mutagenicity of *Z. joazeiro* extracts are needed to characterize the (anti)genotoxic effects and mechanisms, and to determine the potential health risks of this extract in humans. Such investigations will be useful for implementing strategies related to the use of *Z. joazeiro* bark in chemoprevention.

## Acknowledgments

This research was supported by Rede Mineira de Ensaios Toxicológicos e Farmacológicos de Produtos Terapêuticos (REDE MINEIRA TOXIFAR - 2012) and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG).

## References

- Agência Nacional de Vigilância Sanitária (2010) Farmacopéia Brasileira. Fundação Oswaldo Cruz, Brasília, 545 pp.
- Al-Gharably NM (1996) Effect of probucol on the cytological and biochemical changes induced by adriamycin in Swiss albino mice. *Res Commun Mol Pathol Pharmacol* 94:289-303.
- Al-Harbi MM (1993) Effect of captopril on the cytological and biochemical changes induced by adriamycin. *Food Chem Toxicol* 31:209-212.
- Al-Shabanah OA (1993) Inhibition of adriamycin-induced micronuclei by desferrioxamine in Swiss albino mice. *Mutat Res* 301:107-111.
- Alves JM, Munari CC, Neto MABM, Furtado RA, Senedese JM, Bastos JK and Tavares DC (2012) *In vivo* protective effect of *Copaifera langsdorffii* hydroalcoholic extract on micronuclei induction by doxorubicin. *J Appl Toxicol* 33:854-860.
- Alviano WS, Alviano DS, Diniz CG, Antonioli AR, Alviano CS, Faria LM, Carvalho MAR, Souza MMG and Bolognese AM (2008) *In vitro* antioxidant potential of plant extracts and their activities against oral bacteria based on Brazilian folk medicine. *Arch Oral Biol* 53:545-552.
- Ames BN (1983) Dietary carcinogens and anticarcinogens: Oxygen radicals and degenerative diseases. *Science* 221:1256-1264.
- Bachur NR, Gordon SL, Gee MV and Kon HNR (1979) NADPH cytochrome P-450 reductase activation of quinone anticancer agents to free radicals. *Proc Natl Acad Sci USA* 76:954-957.
- Bardeleben RV, Dunkern T, Kaina B and Fritz G (2002) The HMG-CoA reductase inhibitor lovastatin protects cells from the antineoplastic drugs doxorubicin and etoposide. *Int J Mol Med* 10:473-479.
- Bean CL, Armstrong MJ and Galloway SM (1992) Effect of sampling time on chromosome aberration yield for 7 chemicals in Chinese hamster ovary cells. *Mutat Res* 265:31-44.
- Beaver BV, Reed W, Leary S, McKiernan B, Bain F, Schultz R, Bennett BT, Pascoe P, Shull E, Cork LC, *et al.* (2000) Report of the American Veterinary Medical Association panel on euthanasia. *J Am Vet Med Assoc* 218:669-696.
- Braga R (1960) Plantas do Nordeste, Especialmente do Ceará. Centro de Divulgação Universitária, Fortaleza, 540 pp.
- Cartaxo SL, Sousa MMA and Albuquerque UP (2010) Medicinal plants with bioprospecting potential used in semi-arid north-eastern Brazil. *J Ethnopharmacol* 13:326-342.
- Chandrasekaran CV, Sundarajan K, Gupta A, Srikanth HS, Edwin J and Agarwal A (2011) Evaluation of the genotoxic potential of standardized extract of *Glycyrrhiza glabra* (GutGard™). *Regul Toxicol Pharmacol* 61:373-380.
- Chorilli M, Brizante AC, Rodrigues CA and Salgado HRN (2007) Aspectos gerais em sistemas transdérmicos de liberação de fármacos. *Rev Bras Farm* 88:7-13.
- Collaborative Study Group for the Micronucleus Test (CSGMT) (1986). Sex differences in the micronucleus test. *Mutat Res* 172:151-163.
- Costa WF and Nepomuceno JC (2006) Protective effects of a mixture of antioxidant vitamins and minerals on the genotoxicity of doxorubicin in somatic cells of *Drosophila melanogaster*. *Environ Mol Mutagen* 47:18-24.
- Cruz GL (1985) Dicionário das Plantas Úteis do Brasil. Editora Civilização Brasileira S.A., Rio de Janeiro, 405 pp.
- Cruz MCS, Santos AM, Barbosa Jr AM, de Melo DLFM, Alviano CS, Antonioli AR, Alviano DS and Trindade RC (2007) Antifungal activity of Brazilian medicinal plants involved in popular treatment of mycoses. *J Ethnopharmacol* 111:409-412.
- D'Agostini F, Bagnasco M, Giunciuglio D, Albini A and de Flora S (1998) Inhibition by oral N-acetylcysteine of doxorubicin-induced clastogenicity and alopecia, and prevention of primary tumors and lung micrometastases in mice. *Int J Oncol* 3:217-224.
- Delvaeye M, Verovski V, De Neve W and Storme G (1993) DNA breakage, cytotoxicity, drug accumulation and retention in two human ovarian tumor cell lines AZ224 and AZ364 treated with adriamycin, modulated by verapamil. *Anticancer Res* 13:1533-1538.
- Dhawan A, Kayani MA, Parry JM, Parry E and Anderson D (2003) Aneugenic and clastogenic effects of doxorubicin in human lymphocytes. *Mutagenesis* 18:487-490.
- Eliot H, Gianni L and Myers C (1984) Oxidative destruction of DNA by the adriamycin-iron complex. *Biochemistry* 5:928-936.
- Gewirtz DA (1999) A critical evaluation of the mechanisms of action proposed for the antitumor effects of the anthracycline antibiotics adriamycin and daunorubicin. *Biochem Pharmacol* 57:727-741.
- Guano F, Pourquier P, Tinelli S, Binaschi M, Bigioni M, Animati F, Manzini S, Zunino F, Kohlhagen G, Pommier Y, *et al.* (1999) Topoisomerase poisoning activity of novel disaccharide anthracyclines. *Mol Pharmacol* 56:77-84.
- Gulkac MD, Akpınar G, Ustun H and Ozon KA (2004) Effects of vitamin A on doxorubicin-induced chromosomal aberrations in bone marrow cells of rats. *Mutagenesis* 19:231-236.
- Indart A, Viana M, Clapés S, Izquierdo L and Bonet B (2007) Clastogenic and cytotoxic effects of lipid peroxidation products generated in culinary oils submitted to thermal stress. *Food Chem Toxicol* 45:1963-1967.
- Jagetia GC and Aruna R (2000) Correlation between cell survival and micronuclei-induction in HeLa cells treated with adria-

- mycin after exposure to various doses of gamma-radiation. *Toxicol Lett* 115:183-193.
- Jagetia GC and Nayak V (1996) Micronuclei-induction and its correlation to cell survival in HeLa cells treated with different doses of adriamycin. *Cancer Lett* 110:123-128.
- Jagetia GC and Nayak V (2000) Effect of doxorubicin on cell survival and micronuclei formation in HeLa cells exposed to different doses of gamma-radiation. *Strahlenther Onkol* 176:422-428.
- Kiyomiya K, Matsuo S and Kurebe M (2001) Differences in intracellular sites of action of adriamycin in neoplastic and normal differentiated cells. *Cancer Chemother Pharmacol* 47:51-56.
- Konorev EA, Kennedy MC and Kalyanaraman B (1999) Cell-permeable superoxide dismutase and glutathione peroxidase mimetics afford superior protection against doxorubicin-induced cardiotoxicity: The role of reactive oxygen and nitrogen intermediates. *Arch Biochem Biophys* 368:421-428.
- Konstantoupoulou I, Vassilopoulou L, Maviaganitsipido U and Scouras ZG (1992) Insecticidal effects of essential oils. A study of the effects of essential oils extracted from eleven Greek aromatic plants on *Drosophila auraria*. *Experientia* 48:616-619.
- Leal ICR, Santos KRN, Itabaiana Jr I, Antunes OAC, Porzel A, Wessjohann L and Kuster RM (2010) Ceanothane and lupane type triterpenes from *Zizyphus joazeiro* - An anti-staphylococcal evaluation. *Planta Med* 76:47-52.
- Lu HZ, Geng BQ, Zhu YL and Yong DG (1996) Effects of beta-carotene on doxorubicin-induced cardiotoxicity in rats. *Zhongguo Yao Li Xue Bao* 17:317-320.
- Mackenzie D (2001) Complementary medicine, a special report. Swallow it whole. *New Sci* 2292:38-40.
- Mateuca R, Lombaert N, Aka PV, Decordier I and Kirsch-Volders M (2006) Chromosomal changes: Induction, detection methods and applicability in human biomonitoring. *Biochimie* 88:1515-1531.
- Myers C (1998) The role of iron in doxorubicin-induced cardiomyopathy. *Semin Oncol* 25:10-14.
- Nunes PHM, Marinho LC, Nunes MLRL and Soares EO (1987) Antipyretic activity of an aqueous extract of *Zizyphus joazeiro* Mart. (*Rhamnaceae*). *Braz J Med Biol Res* 20:599-601.
- OECD (1997a) Guidelines for the Testing of Chemicals: Bacterial reverse mutation test. Organisation for Economic Cooperation and Development, Paris, Guideline 471, 11 pp.
- OECD (1997b) Guideline for the Testing of Chemicals: *In vitro* mammalian chromosome aberration test. Organisation for Economic Cooperation and Development, Paris, Guideline 473, 14 pp.
- OECD (1997c) Guideline for the Testing of Chemicals: Mammalian erythrocyte micronucleus test. Organisation for Economic Cooperation and Development, Paris, Guideline 474, 10 pp.
- Oliveira AFM, Meirelles ST and Salatino A (2003) Epicuticular waxes from caatinga and cerrado species and their efficiency against water loss. *An Acad Bras Cienc* 75:431-439.
- Painter RB (1978) Inhibition of DNA replicon initiation by 4-nitroquinoline 1-oxide, adriamycin, and ethyleneimine. *Cancer Res* 38:4445-4449.
- Pommier Y, Schwartz RE, Zwelling LA and Kohn KW (1985) Effects of DNA intercalating agents on topoisomerase II induced DNA strand cleavage in isolated mammalian cell nuclei. *Biochemistry* 24:6406-6410.
- Purves D, Harvey C, Tweats D and Lumley CE (1995) Genotoxicity testing: Current practices and strategies used by the pharmaceutical industry. *Mutagenesis* 10:297-312.
- Ribeiro LR (2003) Teste de micronúcleo em medula óssea de roedor *in vivo*. In: Ribeiro LR, Salvadori DMF and Marques EK (eds) *Mutagênese Ambiental*. Ulbra, Canoas, pp 173-198.
- Sadzuka Y, Sugiyama T, Shimoi K, Kinai N and Hirota S (1997) Protective effect of flavonoids on doxorubicin-induced cardiotoxicity. *Toxicol Lett* 92:1-7.
- Schmid W (1976) Chemical mutagens. In: Hollender A (ed) *The Micronucleus Test for Cytogenetic Analysis*. Plenum Press, New York, pp 31-53.
- Schuhly W, Heilmann J, Calis I and Sticher O (1999) New triterpenoids with antibacterial activity from *Zizyphus joazeiro*. *Planta Med* 65:740-743.
- Shan K, Lincoff AM and Young JB (1996) Anthracycline-induced cardiotoxicity. *Ann Inter Med* 125:47-58.
- Silva CR, Vieira PM, Santos SC and Chen-Chen L (2011) Assessment of *Duguetia furfuracea* genotoxic and cytotoxic activity in bacteria and mice. *An Acad Bras Cienc* 84:149-156.
- Tavares W (1996) *Manual de Antibióticos e Quimioterápicos Anti-Infeciosos - Introdução ao Estudo dos Antimicrobianos*. Atheneu, Rio de Janeiro, 792 pp.
- UNESCO (1978) *Comissão de Ética no Uso de Animais, Declaração Universal dos Direitos dos Animais (UNESCO)*, Bruxelas, Bélgica, 2 pp.
- Van Acker SA, Kramer K, Grimbergen JA, Van Den Berg DJ, Van Der Vijgh WJ and Bast A (1995) Monohydroxyethylrutinoside as protector against chronic doxorubicin-induced cardiotoxicity. *Br J Pharmacol* 115:1260-1264.
- Van Acker FA, Van Acker SA, Kramer K, Haenen GR, Bast A and Van Der Vijgh WJ (2000) 7-Monohydroxyethylrutinoside protects against chronic doxorubicin-induced cardiotoxicity when administered only once per week. *Clin Cancer Res* 6:1337-1341.
- Varanda EA (2006) Atividade mutagênica de plantas medicinais. *Rev Ciênc Farm Básica Apl* 27:1-7.
- Venkatesh P, Shantala B, Jagetia GC, Rao KK and Baliga MS (2007) Modulation of doxorubicin-induced genotoxicity by *Aegle marmelos* in mouse bone marrow: A micronucleus study. *Integr Cancer Ther* 6:42-53.
- Vidhya N and Devraj SN (1999) Antioxidant effect of eugenol in rat intestine. *Ind J Exp Biol* 37:1192-1195.
- Zambrano MA, Targa HJ and Rabello-Gay MN (1982) Physiological saline solutions as a useful tool in micronucleus and metaphase slide preparations. *Stain Technol* 57:48-49.

Associate Editor: Daisy Maria Fávero Salvadori